

UNIVERSITY OF PAVIA
DEPARTMENT OF THEORETICAL AND NUCLEAR PHYSICS

Simplicial and modular aspects of string dualities

Advisor:

Prof. Mauro Carfora

Doctoral thesis of:

Valeria L. Gili

Dottorato di Ricerca XVIII Ciclo

Contents

Introduction	7
I Simplicial aspects of string dualities	19
1 Dual uniformizations of triangulated surfaces	21
1.1 Random Regge triangulated surfaces as singular Euclidean structures	21
1.2 Ribbon graphs on Regge polytopes	28
1.3 Dual uniformizations of a random Regge triangulated surface . .	31
2 BCFT on the annuli: Langlands' boundary states	41
2.1 Bulk and Boundary Conformal Field Theory	43
2.1.1 Glueing conditions and boundary fields	46
2.2 The quantum amplitude on the annulus	50
2.3 Generalized Langlands boundary states	57
3 Boundary Insertion Operators	61
3.1 Boundary Insertion Operators	61
3.2 Investigations at the T-duality self dual radius	67
3.3 The algebra of rotated Boundary Insertion Operators	76
3.4 The action of BIOs at the self-dual radius	79
3.5 Open string amplitude on a RRT	81
4 Inclusion of Open String gauge degrees of freedom	83
4.1 Chan-Paton factors on a RRT	84
4.2 From triangulations to strings	87
4.3 Extending the gluing procedure	91
4.3.1 Chiral currents and boundary states	94
4.3.2 Gluing along the Ribbon Graph: a proposal	95
4.4 Conclusions and perspectives	96

II	Cosmological backgrounds in String Theory	99
5	Geodesics on maximally non compact cosets U/H	101
5.1	Decoupling the sigma model from gravity	102
5.2	Geodesic equations in target space and the Nomizu operator . . .	104
6	The A_2 toy model as a paradigm	111
6.1	Structure of the A_2 system	112
6.2	Scalar fields of the A_2 model	116
6.3	Generation of new solutions via the compensator method	117
6.3.1	Solution of the differential equations for the tangent vectors with two Cartan and one nilpotent field	
6.3.2	Solution of the differential equations for the tangent vectors with two Cartan and three nilpotent fields	
7	The E_8 Lie algebra	123
8	Oxidation of the A_2 solutions	131
8.1	Possible embeddings of the A_2 algebra	131
8.2	Choice of one embedding example	133
8.2.1	Oxidation of the solution with only one root switched on	136
8.2.2	Oxidation of the solution with all three roots switched on	138
8.2.3	How the supergravity field equations are satisfied and their cosmological interpretation	142
8.2.4	Properties of the solution with just one root switched on .	144
8.2.5	Properties of the solution with all roots switched on . . .	148
8.2.6	Summarizing the above discussion and the cosmological billiard	152
8.3	Conclusions and perspectives	154
III	Appendices	159
A	Projection of generalized Langland's boundary states	161
B	Conformal properties of Boundary Insertion Operators	165
C	An introduction to truly marginal deformations of boundary conformal field theories	169
C.1	Deformations of a boundary conformal field theory	171
C.1.1	The general prescription	172
C.1.2	Truly marginal operators: the case of chiral fields	174
D	Useful formulae	177
D.1	Simplicial String Duality	177
D.2	Cosmological solution of supergravity	178
E	Listing of the E_8 positive roots	181
E.1	Listing according to height	181
E.2	Listing the roots according to the dimensional filtration	184

Introduction

My Ph.D. work has been dedicated to the study of different aspects of modern unifying theories.

Part of the time has been devoted to search for cosmological backgrounds in superstring theories, thus continuing the research line started in my Laurea thesis.

This work, which has been done in collaboration with Prof. Pietro Fré's group, fits into a wide set of research lines merging string theory, which try to express all fundamental constituents of the universe and their interaction into a unique framework, with the undertaken of cosmological models, whose aim is to describe the origin, the evolution and the structure of the Universe as a whole. The great number of success of both models has been suggesting since middle of 90's a possible unification of their investigation areas. In view of the new observational data which seems to provide evidence for a small but positive cosmological constant [1, 2, 3, 4], there has been wide interest in the context of M-theory/string theory and extended supergravities for the search of de Sitter like vacua (see for instance [5, 6, 7, 8, 9] and references therein). More generally, the analysis of time-dependent backgrounds [10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26] has been done in various approaches and at different levels, namely both from the microscopic viewpoint, considering time-dependent boundary states and boundary CFTs (see for instance [27, 28] and references therein) and from the macroscopic viewpoint studying cosmological solutions in both gauged and ungauged supergravities.

Successively, in collaboration with Prof. Mauro Carfora's group, I started investigating modular and simplicial properties of string dualities. Both collaborations have been characterized by a deep analysis of dynamic symmetries associated with Lie algebras of both finite and affine kind.

The first hint of role Kač-Moody algebras plays in describing dynamical symmetries of gravitational models goes back to early '70, when Geroch showed that the automorphism group of solutions of Einstein Gravity reduced to $D = 2$ is infinite dimensional and it has as underlying Lie algebra A_1 , the untwisted affine Kač-Moody extension of A_1 [29, 30], this being the three dimensional Lie algebra of $SL(2, \mathbb{R})$. This discovery triggered a lot of interest in the analysis of

these symmetries, but it was with the advent of string theory and supergravity that the interest in infinite symmetries received a boost. As a matter of fact, by dimensionally reducing $D = 10$ or $D = 11$ supergravity to lower dimensions, new dynamical symmetries arise. The so called hidden symmetries act as isometries of the metric on the scalar manifold and as generalized electric/magnetic duality rotations on the various p -forms. Formalization of these concepts goes back to the work of Cremmer and Julia, who clarified that $E_{11-D(11-D)}$ is the duality symmetry group of maximal supergravity in $D \geq 3$ dimensions, obtained by a Kaluza-Klein compactification of M -theory on a T^{11-D} torus. In particular, they showed that the massless scalars which emerge from the Kaluza-Klein mechanism in D dimensions, just parametrize the maximally non-compact coset manifold

$$\mathcal{M}_{scalar} = \frac{E_{11-D(11-D)}}{H_{11-D}}$$

where $H_{(11-D)}$ is the maximally compact subgroup of the simple Lie group $E_{11-D(11-D)}$ [31, 32].

Julia noted that the extension of this process to $D < 3$ calls into play algebras which are no longer finite dimensional, but rather they are infinite dimensional Kač-Moody ones[33]. As a matter of fact, $E_{9(9)}$ is the affine extension of $E_{8(8)}$, while $E_{10(10)}$ is its double hyperbolic extension. This last case is particularly intriguing. Since compactification and truncation to the massless modes is an alternative way of saying that we just focus on field configurations THAT DEPEND ONLY THE REMAINING D COORDINATES, the compactification on a T^{10} torus leads to non trivial dependence of supergravity fields just on *one coordinate*, namely *time*, thus linking the arising of cosmological backgrounds to the E_{10} algebra.

In this connection, a very much appealing and intriguing scenario has been proposed in a series of papers by Damour et al. [34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44]: that of cosmological billiards. Studying the asymptotic behavior of ten (eleven) dimensional supergravity field equations near time (space-like) singularities, these authors envisaged the possibility that the nine (ten) cosmological scale factors relative to the different space dimensions of string theory plus the dilaton could be assimilated to the Lagrangian coordinates of a fictitious ball moving in a ten(resp. eleven)-dimensional space.

In their approach, Damour and collaborators analyzed the cosmic billiard phenomenon as an asymptotic regime in the neighborhood of space-like singularities. The billiard walls were seen as the various p -forms of supergravity acting as exponential potential walls which become sharper and sharper in the evolution toward the singularity, ending in infinite potential walls in the $t \rightarrow 0^+$ limit. The main focus attention was centered on establishing whether and under which conditions there may be a chaotic behavior in the evolutions of scale factors of the universe. Eventually, the authors established that the billiard




Figure 1: Time-dependent homogeneous supergravity backgrounds in $D = 10$ can be obtained by first dimensionally reducing to $D = 3$, solving the differential equations of the sigma model and then oxidizing back the result to $D = 10$. This procedure defines also the action of the hidden symmetry group $E_{8(8)}$ on the ten dimensional configurations.

table to be hyperbolic was a necessary condition, where the billiard table was identified with the Weyl chamber of the $E_{(10)}$ algebra.

There is a clear relation between this picture and the duality groups of superstring theories. As a matter of fact, the Cartan generators of the $E_{r(r)}$ algebra are dual to the *radii* of the T^{r-1} torus plus the dilaton. So it is no surprising that the evolution of the cosmological scale factor should indeed represent some kind of motion in the dual of the Cartan subalgebra of E_{10} . Although naturally motivated, the E_{10} billiard picture was so far considered only in the framework of an approximated asymptotic analysis and no exact solution with such a behavior was actually constructed.

In our work, we investigated the problem from a different viewpoint. We focused on three-dimensional maximal supergravity [45]–[46], namely on the dimensional reduction of type II theories on a T^7 torus, instead of going all the way down to reduction to one-dimension. The advantage of this choice is that all the bosonic fields are already scalar fields, described by a non-linear sigma model without, however, the need of considering Kač-Moody algebras which

arise as isometry algebras of scalar manifolds in $D < 3$ space-times. In this way we had been able to utilize the *solvable Lie algebra approach* to the description of the whole bosonic sector which enabled us to give a completely algebraic characterization of the microscopic origin of the various degrees of freedom [47, 48]. Within this framework the supergravity field equations for bosonic fields restricted to only time dependence reduce simply to the geodesic equations in the target manifold $E_{8(8)}/SO(16)$. These latter can be further simplified to a set of differential equations whose structure is completely determined in Lie algebra terms. This was done through the use of the so called *Nomizu operator*. The concept of Nomizu operator coincides with the concept of covariant derivative for solvable group manifolds. The possibility of writing covariant derivatives in this algebraic way as linear operators on solvable algebras relies on the theorem that states that a non-compact coset manifold with a transitive solvable group of isometries is isometrical to the solvable group itself.

The underlying idea for our approach was rooted in the concept of hidden symmetries. Cosmological backgrounds of superstring theory, being effectively one-dimensional fill orbits under the action of a very large symmetry group, possibly E_{10} , that necessarily contains $E_{8(8)}$, as the manifest subgroup in three dimensions. Neither E_{10} nor $E_{8(8)}$ are manifest in 10-dimensions but become manifest in lower dimension. So an efficient approach to finding spatially homogeneous solutions in ten dimensions consists of the process schematically described in fig.1. First one reduces to $D = 3$, then solves the geodesic equations in the algebraic setup provided by the Nomizu-operator-formalism and then oxidizes back the result to a full fledged $D = 10$ configuration. Each possible $D = 3$ solution is characterized by a non-compact subalgebra

$$\mathbf{G} \subset E_{8(8)}$$

which defines the smallest consistent truncation of the full supergravity theory within which the considered solution can be described. The inverse process of oxidation is not unique but leads to as many physically different ten dimensional solutions as there are algebraically inequivalent ways of embedding $\mathbf{G}$ into $E_{8(8)}$. In [49] we illustrated this procedure by choosing for $\mathbf{G}$ the smallest non abelian rank two algebra, namely $\mathbf{G} = A_2$. The solvable Lie algebra formalism allowed us to control, through the choice of the $\mathbf{G}$ -embedding, the physical ten-dimensional interpretation of any given σ -model solution. Focusing on a particular embedding for the subalgebra A_2 , we got a type IIB time dependent background generated by a system of two euclidean D-branes or S-branes [10, 11, 12, 13, 14, 15, 16, 17, 18]: a D3 and a D1, whose world volumes are respectively four and two dimensional. This physical system contains also an essential non trivial B-field reflecting the three positive root structure of the A_2 Lie algebra, one root being associated with the RR 2-form $C^{[2]}$, a second with the RR 4-form $C^{[4]}$ and the last with the NS 2-form $B^{[2]}$. In the time evolution




Figure 2: A ribbon graph with $g = 1$ and $h = 9$ and the associated Riemann surface, in which holes are filled with regions having the topology of a disk

of this exact solution of type II B supergravity we retrieved a smooth realization of the bouncing phenomenon envisaged by the *cosmic billiards* of [35]–[44]. Indeed, in the oxidized picture, the scale factors corresponding to the dimensions parallel to the S-branes first expand and then, after reaching a maximum, contract. The reverse happens to the dimensions transverse to the S-branes. They display a minimum approximately at the same time when the parallel ones are maximal. Transformed to the dual CSA space this is the bouncing of the cosmic ball on a Weyl chamber wall. This is not the full cosmic billiard, but it illustrates the essential physical phenomena underlying its implementation.

In the meanwhile I started, in collaboration with Prof. Mauro Carfora and Dr. Claudio Dappiaggi, to investigate simplicial and modular aspects of string dualities.

Since 't Hooft seminal paper[50], the idea that a large N gauge theory has a dual description in term of a closed string theory has been drawing the attention of large part of theoretical physicists community. According to 't Hooft prescription, if we consider a $U(N)$ gauge theory whose action is written as $\frac{1}{g_{YM}^2} \int \mathcal{L}(A)$ (A is the gauge connection), the group theoretical part of Feynman diagrams arises as a ribbon graph, whose edges acquire a gauge coloring in the first fundamental times first anti-fundamental representation of the gauge group. In this connection, amplitudes and Green functions are obtained summing over all possible (planar and non-planar) diagrams, with the appropriate combinatorial factor. Ribbon graphs associated to vacuum amplitudes can be classified according to their dependence from N and g_{YM} . As a matter of fact, if we consider a ribbon graph with h boundary components, it comes with the factor $(g_{YM}^2)^{-V+E} N^h$, where V and E are respectively the number of vertexes and edges of the graph. Since each ribbon graph can be viewed as a closed Riemann surface with h holes, the dependence of the amplitude from N and g_{YM} can be captured by the topology of the underlying surface. If we consider an associated Riemann surface of genus g , we can write:

$$(g_{YM}^2)^{-V+E} N^h = (g_{YM}^{2g-2}) (g_{YM}^2 N)^h = (g_{YM}^{2g-2}) \lambda^h \quad (1)$$

where we have exploited the topological relation $V - E + h = 2 - 2g$, and

where we have introduced the 't Hooft coupling $\lambda = g_{YM}^2 N$. Thus, the sum over all ribbon graphs remarkably translates into a sum over all topologies. The associated amplitude is then given by:

$$\mathcal{F} = \sum_{g=0}^{\infty} \sum_{h=1}^{\infty} (g_{YM}^{2g-2}) \lambda^h \mathcal{F}_{g,h} = \mathcal{F} = \sum_{g=0}^{\infty} (g_{YM}^{2g-2}) \mathcal{F}_g, \quad (2)$$

where we have defined

$$\mathcal{F}_g = \sum_{h=1}^{\infty} \lambda^h \mathcal{F}_{g,h} \quad (3)$$

and the coefficients $\mathcal{F}_{g,h}$ being functions of other parameters of the theory. 't Hooft's conjecture states that, since at fixed λ formula (3) can be seen both as an $g_{YM} \rightarrow 0$ or as a $N \rightarrow \infty$ expansion, in the large N (and fixed λ) limit an equivalent description arises, involving closed Riemann surfaces which are obtained "filling holes of the ribbon graphs with disks". This naturally promotes the expansion (2) to a sum over genus, *i.e.* a closed string expansion.

Furthermore, at the beginning of 90s', Susskind[51] and 't Hooft[52] himself introduced the HOLOGRAPHIC PRINCIPLE, according to which classical space-time geometry and its matter contents arise from an underling (non a priori gravitational) quantum theory in such a way that the covariant entropy bound should be satisfied.

In this connection, Maldacena's conjecture about *AdS/CFT* correspondence [53] had merged elegantly these two aspects. It is perhaps the best example of such a kind of holographic correspondence, in which the supergravity limit of a string theory in d dimensions admits an holographically dual description in term of a gauge theory living on a codimension one manifold, and where the connection between the gauge theory in the open string sector and the closed string theory is guided by a large N transition. Not less important, it provides a powerful tool in studying non-perturbative effects of a quantum field theory by means of perturbative techniques proper of string theory.

However, dealing with *AdS/CFT* correspondence, a strong working assumption is usually performed. The large N transition is, a priori, a worldsheet effect: as remarked above, the leading (gauge theory) large N correlator functions arise from planar open string diagrams with some insertion on their boundaries. In the expected worldsheet scenario, the loops of the original conformal field theory glue up to form a closed Riemann surface with n closed string insertions on a different background. Thus, the open-closed string equivalence works at the level of the worldsheet and it has to be implemented at a worldsheet moduli space level, though the very process is unknown.

On the contrary, in a topological setup, even if the duality is guided by a geometrical transition at target space level (for a review see [54, 55]), the duality process acting at a worldsheet level was identified. As a matter of fact, in their

original definition, topological large N open/closed string duality are based on the gauge theory description of topological string theory. In particular, when the A topological string model is defined on a Calabi-Yau $X = T^*M$, M being a three-manifold, and there are N topological D -branes wrapping on M , the full Open String Field Theory is equivalent to a $U(N)$ Chern-Simons on M . The open string amplitudes are then just numbers computed by the fatgraphs of the corresponding gauge theories. In this background, the large N transition act at a geometric level as a conifold transition [55] relating the open string Calabi-Yau background X underlying the gauge theory and the closed string Calabi-Yau background Y . As usual, at worldsheet level, the scenario one expects is that the boundary of the open Riemann surface get glued up to form a closed surface with n close string insertion. In this connection, at worldsheet level, when the 't Hooft coupling is small, a new branch of the conformal theory arises and the compact domains of this new phase can be viewed as holes for an open string theory living in a different regime. In such a context, assumptions are no longer necessary. However, the theory is topological and it excludes a dynamical coupling with gravity in the target space. As pointed out, open string amplitudes are just numbers which the duality process rereads in term of the topological invariants of the closed surface.

In this connection, a paradigmatical result has been recently established by Gopakumar [56, 57] in the setup of AdS/CFT correspondence. His intuition has as starting point $\mathcal{N} = 4$ SYM theory, namely the $\alpha' \rightarrow 0$ limit of open string theory. Starting from a Schwinger parametrization of free fields correlators, the author was able to reorganize n points amplitudes (at genus g in a 't Hooft sense) in terms of skeleton diagrams. These are simply the graphs obtained by merging together all the homotopically equivalent contractions between any two pairs of vertexes. In this connection, the generic skeleton diagram associated to a particular merging of contractions is simply a triangulation of the underlying genus g surface, with as many vertexes there are internal and external ones. As a consequence, the entire expansion for the n -points function can be completely expressed as a sum over all the inequivalently connected skeleton graphs contributing to the amplitude.

The counting over degrees of freedoms allowed him to interpret the integral over Schwinger times plus the sum over inequivalent skeleton graphs as sum over the decorated moduli space $\mathcal{M}_{g,n} \times \mathbb{R}_+^n$. Thus, the arising of a closed string moduli space, was claimed as the manifestation that a glueing process might be seen as a change of variables in the integrand of amplitudes.

This transition was actually evident in a target space connection: the author introduced an algorithm which, exploiting the *bulk-to-boundary* propagator introduced by Witten, Gubser, Klebanov and Polyakov in [58, 59], allows to recast gauge amplitudes as amplitudes in *AdS*[57].

In view of these results, in order to analyze string dualities in a general framework, it seems that a discretized approach can play a fundamental role, in particular dealing with triangulations with varying connectivity. Thus, motivated by the ubiquitous role that simplicial methods play in the above results, in collaboration with Prof. Mauro Carfora and Dr. Claudio Dappiaggi, we have tried to implement examples of open/closed string duality in a recently introduced new geometrical framework[60].

Our aim is to extend the large N process to the appealing context of dynamical background fields, while retaining the *open-to-closed* transition to be implemented at a worldsheet level.

This approach is based on a careful use of uniformization theory for triangulated surfaces carrying curvature degrees of freedom. Starting from a closed string theory point of view i.e. an n -marked Riemann surface with conical singularities, we can switch through geometrical arguments to a open string theory point of view where the "cones" are traded with finite cylindrical ends. This transition between the open and the closed Riemann surface exploits a particular connection between the N_0 localized curvature degrees of freedom in the closed sector and the modular data associated to finite cylindrical ends in the open sector[60].

In order to show how this uniformization arises, let us consider the dual polytope associated with a Random Regge Triangulation Triangulation [61] $|T_l| \rightarrow M$ of a Riemannian manifold M . The singular Euclidean structure around each puncture can be uniformized by a conformal class of conical metrics (see formula (1.10)), while the edge-refinement of the Regge dual polytope 1-skeleton is in one-to-one correspondence with trivalent Ribbon graphs.

Using properties of Jenkins-Strebel quadratic differentials [60], it is possible to fix a point in the conformal class decorating the neighborhood of each curvature supporting vertex, uniformizing it with a punctured disk endowed with a conical metric

$$ds_{(k)}^2 \doteq \frac{[L(k)]^2}{4\pi^2} |\zeta(k)|^{-2(\frac{\varepsilon(k)}{2\pi})} |d\zeta(k)|^2.$$

Moreover, exploiting the Stebel theorem[62], it is possible to introduce an uniformizing coordinate both along the edges of the Ribbon Graph and on each trivalent vertex.

Alternatively, we can blow up every cone into a corresponding finite cylindrical end, by introducing a finite annulus

$$\Delta_{\varepsilon(k)}^* \doteq \left\{ \zeta(k) \in \mathbb{C} \mid \exp -\frac{2\pi}{2\pi - \varepsilon(k)} \leq |\zeta(k)| \leq 1 \right\}$$

endowed with the cylindrical metric:

$$|\phi(k)| \doteq \frac{[L(k)]^2}{4\pi^2} |\zeta(k)|^{-2} |d\zeta(k)|^2$$

It is important to stress the different role that the deficit angle plays in such two uniformization. In the “closed” uniformization the deficit angles $\varepsilon(k)$ plays the usual role of localized curvature degrees of freedom and, together with the perimeter of the polytopal cells, provide the geometrical information of the underlying triangulation. Conversely, in the “open” uniformization, the deficit angle associated with the k -th polytope cell defines the geometric moduli of the k -th cylindrical end. As a matter of fact each annulus can be mapped into a cylinder of circumference $L(k)$ and height $\frac{L(k)}{2\pi-\varepsilon(k)}$, thus $\frac{1}{2\pi-\varepsilon(k)}$ is the geometrical moduli of the cylinder. This shows how the uniformization process works quite differently from the one used in Kontsevich-Witten models, in which the whole punctured disk is uniformized with a cylindrical metric. In this case the disk can be mapped into a semi-infinite cylinder, no role is played by the deficit angle and the model is topological; conversely, in our case, we are able to deal with a non topological theory.

Very recently[63] Gopakumar pursued further the identification of the field theory expression for the integrand over the Schwinger parameters and with a correlator of closed string vertex operator. In his construction, he exploited the isomorphism between the space of metric Ribbon graphs and $\mathcal{M}_{g,n} \times \mathbb{R}_+^n$ to associate to each skeleton diagram naturally coupled to the Schwinger parametrization of gauge amplitudes the unique dual Ribbon graph. To this end, he proposed a precise dictionary between Strebel lengths l_r (*i.e.* lengths of ribbon graph edges) and Schwinger times τ_r , namely:

$$l_r = \frac{1}{\tau_r} \quad (4)$$

With this dictionary in mind, exploiting the Strebel theorem, which uniquely associate a closed Riemann surface uniformization to a given ribbon graph, he gave a concrete proposal to reconstruct a particular closed Riemann worldsheet associated to a given gauge amplitude.

We would like to remark differences between this approach and our construction. Gopakumar starts with gauge theory correlator, and, identifying Schwinger times with the inverse of Strebel lengths, he is able to associate to each correlator a dual closed surface. On the contrary, in our construction, Strebel theorem allows to introduce a suitable uniformization of both the open and the closed surfaces, while the transition between the two geometries is deeply rooted into the discrete structure of the surface. It is obtained exploiting conformal properties of the singular Euclidean metric around each vertex in order to trade curvature assignment of the closed surface into moduli of the open surface M_∂ , which thus inherits a precise discrete structure. As a final result, in the open sector the overall picture sees the decomposition of the Riemann surface into its fundamental cylindrical components.

In our picture, each cylindrical end can be interpreted as an open string

connected at one boundary to the ribbon graph associated to the discretized worldsheet and, at the other boundary, to a D-brane which acts naturally as a source for gauge fields. This provides a new kinematical set-up for discussing gauge/gravity correspondence.

While, at fixed genus g and number of vertexes N_0 , in the closed sector both the coupling of the geometry of the triangulation with D bosonic fields and the quantization of the theory can be performed under the paradigm of critical field theory, the coupling of the two-dimensional open Riemann surface M_∂ in formula (1.48) with a quantum bosonic open string has called into play Boundary Conformal Field Theory (BCFT) techniques. We have discussed the quantization of bosonic fields on each finite cylindrical end, then we have glued together the resulting BCFTs along the intersection pattern defined by the ribbon graph naturally associated to the Regge polytope dual to the original triangulation[64, 65].

In particular, the unwrapping of the cones into finite cylinders has suggested to compactify each field defined on the k -th cylindrical end along a circle of radius $\Omega(k) = \frac{R(k)}{l(k)}$:

$$X^\alpha(k) \xrightarrow{\vartheta(k) \rightarrow \vartheta(k) + 2\pi} X^\alpha(k) + 2\pi\nu^\alpha(k) \frac{R^\alpha(k)}{l(k)} \quad \nu(k) \in \mathbb{Z}$$

where $l(k)$ is an unspecified length defined as a function of all characteristic scales defining the geometry of the underlying triangulation.

Under these assumptions, it is possible to quantize the theory and to compute the quantum amplitude over each cylindrical end: writing it as an amplitude between an initial and final state, we can extract suitable boundary states which arise as a generalization of the states introduced by Langlands in [66]. As they stand, these boundary state did not preserve neither the conformal symmetry nor the $U(1)_L \times U(1)_R$ symmetry generated by the cylindrical geometry. It has then been necessary to impose on them suitable gluing conditions relating the holomorphic and anti-holomorphic generators on the boundary. These restrictions generated the usual families of Neumann and Dirichlet boundary states.

Within this framework, the next step in the quantization of the theory has been to define the correct interaction of the N_0 copies of the cylindrical CFT on the ribbon graph associated with the underlying Regge Polytope. This has been achieved via the introduction over each strip of the graph of Boundary Insertion Operators (BIO) $\psi_{\lambda(p,q)}^{\lambda(p)\lambda(q)}$ which act as a coordinate dependent homomorphism from $V_{\lambda(p)} \star V_{\lambda(p,q)}$ and $V_{\lambda(q)}$, so mediating the changing in boundary conditions. Here $V_{\lambda(\bullet)}$ denotes the Verma module generated by the action of the Virasoro generators over the $\lambda(\bullet)$ highest weight and $\star$ denotes the fusion of the two representations.

In the limit in which the theory is rational (*i.e.* when the compactification ra-

dius is an integer multiple of the self dual radius $\Omega_{s.d.} = \sqrt{2}$) the Hilbert space of the BCFT can be rewritten as an $SU(2)_{k=1}$ WZW model Hilbert space. It is then possible to identify BIO as marginal deformations of boundary conditions changing operators. Thus they are primary operators with well defined conformal dimension and correlators. Moreover, considering the coordinates of three points in the neighborhood of a generic vertex of the ribbon graph, we can write the OPEs describing the insertion of such operators in each vertex. Considering four adjacent boundary components, it has then been possible to show that the OPE coefficients $C_{j_{(r,p)}j_{(q,r)}j_{(p,q)}}^{j_p j_r j_q}$ are provided by the fusion matrices $F_{j_r j_{(p,q)}} \begin{bmatrix} j_p & j_q \\ j_{(r,p)} & j_{(q,r)} \end{bmatrix}$, which in WZW models coincide with the $6j$ -symbols of the quantum group $SU(2)_{e^{\frac{\pi}{3}i}}$:

$$C_{j_{(r,p)}j_{(q,r)}j_{(p,q)}}^{j_p j_r j_q} = \left\{ \begin{matrix} j_{(r,p)} & j_p & j_r \\ j_q & j_{(q,r)} & j_{(p,q)} \end{matrix} \right\}_{Q=e^{\frac{\pi}{3}i}}$$

From these data, through edge-vertex factorization we have characterized the general structure of the partition function for this model [67] as a sum over all possible $SU(2)$ primary quantum numbers describing the propagation of the Virasoro modes along the N_0 cylinders $\{\Delta_{\varepsilon(k)}^*\}$.

The emerging overall picture is that of N_0 cylindrical ends glued through their inner boundaries to the ribbon graph, while their outer boundaries lay on D-branes. Each D-brane acts naturally as a source for gauge fields: it has allowed us to introduce open string degrees of freedom whose information is traded through the cylinder to the ribbon graph, whose edges thus acquire naturally a gauge coloring. This provides a new kinematical set-up for discussing gauge/gravity correspondence.

Outline

In writing this thesis, I have separated the two different topics in two distinct parts.

The first part has been dedicated to the investigation of simplicial and modular aspects of string dualities.

Chapter 1 contains an introduction to the peculiar geometry arising when we uniformize Riemann surfaces carrying curvature degrees of freedom. After a short presentation of fundamental concepts of simplicial geometry, I have reviewed the two dual uniformizations of a RRT cited above, thus summarizing results in [60, 61].

With chapter 2 it starts the description of the coupling between non-critical Polyakov string with the geometry defined by the dual uniformized open Riemann surface ∂M . In particular, this chapter is devoted to the quantization of the BCFT arising over each cylindrical end $\Delta_{\varepsilon(k)}^*$.

Chapter 3 introduces Boundary Insertion Operators and their conformal properties, laying the basis of the description of interaction of the $N_T(0)$ dis-

tinct copies of the BCFT along the pattern defined by the ribbon graph Γ . This project is completed in chapter C, where we will show that, at enhanced symmetry point, we are able to coherently describe the dynamical glueing calling into play truly marginal deformation of the original boundary conformal field theory.

Eventually, chapter 4 is devoted to the analysis of the coupling of the model with background gauge fields.

Part II reports results obtained in [49] in searching for cosmological background of superstring theories. Chapter 5, after summarizing the Kaluza-Klein reduction of the bosonic sector of type *IIA/B* superstring theory on T^7 , introduces a particular ansatz for the background metric in $D = 3$, allowing the decoupling on the scalar sigma model from gravity. In particular, it shows how to rephrase the supergravity scalar fields equations of motions as geodesic equations on $\mathcal{M}_{128}$. Moreover, it presents the compensator method which allows, once we have found a solution of the geodesic problem via the solvable Lie algebra parametrization of the above coset manifold, to generate new solutions exploiting isometries of the model.

In chapter 6 we have applied the compensator method to the simplest (non-trivial) coset manifold $\mathcal{M}_5 = \exp[SolvA_2]$, while in chapter 7 shows that the hierarchical dimensional reduction/oxidation of supergravity backgrounds is algebraically encoded in the hierarchical embeddings of subalgebras into E_8 algebra. Thus, in chapter 8, after analyzing possible regular embeddings $A_2 \hookrightarrow E_{8(8)}$, promoting the solution found with the compensator method to solutions of type *IIA/B* supergravities, we have applied the oxidation process to derive the $D = 10$ supergravity cosmological backgrounds associated to the previous solutions.

Part I

SIMPLICIAL ASPECTS OF STRING DUALITIES

Chapter 1

Dual uniformizations of triangulated surfaces

In this chapter, after a short introduction to the concepts of triangulation of a Riemann surface and of random Regge triangulations (an exhaustive introduction to simplicial geometry can be found in [68], while a statistical field theory approach to discrete geometry can be found in [69]), we will review explicitly the geometrical framework arising when we uniformize triangulated surfaces carrying curvature degrees of freedom. This framework has been developed in [60, 61] and we refer to these papers and to references therein for a deeper analysis.

1.1 Random Regge triangulated surfaces as singular Euclidean structures

One way to define the topological properties of a surface is to construct a polyhedron homeomorphic to it. Then we will be able to define the Euler characteristic and the homological properties of the given surface via the properties of the associated polyhedron. The main procedure, then, is to associate to each surface a collection of standard object (triangles in two dimensions, simplexes in higher ones) in such a way that it will become possible to associate to each surface a standard abelian structure.

Simplexes are defined as the building blocks of polyhedron. An r -simplex $\langle p_0 p_1 \dots p_r \rangle$ is an r -dimensional object whose vertexes p_i are geometrically independent, that is, no $(r-1)$ -dimensional hyperplane contains all the $r+1$ points. Let $p_0, \dots, p_r$ be geometrically independent points in $\mathbb{R}^m$, with $m \geq r$. The r -simplex $\sigma_r = \langle p_0 p_1 \dots p_r \rangle$ is expressed as:

$$\sigma_r = \left\{ x \in \mathbb{R}^m \mid x = \sum_{i=0}^r c_i p_i \geq 0, \sum_{i=0}^r c_i = 1 \right\}$$

where the $(c_1, \dots, c_r)$ are the barycentric coordinates of x . Since σ_r is a bounded closed subset of $\mathbb{R}^m$, it is compact.

Let q be an integer with $0 \leq q \leq r$. If we choose $q + 1$ points $p_{i_0}, \dots, p_{i_q}$ out of $p_0, \dots, p_r$, these $q + 1$ points define a q -simplex $\sigma_q = \langle p_{i_0}, \dots, p_{i_q} \rangle$, which is called a q -face of σ_r and is denoted by $\sigma_q \leq \sigma_r$. If $\sigma_q < \sigma_r$, then σ_q is called a proper face of σ_r . We can further define the star of a face σ_p , denoted with $st(\sigma_p)$ as the union of all simplices of which σ_p is a face, and the link of a face σ_p , $lk(\sigma_p)$, as the union of all faces σ_f , $f \leq p$ in $st(\sigma_p)$ such that $\sigma_f \cap \sigma_p = \emptyset$.

Let T be a finite set of simplices. We define a SIMPLICIAL COMPLEX in $\mathbb{R}^m$ a finite set of simplexes T nicely fitted together, *i.e.*:

1. If $\sigma \in T$ and $\sigma' \leq \sigma$, then $\sigma' \in T$.
2. If $\sigma, \sigma' \in T$, then either $\sigma \cap \sigma' = \emptyset$ or $\sigma \cap \sigma' \leq \sigma$ and $\sigma \cap \sigma' \leq \sigma'$.

The dimension of a simplicial complex T is defined to be the maximum dimension of simplexes in T .

Thus, a simplicial complex is a collection of simplexes. If each simplex is regarded as a subset of $\mathbb{R}^m$, $m \geq \dim T$, the union of all the elements in T is a subset of $\mathbb{R}^m$ too. It is called the polyhedron $|T|$ associated to a simplicial complex T and it holds $\dim |T| = \dim T$.

Let X be a topological space. If there exist a simplicial complex T and an homeomorphism $f : |T| \rightarrow X$, then X is said to be triangulable and the pair (T, f) is called a TRIANGULATION OF X .

To be more precise let us consider a two dimensional simplicial complex T with underlying polyhedron $|T|$ and a vector $(N_0(T), N_1(T), N_2(T))$, where $N_i(T) \in \mathbb{N}$ is the number of i -dimensional simplexes in T . A Regge triangulation of a two dimensional Piecewise Linear (PL from now on) manifold M is an homeomorphism $|T| \rightarrow M$, where each face of T is realized by a rectilinear simplex of variable edge length $l(\sigma_1(i))$, $i = 1, \dots, N_1(T)$ of the appropriate dimension.

A dynamical triangulation is a particular case $|T_{l=a}| \rightarrow M$ of a Regge triangulation in which a PL manifold is realized by rectilinear and equilateral simplexes of constant edge length: $l(\sigma_1(i)) = a$, $\forall i = 1, \dots, N_1(T)$.

A triangulation of a two dimensional PL surface can be further described by its connectivity. This is specified via an adjacency matrix, *i.e.* the $N_0 \times N_0$ matrix:

$$B_{ij} = \begin{cases} 1 & \text{if } \langle p_i p_j \rangle \text{ exists} \\ 0 & \text{otherwise} \end{cases} \quad (1.1)$$

A Regge Triangulation has connectivity fixed *a priori*. However, in the following we will deal with triangulations with both variable edge length (thus

not dynamical) and variable connectivity. These structure have been first introduced in [60], where they have been baptized Random Regge Triangulations (RRT).

The metric structure of a RRT is locally Euclidean everywhere, with the exception of the vertexes $\sigma_0(i)$ (also called *the bones*), where the sum of the dihedral angles $\theta(\sigma_2)$ of the incident triangles σ can be in excess or in defect respect to the flatness condition $\theta(i) = \sum_{\sigma_2 \text{ inc}} \theta(\sigma_2) = 2\pi$, where the sum extends to all the triangles incident in the i -th bone. The first case is the negative curvature one, while in the second case, we speak of positive curvature. The correspondent deficit angle is defined as:

$$\varepsilon(i) = 2\pi - \sum_{\sigma_2 \text{ inc}} \theta(\sigma_2), \quad i = 1, \dots, N_0(T) \quad (1.2)$$

If we denote with $K_0(T) = \{\sigma_0(i)\}, i = 1, N_0(T)$ the 0-skeleton of the triangulation, then $M \setminus K_0(T)$ is a flat Riemannian manifold, and any point in the interior of an r -simplex σ_r has a neighborhood homeomorphic to $B^r \times C(lk(\sigma_r))$, where B^r is the r -dimensional ball in $\mathbb{R}^m$ and $C(lk(\sigma_r))$ is the cone over the link of σ_r (*i.e.* the product $lk(\sigma_r) \times [0, 1]$, with $lk(\sigma_r) \times \{1\}$ shrunked to a point).

For the simpler case of dynamical triangulations, the deficit angles are generated by the CURVATURE ASSIGNMENT, *i.e.* the string of integers $\{q(i)\}_{i=1}^{N_0(T)}$ which specifies the number of equilateral triangles incident in the i -th bone, via the relation:

$$\varepsilon(i) = 2\pi - q(i) \arccos \frac{1}{2} \quad i = 1, \dots, N_0(T). \quad (1.3)$$

For a regular (not necessary dynamical) triangulation we have $q(k) \geq 3 \forall k$ and, since each triangle has three internal angles, the curvature assignments obey the following constraint[60]:

$$\sum_{k=1}^{N_0(T)} q(k) = 3N_2(T) = 6 \left[1 - \frac{\chi(M)}{N_0(T)} \right] N_0(t) \quad (1.4)$$

where $\chi(M)$ is the Euler-Poincaré characteristic of the surface and $6 \left[1 - \frac{\chi(M)}{N_0(T)} \right]$ is the average value of the curvature assignments. In particular, if we remove the constraint $q(k) \geq 3$ we are dealing with generalized RRT, namely configurations in which the star of a vertex contains only two or one bidimensional simplexes.

In what follow, we will summarize the main results of [61] showing that it is possible to geometrically characterize the metrical structure of a generalized RRT described above as a particular case of the theory of singular Riemann surfaces endowed with a SINGULAR EUCLIDEAN STRUCTURE[70, 71]. In the next section, we will describe results, of [60], showing that this structure can be described in terms of complex functions theory.

Let us consider the first barycentric subdivision of $|T_i| \rightarrow M$ and, within this subdivision, the closed stars of the vertexes of the original triangulation.




Figure 1.1: Relations between the edge-lengths of the conical polytope and the edge-lengths of the triangulation

They form a collection of two-cells $\{\rho^2(i)\}_{i=1}^{N_0(T)}$ characterizing a polytope P baricentrically dual to T . As remarked in [61], it is important to stress that we are not dealing with a rectilinear presentation of P , where the cells are realized by rigid polytopes, but with a geometrical presentation of P , where the two-cells retain the conical geometry induced by the deficitary sum of dihedral angles of the original triangulation. To this end, we endow each cell $\{\rho^2(i)\}_{i=1}^{N_0(T)}$ with a polar reference frame, centered on the vertex $\sigma_0(i)$. Denoting with $(\lambda(i), \chi(i))$ the polar coordinate of a point $p \in \rho^2(i)$, then $\rho^2(i)$ is geometrically realized as the space[61]

$$\rho^2(i) = \frac{\{(\lambda(i), \chi(i)) \mid \lambda(i) \geq 0; \chi(i) \in \mathbb{R}/(2\pi - \varepsilon)\mathbb{Z}\}}{(0, \chi(i)) \sim (0, \chi'(i))} \quad (1.5)$$

endowed with the metric

$$d^2\lambda(i) + \lambda^2(i) d^2\chi(i) \quad (1.6)$$

This definition characterize the conical Regge polytope $|P_{T_l}| \rightarrow M$ barycentrically dual to $|T_l| \rightarrow M$.

The relation between the conical geometry of the barycentrically dual polytope $|P_{T_l}| \rightarrow M$ and the original triangulation data are fixed once they are given the lengths of the 1-simplexes of $|T_l| \rightarrow M$. A direct calculation provides the lengths of the (third part of the) medians connecting the barycenter $\rho_0(i, j, k)$ of the triangle defined by the bones $\sigma_0(i)$, $\sigma_0(j)$ and $\sigma_0(k)$, $i, j, k = 1, \dots, N_2(T)$, to the middle points of its edges. Following notation in [60], let us denote with $l(i, j)$ the length of the edge of the triangle connecting the bones $\sigma_0(i)$ and $\sigma_0(j)$, while $\hat{L}(k)$ is the length of the segment connecting its middle point with

$\rho_0(i, j, k)$ (*i.e.* the third part of the median drawn from $\sigma_0(k)$, see fig. 1.1). We can write:

$$\begin{aligned}\hat{L}^2(i) &= \frac{1}{18}l^2(i, j) + \frac{1}{18}l^2(k, i) - \frac{1}{36}l^2(j, k) \\ \hat{L}^2(j) &= \frac{1}{18}l^2(j, k) + \frac{1}{18}l^2(i, j) - \frac{1}{36}l^2(k, i) \\ \hat{L}^2(k) &= \frac{1}{18}l^2(j, k) + \frac{1}{18}l^2(k, i) - \frac{1}{36}l^2(i, j)\end{aligned}$$

and

$$\begin{aligned}l^2(i, j) &= 8\hat{L}^2(i) + 8\hat{L}^2(j) - 4\hat{L}^2(k) \\ l^2(j, k) &= 8\hat{L}^2(j) + 8\hat{L}^2(k) - 4\hat{L}^2(i) \\ l^2(k, i) &= 8\hat{L}^2(k) + 8\hat{L}^2(i) - 4\hat{L}^2(j)\end{aligned}$$

In such a geometrical framework, let $\rho^2(k)$ be the generic 2-cell barycentrically dual to the bone $\sigma_0(k)$, $k = 1, \dots, N_0(T)$, and let us denote with:

$$L(k) \doteq \sum_{i=1}^{q(k)} L(\rho^1(i)) \quad (1.7)$$

the length of its boundary, where $L(\rho^1(i))$ is the length of its $q(k)$ ordered edges $\rho^1(i) \in |P_{T_l}| \rightarrow M$. If $\varepsilon(k)$ denotes the deficit angle associated to the bone $\sigma_0(k) \in |T_l| \rightarrow M$, the slant radius associated with the cell $\rho^2(k) \in |P_{T_l}| \rightarrow M$ is defined as:

$$r(k) \doteq \frac{L(k)}{2\pi - \varepsilon(k)}, \quad k = 1, \dots, N_0(T) \quad (1.8)$$

We can associate to each two-cell $\rho^2(k) \in |P_{T_l}| \rightarrow M$ an open ball $B^2(k) = \{p \in \rho^2(k)/\partial(\rho^2(k))\}$; it is contained in $st(\sigma_0(k))$. To any vertex we can associate a complex uniformizing coordinate $\zeta(k)$ defined on an open disk of radius $r(k)$:

$$B^2(k) \longrightarrow D_k(r(k)) \doteq \{\zeta(k) \in \mathbb{C} \mid 0 \leq \zeta(k) \leq r(k)\} \quad (1.9)$$

In terms of $\zeta(k)$ we can write explicitly the singular Euclidean metric characterizing the singular Euclidean structure on each open ball as[61]:

$$ds_k^2 \doteq e^{2u} |\zeta(k) - \zeta(k)(\sigma_0(k))|^{-2\frac{\varepsilon(k)}{2\pi}} |d\zeta(k)|^2, \quad (1.10)$$

where $u : B^2 \rightarrow \mathbb{R}$ is a continuous function such that

$$|\zeta(k) - \zeta(k)(\sigma_0(k))| \frac{\partial u}{\partial \zeta(k)} \xrightarrow{\zeta(k) \rightarrow \zeta(k)(\sigma_0(k))} 0 \quad (1.11a)$$

$$|\zeta(k) - \zeta(k)(\sigma_0(k))| \frac{\partial u}{\partial \bar{\zeta}(k)} \xrightarrow{\zeta(k) \rightarrow \bar{\zeta}(k)(\sigma_0(k))} 0 \quad (1.11b)$$

Up to the conformal factor e^{2u} , (1.10) is the metric of a Euclidean cone of total angle $\theta(k) = 2\pi - \varepsilon(k)$. We can glue together the uniformizations $\{D_k(r(k))\}_{k=1}^{N_0(T)}$ along the pattern defined by the 1-skeleton of the dual Regge polytope and generate on M the quasi conformal structure:

$$(M, C_{sg}) \doteq \bigcup_{|P_{T_l}| \rightarrow M} \{D_k(r(k)); ds_{(k)}^2\}_{k=1}^{N_0(T)} \quad (1.12)$$

If $|dt^2|$ is a conformally flat metric on M , then the quasi-conformal structure (M, C_{sg}) can be locally represented by the metric[61]:

$$ds_T^2 = e^{2v}|dt^2| \quad (1.13)$$

with conformal factor $v \equiv u - \sum_{k=1}^{N_0(T)} \left(-\frac{\varepsilon(k)}{2\pi}\right) \ln|\zeta(k) - \zeta(k)(\sigma_0(k))|$. As there are no natural choice for u , the metric is only well defined up to conformal symmetry: this allows to move within the conformal class of all the metrics possessing the same singular structure of the triangulated surface $|T_l| \rightarrow M$.

This quasi conformal structure defines a RRT as a particular case of the theory of singular Riemann surfaces[61]. As a matter of fact, the singular structure described above can be summarized introducing the associated real divisor:

$$Div(T) \doteq \sum_{k=1}^{N_0(T)} \left(-\frac{\varepsilon(k)}{2\pi}\right) \sigma_0(k) = \sum_{k=1}^{N_0(T)} \left(\frac{\theta(k)}{2\pi} - 1\right) \sigma_0(k) \quad (1.14)$$

supported on the set of bones $\{\sigma_0(k)\}_{k=1}^{N_0(T)}$. The degree of such a divisor is $|Div(T)| = \sum_{k=1}^{N_0(T)} \left(\frac{\theta(k)}{2\pi} - 1\right) = -\chi(M)$. The real divisor characterizes the Euler class of the pair $(|T_l| \rightarrow M, Div(T))$ (or, shortly, $(T, Div(T))$): Associating to $(T, Div(T))$ the Euler number[70]

$$e(T, Div(T)) = \chi(M) - |Div(T)| \quad (1.15)$$

it is possible to rewrite the Gauss-Bonnet formula as:

Lemma 1 *Gauss-Bonnet formula for triangulated surfaces[61]*

Let $(T, Div(T))$ be a triangulated surface with divisor:

$$Div(T) = \sum_{k=1}^{N_0(T)} \left(\frac{\theta(k)}{2\pi} - 1\right) \sigma_0(k)$$

associated with the vertexes incidences $\{\sigma_0(k)\}_{k=1}^{N_0(T)}$. Let ds^2 be the conformal metric (1.13) representing the divisor $Div(T)$. Then

$$\frac{1}{2\pi} \int K dA = e(T, Div(T))$$

where K and dA are respectively the curvature and the area element corresponding to the metric (1.13).

Since for a RRT it holds $e(T, \text{Div}(T)) = 0$, then $\frac{1}{2\pi} \int K dA = 0$, i.e. a triangulation $|T_l| \leftarrow M$ naturally carries a conformally flat structure. This result admits a non-trivial converse:

Theorem 1 *Troyanov-Picard[70]*

Let $((M, C_{sg}), \text{Div})$ be a singular Riemann surface with a divisor such that $e(T, \text{Div}(T)) = 0$. Then there exists on M a unique (up to homothety) conformally flat metric representing the divisor $\text{Div}(T)$.

With these last statements, authors of [61] fully characterized the metric triangulations as a particular case of the theory of singular Riemann surfaces. Moreover, thanks to the introduction of the real divisor (1.14) and exploiting the Poncaré-Klein-Koebe uniformization theorem, they showed that the singular Riemann surface $((M, C_{sg}), \text{Div})$ is a particular case of singular uniformizations of $(M, N_0) = M - \sum_{i=1}^{N_0(T)} \sigma_0(i)$, thus including the study of RRT into the theory of punctured surfaces[61]. However, they stressed that in due to the presence of null euler number, this approach would have involved non standard techniques to be dealt with. Thus, they pursued further this analysis relating the space of inequivalent singular Euclidean structure to the theory of uniformization of singular Euclidean surfaces via the properties of Jenkins-Strebel quadratic differentials.

Remark 1 *We would like to stress that the construction described above includes naturally particular limiting cases of Regge triangulation like those situations in which some of the vertexes are characterized by deficit angles $\varepsilon(K) \rightarrow 2\pi$, i.e. $\theta(k) = 0$. Such a situation correspond to having the cone $C[lk(\sigma_0(k))]$ over the link $lk(\sigma_0(k))$ realized as an Euclidean cone of total angle 0. These situations, which had always been considered as pathological, are no longer so in the connection introduced above: the corresponding two-cells $\rho^2(k) \in |P_{T_l}| \rightarrow M$ can be naturally endowed with the conformal Euclidean structure obtained by setting $\frac{\varepsilon(k)}{2\pi} = 1$ in (1.10). The remaining object is then (up to the conformal factor e^{2u}) the flat metric on the semi-infinite cylinder $\mathbb{S}^1 \times \mathbb{R}_+$ (a cylindrical end). Alternatively, it is possible to consider $\rho^2(k)$ endowed with the geometry of an hyperbolic cusp, i.e. an half-infinite cylinder endowed with the hyperbolic metric $\lambda(k)^{-2}(d\lambda(k)^2 + d\chi(k)^2)$. The triangles incident in $\sigma_0(k)$ are then realized as hyperbolic triangles with the vertex located at $\lambda(k) = \infty$ and null incident angle. The two points of view are strictly related: one can switch from the Euclidean representation to the hyperbolic one simply setting the conformal factor to $e^{2u} = \left(\ln \frac{1}{|\zeta(k) - \zeta(k)(\sigma_0(k))|} \right)^{-2}$. The presence of cylindrical ends is consistent with a singular Euclidean structure as long as the associated divisor satisfies the topological constraint $|\text{Div}(t)| = -\chi(M)$ [60].*




Figure 1.2: The two-cells $\rho^2(k)$ of the dual polytope $|P_{T_l}| \rightarrow M$ can be geometrically represented by a punctured disk uniformized by a conical metric, while its 1-skeleton is in one-to-one correspondence with a trivalent ribbon graph[67]

1.2 Ribbon graphs on Regge polytopes: moduli space parametrization

In the previous section we have shown that, given a triangulation $|T_l| \rightarrow M$ of a Riemann surface M , the two-cells of the dual Regge polytope $|P_{T_l}| \rightarrow M$ are geometrically realized by punctured disks, uniformized by a conical metric. Moreover, the geometrical realization of the 1-skeleton of the dual Regge polytope is a trivalent ribbon graph:

$$\Gamma = (\{\rho^0(k)\}, \{\rho^1(k)\}) \quad (1.16)$$

where the $\{\rho^0(k)\}_{k=1}^{N_2(t)}$ are the barycenter of the triangles $\{\sigma_2(k)\}_{k=1}^{N_2(t)} \in |T_l| \rightarrow M$, and $\{\rho^1(k)\}$ is the set of edges of $|P_{T_l}| \rightarrow M$. These are generated by the pairwise joining of the half edges $\rho^1(k)^+$ and $\rho^1(k)^-$ in through the barycenter $W(k)$, $k = 1, \dots, N_1(T)$ of the edges $\{\sigma_1(k)\}_{k=1}^{N_1(t)} \in |T_l| \rightarrow M$. If we formally introduce a degree-2 ghost-vertex on each $W(k)$, then the underling graph of $|P_{T_l}| \rightarrow M$ is the edge refinement of Γ (see fig. 1.2):

$$\Gamma_{\text{ref}} = \left(\{\rho^0(k)\} \bigsqcup_{h=1}^{N_1(T)} \{W(h)\}, \{\rho^1(j)^+\} \bigsqcup_{j=1}^{N_1(T)} \{\rho^1(j)^-\} \right). \quad (1.17)$$

The natural automorphism group $\text{Aut}(P_l)$ of $|P_{T_l}| \rightarrow M$ (*i.e.* the set of bijective maps $\Gamma = (\{\rho^0(k)\}, \{\rho^1(k)\}) \rightarrow \tilde{\Gamma} = (\{\tilde{\rho}^0(k)\}, \{\tilde{\rho}^1(k)\})$ preserv-

ing the incidence matrix of the graph), is the automorphism group of its edge refinement [62]: $Aut(P_{T_l}) \doteq Aut\Gamma_{\text{ref}}$.

The local uniformizing complex coordinate $\zeta(k)$, in terms of which we have explicitly express the conical metric uniformizing the neighborhood of conical singularities, provides a counterclockwise orientation of the two cells of $|P_{T_l}| \rightarrow M$. Such an orientation provides a cyclic ordering of the set of half edges incidents on each vertex $\rho^0(k)_{k=1}^{N_2(T)}$: according to these remarks, the 1-skeleton of $|P_{T_l}| \rightarrow M$ is a ribbon (or fat) graph, *i.e.* a graph together with a cyclic ordering of the set of half edges incident to each vertex of Γ . Conversely, any ribbon graph Γ characterize a Riemann surface $M(\Gamma)$ possessing Γ as a spine. Thus, the edge refinement of the 1-skeleton of the Regge dual graph of a RRT is in one-to-one correspondence with trivalent metric ribbon graphs.

The set of all such trivalent metric Ribbon graphs Γ with given edge set $e(\Gamma)$ can be characterize as a space homeomorphic to $\mathbb{R}_+^{|e(\Gamma)|}$ (where $|e(\Gamma)|$ is the number of edges in $e(\Gamma)$). The automorphism group $Aut_{P_{T_l}}$ acts naturally on $\mathbb{R}_+^{|e(\Gamma)|}$ via the homomorphism $Aut_{P_{T_l}} \rightarrow \mathfrak{S}_{e(\Gamma)}$ ($\mathfrak{S}_{e(\Gamma)}$ being the symmetric group over $|e(\Gamma)|$ elements).

In this connection, let $Aut_{\partial}(P_l) \subset Aut(P_l)$, denote the subgroup of ribbon graph automorphisms of the (trivalent) 1-skeleton Γ of $|P_{T_l}| \rightarrow M$ that preserve the (labeling of the) boundary components of Γ . Then, the space K_1RP_{g,N_0}^{met} of 1-skeletons of conical Regge polytopes $|P_{T_l}| \rightarrow M$, with $N_0(T)$ labelled boundary components, on a surface M of genus g can be defined by [62]

$$K_1RP_{g,N_0}^{met} = \bigsqcup_{\Gamma \in RGB_{g,N_0}} \frac{\mathbb{R}_+^{|e(\Gamma)|}}{Aut_{\partial}(P_l)}, \quad (1.18)$$

where the disjoint union is over the subset of all trivalent ribbon graphs (with labelled boundaries) satisfying the topological constraint $2 - 2g - N_0(T) < 0$, and which are dual to generalized triangulations. It follows, (see [62] theorems 3.3, 3.4, and 3.5), that the set K_1RP_{g,N_0}^{met} is locally modeled on a stratified space constructed from the components $\mathbb{R}_+^{|e(\Gamma)|}/Aut_{\partial}(P_l)$ by means of a (Whitehead) expansion and collapse procedure for ribbon graphs, which amounts to collapsing edges and coalescing vertexes, (the Whitehead move in $|P_{T_l}| \rightarrow M$ is the dual of the flip move [69] for triangulations). Explicitly, if $l(t) = tl$ is the length of an edge $\rho^1(j)$ of a ribbon graph $\Gamma_{l(t)} \in K_1RP_{g,N_0}^{met}$, then, as $t \rightarrow 0$, we get the metric ribbon graph $\hat{\Gamma}$ which is obtained from $\Gamma_{l(t)}$ by collapsing the edge $\rho^1(j)$. By exploiting such construction, we can extend the space K_1RP_{g,N_0}^{met} to a suitable closure $\overline{K_1RP_{g,N_0}^{met}}$ [72], (this natural topology on K_1RP_{g,N_0}^{met} shows that, at least in two-dimensional quantum gravity, the set of Regge triangulations with *fixed connectivity* does not explore the full configuration space of the theory). The open cells of K_1RP_{g,N_0}^{met} , being associated with trivalent graphs, have dimension

provided by the number $N_1(T)$ of edges of $|P_{T_l}| \rightarrow M$, *i.e.*

$$\dim [K_1 RP_{g,N_0}^{met}] = N_1(T) = 3N_0(T) + 6g - 6. \quad (1.19)$$

There is a natural projection[60]

$$\begin{aligned} p : K_1 RP_{g,N_0}^{met} &\longrightarrow \mathbb{R}_+^{N_0(T)} \\ \Gamma &\longmapsto p(\Gamma) = (l_1, \dots, l_{N_0(T)}), \end{aligned} \quad (1.20)$$

where $(l_1, \dots, l_{N_0(T)})$ denote the perimeters of the polygonal 2-cells $\{\rho^2(j)\}$ of $|P_{T_l}| \rightarrow M$. With respect to the topology on the space of metric ribbon graphs, the orbifold $K_1 RP_{g,N_0}^{met}$ endowed with such a projection acquires the structure of a cellular bundle. For a given sequence $\{l(\partial(\rho^2(k)))\}$, the fiber

$$p^{-1}(\{l(\partial(\rho^2(k)))\}) = \{|P_{T_l}| \rightarrow M \in K_1 RP_{g,N_0}^{met} : \{l_k\} = \{l(\partial(\rho^2(k)))\}\} \quad (1.21)$$

is the set of all generalized conical Regge polytopes with the given set of perimeters[60]. If we take into account the $N_0(T)$ constraints associated with the perimeters assignments, it follows that the fibers $p^{-1}(\{l(\partial(\rho^2(k)))\})$ have dimension provided by

$$\dim [p^{-1}(\{l(\partial(\rho^2(k)))\})] = 2N_0(T) + 6g - 6, \quad (1.22)$$

which again corresponds to the real dimension of the moduli space $\mathfrak{M}_{g,N_0}$ of N_0 -pointed Riemann surfaces of genus g .

The complex analytic geometry of the space of conical Regge polytopes which we will discuss in the next section generalizes the well-known bijection (a homeomorphism of orbifolds) between the space of metric ribbon graphs $K_1 RP_{g,N_0}^{met}$ (which forgets the conical geometry) and the moduli space $\mathfrak{M}_{g,N_0}$ of genus g Riemann surfaces $((M; N_0), \mathcal{C})$ with $N_0(T)$ punctures [62], [72]. This bijection results in a local parametrization of $\mathfrak{M}_{g,N_0}$ defined by

$$\begin{aligned} h : K_1 RP_{g,N_0}^{met} &\rightarrow \mathfrak{M}_{g,N_0} \times R_+^N \\ \Gamma &\longmapsto [((M; N_0), \mathcal{C}), l_i] \end{aligned} \quad (1.23)$$

where $(l_1, \dots, l_{N_0})$ is an ordered n -tuple of positive real numbers and Γ is a metric ribbon graphs with $N_0(T)$ labelled boundary lengths $\{l_i\}$ (figure 8).

If $\overline{K_1 RP_{g,N_0}^{met}}$ is the closure of $K_1 RP_{g,N_0}^{met}$, then the bijection h extends to $\overline{K_1 RP_{g,N_0}^{met}} \rightarrow \overline{\mathfrak{M}_{g,N_0}} \times R_+^{N_0}$ in such a way that a ribbon graph $\Gamma \in \overline{K_1 RP_{g,N_0}^{met}}$ is mapped in two (stable) surfaces M_1 and M_2 with $N_0(T)$ punctures if and only if there exists an homeomorphism between M_1 and M_2 preserving the (labeling of the) punctures, and is holomorphic on each irreducible component containing one of the punctures.

As remarked in [60], it is important to stress that even if ribbon graphs can be thought of as arising from Regge polytopes (with variable connectivity), the

morphism (1.23) only involves the ribbon graph structure and the theory can be (and actually is) developed with no reference at all to a particular underlying triangulation. In such a connection, the role of dynamical triangulations has been slightly overemphasized, they simply provide a convenient way of labeling the different combinatorial strata of the mapping (1.23), but, by themselves they do not define a combinatorial parametrization of $\overline{\mathfrak{M}}_{g,N_0}$ for any finite N_0 . However, it is very useful, at least for the purposes of quantum gravity, to remember the possible genesis of a ribbon graph from an underlying triangulation and be able to exploit the further information coming from the associated conical geometry. Such an information cannot be recovered from the ribbon graph itself (with the notable exception of equilateral ribbon graphs, which can be associated with dynamical triangulations), and must be suitably codified by adding to the boundary lengths $\{l_i\}$ of the graph a further decoration. This can be easily done by explicitly connecting Regge polytopes to punctured Riemann surfaces[60].

1.3 Dual uniformizations of a random Regge triangulated surface

The genesis of a ribbon graph as underlying structure of a RRT has been shown very useful because it has allowed to parametrize the moduli space of a RRT via the morphism between the space of trivalent metric ribbon graphs and $\mathcal{M}_{g,N} \times \mathbb{R}_+^N$. Nonetheless, the full set of data we are handling include the conical geometry defined on the Regge polytopes. This can be codified in the above context by adding to the ribbon graph data (*i.e.* to the boundary lengths $\{L(k)\}_{k=1}^{N_0(T)}$) a suitable decoration which arises naturally once we connect the dual Regge polytopes $|P_{T_i}| \rightarrow M$ to the theory of punctured Riemann surfaces.

In [62] it was introduced a ribbon graph uniformization which, exploiting properties of Jenkins-Strebel quadratic differential, allows to associate to a ribbon graph with given edge lengths $\{L(k)\}_{k=1}^{N_0(T)}$ a complex structure $((M, N_0), \mathcal{C})$, *i.e.* a punctured Riemann surface. This construction is a priori completely general: given a smooth Riemann surface with N marked points and $(C, (p_1, \dots, p_N))$ and an ordered N -tuple $(a_1, \dots, a_N) \in \mathbb{R}_+^N$, there is a unique meromorphic quadratic differential ¹ ϕ on M satisfying the following conditions:

¹Let us remember that an HOLOMORPHIC QUADRATIC DIFFERENTIAL defined on a compact Riemann surface M is an element of $H^0(C, K_C^{\otimes 2})$ (being $K_C^{\otimes 2}$ the second symmetric tensor product of the canonical sheaf on M , K_C). In a local coordinate frame on M the quadratic differential is represented by $\phi = f(z)(dz)^2$, $f(z)$ being a locally defined holomorphic function. Under a coordinate change $w = w(z)$ the local expression $\phi = f(z)(dz)^2 = g(w)(dw)^2$ transforms as

$$f(z) = g(w(z)) \left(\frac{dw(z)}{dz} \right)^2. \quad (1.24)$$

1. ϕ is holomorphic on $C/(p_1, \dots, p_N)$.
2. ϕ has a double pole at each $p_j, j = 1, \dots, N$.
3. The union of all non-compact horizontal leaves² form a zero-measure close subset of M .
4. Every compact horizontal leaf γ_j is a simple loop surrounding one of the poles and satisfying:

$$a_j = \int_{\gamma_j} \sqrt{\phi} \quad (1.25)$$

where the branch of the square root is chosen so that the integral has a positive value with respect to the positive orientation of γ_j , which is determined by the complex structure on M .

This quadratic differential is called STREBEL DIFFERENTIAL. The collection of all its compact horizontal leaves surrounding a pole forms a punctured disk centered on the pole itself. The punctured disk ends on the set non-compact horizontal leaves connecting pairwise the zeros of the differential. The latters are joined to the pole by a vertical leaf(see [62]). The union of the punctured disks (with the punctures filled by the points $(p_1, \dots, p_n)$), of the zeros of ϕ and of the non-compact horizontal leaves form the underlying topological surface of the Riemann surface M .

In this sense, a Strebel quadratic differential defined on a nonsingular Riemann surface M of genus g determines a unique cell decomposition $\square_\phi$ of M consisting of N_0 2-cells, N_1 1-cells and N_2 0-cells, where N_2 is the number of zeros of the Strebel differential and it holds $N_0 - N_1 + N_2 = 2 - 2g$. The 1-skeleton

A MEROMORPHIC quadratic differential on M is an holomorphic quadratic differential with the exception of a finite set $(p_1, \dots, p_N)$ of points of M such that at each singularity p_j of ϕ there is a local expression $\phi = f_j(z)(dz)^2$ with a meromorphic function $f_j(z)$ having a pole at $z = z(p_j)$. If $f_j(z)$ has a pole of order r at $z = z(p_j)$, then we say that ϕ has an order r pole at $z = z(p_j)$.

²Let us recall that given a meromorphic quadratic differential $\phi = f(z)(dz)^2$ defined on M , a real parametric curve

$$\gamma : (a, b) \ni t \mapsto \gamma(t) = z \in \mathbb{C}$$

parametrized on an open interval (a, b) of the real axis is

- an HORIZONTAL LEAF (OR HORIZONTAL TRAJECTORY) of ϕ if


$$f(\gamma(t)) \left(\frac{d\gamma(t)}{dt} \right)^2 > 0$$

for every value of $t \in (a, b)$;

- a VERTICAL LEAF (OR VERTICAL TRAJECTORY) of ϕ if

$$f(\gamma(t)) \left(\frac{d\gamma(t)}{dt} \right)^2 > 0$$

for every value of $t \in (a, b)$.


 Figure 1.3: Differences between $|T_l| \rightarrow M$ and Δ_ϕ

of such a cellular decomposition, *i.e.* the union of the 0-cells and 1-cells of the cell decomposition which is defined by the Strebel quadratic differential is a metric ribbon graph.

Moreover, the set of all the vertical leaves that connect zero and poles of the Strebel differential, together with the cell decomposition $\square_\phi$ induced on M by the non-compact horizontal leaves, form a canonical triangulation (in the sense described above) Δ_ϕ of M .


With these remarks, in [62] it had been shown how to construct a canonical coordinate system on a Riemann surface M : once they are given a N marked points M and an N -tuple of real numbers, we simply have to introduce the local expression of the associated Strebel differential, where local means valid in each 0,1,2-cell of $\square_\phi$.

In [60], this construction had been directly applied to the triangulation $|T_l| \rightarrow M$, once they have been performed the following identifications:

(M, ϕ)		RRT
$(p_1, \dots, p_N)$	$\longleftrightarrow$	$\{\sigma_0(i)\}_{i=1}^{N_0(T)}$
$(a_1, \dots, a_N)$	$\longleftrightarrow$	$\{L(i) = L(\partial\rho^2(i))\}_{i=1}^{N_0(T)}$
$\square_\phi$	$\longleftrightarrow$	$ P_{T_l} \rightarrow M$

Let us stress that it does not hold the identification between Δ_ϕ and $|T_l| \rightarrow M$: the canonical triangulation Δ_ϕ is obtained via the triangulation of each polygon of $\square_\phi$, while $|T_l| \rightarrow M$ is the Regge dual of $|P_{T_l}| \rightarrow M$ (see fig. 1.3).

Let us consider the dual Regge polytope $|P_{T_l}| \rightarrow M$ and its underlying ribbon graph Γ . Let $\rho^2(i) \in |P_{T_l}| \rightarrow M$, $\rho^2(j) \in |P_{T_l}| \rightarrow M$ and $\rho^2(k) \in |P_{T_l}| \rightarrow M$ be three pairwise adjacent two-cell barycentrically dual to the vertexes $\sigma_0(i) \in |T_l| \rightarrow M$, $\sigma_0(j) \in |T_l| \rightarrow M$, $\sigma_0(k) \in |T_l| \rightarrow M$ (see fig 1.4). Each edge of the ribbon graph Γ is shared by two triangles of Δ_ϕ which can be gluing together to get the purple diamond shape in fig. 1.4. This is the set of vertical leaves intersecting $\rho_1(i, j)$. Let $\rho_0(i, j, k)$ and $\rho_0(j, l, i)$ be the two endpoints of the

Figure 1.4: Dual Regge polytope associated to $|T_l| \rightarrow M$

strip and let us assign it an arbitrary direction, for example from $\rho_0(i, j, k)$ to $\rho_0(j, l, i)$. For an arbitrary point P the diamond, the canonical coordinate $z(i, j) = \int_{\rho_0(i, j, k)}^P \sqrt{\phi|_{\rho_1(i, j)}}$ maps the diamond to the strip

$$U_{\rho_1(i, j)} = \{z(i, j) \in \mathbb{C} \mid 0 \leq z(i, j) \leq l(\rho^1(i, j))\} \quad (1.26)$$

of infinite height and width $l(\rho^1(i, j))$, where $l(\rho^1(i, j))$ is the length of the considered edge, *i.e.* the strip is the local uniformization on the Riemann surface of the union of the two triangles. The local expression $\phi|_{\rho_1(i, j)}$ of the Strebel differential is

$$\phi|_{\rho_1(i, j)} = (dz(i, j))^2 \quad (1.27)$$

On the other hand, the vertexes of the ribbon graphs are, according to the cellular decomposition induced by the Strebel differential, the zero of the differential itself. Every quadratic differential has an expression $\phi = \frac{m^2}{4} \omega^{m-2} (d\omega)^2$ around a zero of degree $m - 2$. Thus we can use it as the expression of the Strebel differential on an open neighborhood

$$U_{\rho_0(i, j, k)} = \{\omega(i, j, k) \in \mathbb{C} \mid |\omega(i, j, k)| \leq \delta, \omega(i, j, k)(\rho_0(i, j, k)) = 0\} \quad (1.28)$$

surrounding $\rho_0(i, j, k)$. Since the 1-skeleton of $|P_{T_l}| \rightarrow M$ is a trivalent graph, $m = 3$ for each vertex, thus:

$$\phi|_{\rho_0(i, j, k)} = \frac{9}{4} \omega(i, j, k) (d\omega(i, j, k))^2 \quad (1.29)$$

The transformation law (1.24) on the intersection $U_{\rho_0(i, j, k)} \cap U_{\rho_1(i, j)}$ gives the transition function between the two coordinates patches:

$$\omega(h, j, k) = \begin{cases} z(h, j)^{\frac{2}{3}} \\ e^{\frac{2\pi}{3}i} z(j, k)^{\frac{2}{3}} \\ e^{\frac{4\pi}{3}i} z(k, h)^{\frac{2}{3}}. \end{cases} \quad (1.30)$$




Figure 1.5: The complex coordinate neighborhoods associated to the dual polytope[60]

Due to the conical geometry retained by the deficit angles, the extension the uniformization on the 2-cells of the cellular decomposition (*i.e.* on the Regge polytopes $\rho_2(i)$) is a little more subtle. We will show that is possible to keep track of the conical geometry of the polytopes in two different ways, thus leading to two dual uniformizations.

Since the condition (1.25), we can choose a local coordinate $\zeta(i)$ on an open disk $U_{\rho_2(i)}$ defined by the union of all the compact horizontal leaves homotopic to $\sigma_0(i)$ such that³

$$\phi|_{\rho_2(i)} = -\frac{L(i)^2}{4\pi^2} \frac{d\zeta(i)^2}{\zeta(i)^2} \quad (1.31)$$

If $z(i, j_\nu)$, $\nu = 1, q(i)$ are the coordinates defined on the edges surrounding the given 2-cell $\rho_2(i)$, it holds on the intersections $U_{\rho_2(i)} \cap U_{\rho_1(i, j_n u)}$:

$$(dz(i, j_\nu))^2 = -\frac{L(i)^2}{4\pi^2} \frac{d\zeta(i)}{\zeta(i)^2} \quad (1.32)$$

If we fix the integration constants such that the uniformizing unit disk

$$U_{\rho_2(i)} = \{\zeta(i) \in \mathbb{C} \mid |\zeta(i)| < 1, \zeta(i)(\sigma_0(i)) = 0\} \quad (1.33)$$

covers the full polytope $\rho_2(i)$, we get the following transition functions on $U_{\rho_2(i)} \cap U_{\rho_1(i, j_n u)}$ [60]:

$$\zeta(i) = \exp \left(\frac{2\pi i}{L(i)} \sum_{\beta=1}^{\nu-1} (L(i, j_\beta) + z(i, j_\nu)) \right) \quad \nu = 1, \dots, q(i), \quad (1.34)$$

where $L(i, j) = l(\rho^1(i, j))$ and $\sum_{\beta=1}^{\nu-1} \doteq 0$ if $\nu = 1$.

Since for any closed curve $c : \mathbb{S}^1 \rightarrow U_{\rho_2(i)}$ homotopic to the boundary of $\overline{U}_{\rho_2(i)}$ it holds

$$\oint_c \sqrt{\phi|_{\rho_2(i)}} = L(i), \quad (1.35)$$

³Let us remember the identification $a_i \leftrightarrow L(i)$, $i = 1, \dots, N_0(T)$

then the geometry associated with $\phi|_{\rho^2(i)}$ is described by a cylindrical metric associated with a quadratic differential with a second order pole, *i.e.*

$$|\phi(i)| \doteq |\phi|_{\rho^2(i)} = \frac{L(i)^2}{4\pi^2} \frac{|d\zeta(i)|^2}{|\zeta(i)|^2}. \quad (1.36)$$

We are dealing with such a limiting situation like those described at the end of section 1.1: the punctured disk $\Delta_i^* \subset U_{\rho^2(i)}$

$$\Delta_i^* = \{\zeta(i) \in \mathbb{C} \mid 0 < |\zeta(i)| < 1\} \quad (1.37)$$

endowed with the cylindrical metric (1.36) is isometric to a flat semi-infinite cylinder.

To keep track of the conical geometry associated to the polygonal cell $\rho^2(i)$ we can recall that the conformal factor in (1.10) defines the metric on the polytopes only up to conformal symmetry. Thus, for a given deficit angle $\varepsilon(i) = 2\pi - \theta(i)$ we can conformally relate the cylindrical geometry (1.36) and the conical one (1.10) fixing the conformal factor u such that:

$$ds_{(i)}^2 \doteq \frac{L(i)^2}{4\pi^2} |\zeta(i)|^{-2\frac{\varepsilon(i)}{2\pi}} |d\zeta(i)|^2 \quad (1.38)$$

We can eventually state:

Proposition 1 [60] *If $\{p_k\}_{k=1}^{N_0} \in M$ denotes the set of punctures corresponding to the decorated vertexes $\{\sigma^0(k), \frac{\varepsilon(k)}{2\pi}\}_{k=1}^{N_0}$ of the triangulation $|T_l| \rightarrow M$, let RP_{g,N_0}^{met} be the space of conical Regge polytopes $|P_{T_l}| \rightarrow M$, with $N_0(T)$ labelled vertexes, and let $\mathfrak{P}_{g,N_0}$ be the moduli space $\mathfrak{M}_{g,N_0}$ decorated with the local metric uniformizations $(\zeta(k), ds_{(k)}^2)$ around each puncture p_k . Then the map*

$$\Upsilon : RP_{g,N_0}^{met} \longrightarrow \mathfrak{P}_{g,N_0}$$

provided by

$$\begin{aligned} \Upsilon : (|P_{T_l}| \rightarrow M) &\longrightarrow ((M; N_0), \mathcal{C}); \{ds_{(k)}^2\} = \\ &= \bigcup_{\{\rho^0(i,j,k)\}}^{N_2(T)} U_{\rho^0(i,j,k)} \bigcup_{\{\rho^1(i,j)\}}^{N_1(T)} U_{\rho^1(i,j)} \bigcup_{\{\rho^2(k)\}}^{N_0(T)} (\Delta_k^*, ds_{(k)}^2), \end{aligned} \quad (1.39)$$

defines the decorated, N_0 -pointed, Riemann surface $((M; N_0), \mathcal{C})$ canonically associated with the conical Regge polytope $|P_{T_l}| \rightarrow M$.

The dual decoration: trading curvature for moduli

The dual decoration, which was introduced in [60], is quite more subtle and arises when we open the cone over $\rho^2(i)$ into its constituent conical sectors. To show how it works, let $W_\alpha(k)$, $\alpha = 1, \dots, q(k)$ be the barycenters of the edges $\sigma^1(\alpha) \in |T_l| \rightarrow M$ incident on $\sigma^0(k)$, and intercepting the boundary $\partial(\rho^2(k))$ of

the polygonal cell $\rho^2(k)$. Let us denote with $\widehat{L}_\alpha(k)$ the length of the polygonal lines between the points $W_\alpha(k)$ and $W_{\alpha+1}(k)$ (with α defined mod $q(k)$). Let us go back for a while. In the uniformization (1.9), endowed with the metric (1.10) (where we have set $\zeta(k)(\sigma_0(k)) = 0$) the points $\{W_\alpha(k)\}$ characterize a corresponding set of points on the circumference $\{\zeta(k) \in \mathbb{C} \mid |\zeta(k)| = l(\partial(\rho^2(k)))\}$, and an associated set of $q(k)$ generators $\{\overline{W_\alpha(k)\sigma^0(k)}\}$ on the cone $C[lk(\sigma^0(k))]$. Such generators mark $q(k)$ conical sectors

$$S_\alpha(k) \doteq \left(c_\alpha(k), \frac{L(k)}{\theta(k)}, \vartheta_\alpha(k) \right), \quad (1.40)$$

with base

$$c_\alpha(k) \doteq \{|\zeta(k)| = L(k), \arg W_\alpha(k) \leq \arg \zeta(k) \leq \arg W_{\alpha+1}(k)\}, \quad (1.41)$$

slant radius $\frac{L(k)}{\theta(k)}$, and with angular opening

$$\vartheta_\alpha(k) \doteq \frac{\widehat{L}_\alpha(k)}{L(k)} \theta(k), \quad (1.42)$$

where $\theta(k) = 2\pi - \varepsilon(k)$ is the given conical angle. The cone $(D_k(r(k)), ds_{(k)}^2)$ (see eq. (1.9)) can be formally represented as:

$$(D_k(r(k)), ds_{(k)}^2) = \cup_{\alpha=1}^{q(k)} S_\alpha(k). \quad (1.43)$$

If we split the vertex of the cone and of the associated conical sectors $S_\alpha(k)$, then the conical geometry of $(D_k(r(k)), ds_{(k)}^2)$ can be equivalently described by a cylindrical strip of height $\frac{l(\partial(\rho^2(k)))}{\theta(k)}$ decorating the boundary of $\rho^2(k)$. Each sector $S_\alpha(k)$ in the cone gives rise, in such a picture, to a rectangular region in the cylindrical strip[60]. Introducing a complex variable $v(k) = x(k) + i y(k) \in \mathbb{C}$, we can explicitly parametrize any such region as:

$$R_{\vartheta_\alpha(k)}(k) \doteq \left\{ v(k) \in \mathbb{C} \mid 0 \leq x(k) \leq \widehat{L}_\alpha(k), 0 \leq y(k) \leq \frac{L(k)}{\theta(k)} \right\}. \quad (1.44)$$

In this connection, we can go a step further. If we consider the full cylinder, we can think to parametrize it with the complex coordinate z introduced above. In this case the associated rectangular region is:

$$R_{\theta(k)}(k) \doteq \left\{ v(k) \in \mathbb{C} \mid 0 \leq x(k) \leq L(k), 0 \leq y(k) \leq \frac{L(k)}{\theta(k)} \right\}. \quad (1.45)$$

By means of the canonical conformal transformation:

$$\zeta(k) = e^{\frac{2\pi i}{L(k)} v(k)} \quad (1.46)$$

we can conformally map the cylinder into the annulus:

$$\Delta_{\varepsilon(k)}^* \doteq \left\{ \zeta(k) \in \mathbb{C} \mid e^{-\frac{2\pi}{2\pi - \varepsilon(k)}} \leq |\zeta(k)| \leq 1 \right\} \subset \overline{U_{\rho^2(k)}}. \quad (1.47)$$




Figure 1.6: Opening the cones[60]

Then the surface with boundary:

$$M_{\partial} \doteq ((M_{\partial}, N_0), \mathcal{C}) = \bigcup_{\{\rho^0(i,j,k)\}}^{N_2(T)} U_{\rho^0(i,j,k)} \bigcup_{\{\rho^1(i,j)\}}^{N_1(T)} U_{\rho^1(i,j)} \bigcup_{\{\rho^2(k)\}}^{N_0(T)} (\Delta_{\varepsilon(k)}^*, |\phi(k)|) \quad (1.48)$$

defines the blowing up of the conical geometry of $((M, N_0), \mathcal{C}, ds_{(k)}^2)$ along the ribbon graph Γ [60].

The metrical geometry of $(\Delta_{\varepsilon(k)}^*, |\phi(k)|)$ is that of a flat cylinder with a circumference lenght given by $L(k)$ and height given by $\frac{L(k)}{\theta_k}$ ⁴. We also have

$$\partial M_{\partial} = \bigsqcup_{k=1}^{N_0(T)} S_{\theta(k)}^+ \quad \partial \Gamma = \bigsqcup_{k=1}^{N_0(T)} S_{\theta(k)}^- \quad (1.49)$$

where the circles

$$S_{\theta(k)}^+ = \left\{ \zeta(k) \in \mathbb{C} \mid |\zeta(k)| = e^{-\frac{2\pi}{2\pi - \varepsilon(k)}} \right\} \quad (1.50a)$$

$$S_{\theta(k)}^- = \{ \zeta(k) \in \mathbb{C} \mid |\zeta(k)| = 1 \} \quad (1.50b)$$

denote respectively the inner and outer boundary of the annulus $\Delta_{\varepsilon(k)}^*$. Note that collapsing $S_{\theta(k)}^+$ to a point we get the original cones $(\Delta_k^*, ds_{(k)}^2)$. Thus, the surface with boundary M_{∂} naturally correspond to the ribbon graph Γ , naturally associated with the 1-skeleton of the dual Regge polytope $|P_{T_l}| \rightarrow M$, decorated with the *finite* cylindrical ends $\Delta_{\varepsilon(k)}^*$ [60].

This latter uniformization had proven invaluable in discussing the modular aspects of simplicial quantum gravity, in particular, it allowed a detailed analysis of the connection between the Weil-Petersson volume of the (compactified) moduli space of genus g Riemann surfaces with N_0 punctures $\overline{\mathfrak{M}}_{g, N_0}$ and the dynamical triangulation partition function for pure gravity [60].

According to the above description, it follows that a metric triangulation can be seen either as the (singular Euclidean) uniformization $((M, N_0), \mathbb{C}; ds_{(k)}^2)$ of a punctured Riemann surface, or as the uniformization of an open Riemann surface $((M, \partial M_{(k)}), \mathbb{C}; |\phi(k)|)$ with finite cylindrical ends $\Delta_{\varepsilon(k)}^*$ [60]. Underlying

⁴Let us recall that this is the slant radius of the generalized euclidean cone $(\Delta_k^*, ds_{(k)}^2)$ of base circumference $L(k)$ and vertex conical angle $\theta(k)$

this geometric duality there is the peculiar role played by the deficit angles $\varepsilon(k)$ associated with the vertexes of the triangulation. They represent the localized curvature degrees of freedom of the triangulation, and together with the perimeter length $L(k)$ of the 2-cell dual to the given vertex (puncture), they provide the relevant geometric information for uniformizing (with a conical Euclidean metric) the pointed Riemann surface around each puncture. Conversely, in the open Riemann surface with (N_0) boundaries $\partial M_{(k)}$, the deficit angles (again together with the corresponding 2-cell perimeters) are directly turned into geometric moduli of the associated annuli[67]:

$$m(k) \doteq \frac{1}{2\pi} \ln \frac{1}{e^{-\frac{2\pi}{2\pi - \varepsilon(k)}}} = \frac{1}{2\pi - \varepsilon(k)}. \quad (1.51)$$

In other words, *localized curvature can be traded into modular data*. These two points of view merge one into the other when the deficit angles $\varepsilon(k)$ all degenerate into 2π . This corresponds to the limiting situation when the triangles incident on the given vertex of the triangulation are no longer Euclidean but rather ideal hyperbolic triangles: each finite cylindrical end turns into a hyperbolic cusp. From a modular point of view we may say that the two dual singular Euclidean uniformization representing a given random Regge triangulation merge in a unique hyperbolic uniformization of the same underlying closed (pointed) Riemann surface.

Chapter 2

BCFT on the annuli: Langlands' boundary states

Starting from this chapter, we develop the natural framework to describe the coupling of a matter field theory with two dimensional quantum gravity in the framework described in the previous chapter, *i.e.* with the peculiar geometry defined the the dual uniformized open Riemann surface M_{∂} (see eq. (1.48)) associated with the random Regge triangulation $|T| \rightarrow M$.

The aim of investigating string dualities in such a discretized approach, leads us to deal with non-critical Polyakov string theory. Thus, let us consider D scalar fields X^{α} , $\alpha = 1, \dots, D$, defined as injection maps from the Riemann surface M_{∂} to an unspecified target space $\mathcal{T}$. In this connection, at fixed N_0 and fixed genus g , the strategy we will follow to quantize the bosonic fields on the open Riemann surface with boundaries defined in equation (1.48) calls into play Boundary Conformal Field Theory (BCFT). As a matter of fact, we will discuss the quantization of bosonic fields on each annular region $\Delta_{\varepsilon(k)}^*$, defined in (1.47), then we shall glue together the resulting BCFTs along the intersection pattern defined by the ribbon graph Γ associated with the triangulation. Finally, as usual in (dynamical or random) triangulation, the coupling with two-dimensional gravity on the Riemann surface will be (at least formally) obtained by summing over all possible triangulation of the original Riemann surface M , *i.e.* in the dual picture, over all possible ribbon graphs Γ .

When we describe the metric geometry of the triangulation $|T_l| \rightarrow M$ as the dual open Riemann surface M_{∂} , we are actually unwrapping each conical 2-cell $\rho^2(k)$ of the dual polytope into a corresponding finite cylindrical end. The restriction of each bosonic field to the outer boundary, $X^{\alpha}(k)|_+$, viewed as embedding variables, a priori injects $S_{\varepsilon(k)}^{(+)}$ on a circle of radius $R(k)$ whose length $2\pi R(k)$ is not necessarily equal to the circumference of the boundary itself, *i.e.* $L(k)$. The same is true for the behavior of the fields (injection maps)

on the inner boundary $S_{\varepsilon(k)}^{(-)}$, so we can extend this behavior to the fields defined on the whole cylinder and assume that in general they are compactified on a circle of circumference $2\pi \frac{R^\alpha(k)}{l(k)}$, where $l(k)$ is a length parameter built on the characteristic data of the triangulation. In order to not impose any restrictions over this “unwrapping” process, we will adopt the general condition:

$$X^\alpha(k)(e^{2\pi i}\zeta, e^{-2\pi i}\bar{\zeta}) = X^\alpha(k)(\zeta, \bar{\zeta}) + 2\nu^\alpha(k)\pi \frac{R^\alpha(k)}{l(k)}, \quad \nu^\alpha(k) \in \mathbb{Z} \quad (2.1)$$

stating that each field $X^\alpha(k)$ winds $\nu(k)$ times around the toric cycles of the compact target space $\mathcal{T}$.

Once we are given a Conformal Field Theory (CFT from now on) on a closed Riemann Surface (*i.e.* the Riemann sphere), the problem to extend this theory to a theory defined on surfaces with boundary is encoded into find which boundary conditions can be imposed consistently at the boundaries (in a more stringy language, this is the question of how many consistent open string theories can be added to a given closed string one). In our connection, we have to specify consistent boundary conditions for the fields $X^\alpha(\zeta(k), \bar{\zeta}(k))$, $\alpha = 1, \dots, D$ on the boundaries of the cylindrical region $\Delta_{\varepsilon(k)}^*$:

$$S_{\varepsilon(k)}^{(+)} \doteq \left\{ \zeta(k) \in \mathbb{C} \mid |\zeta(k)| = e^{\frac{2\pi}{2\pi-\varepsilon(k)}} \right\} \quad (2.2a)$$

$$S_{\varepsilon(k)}^{(-)} \doteq \{ \zeta(k) \in \mathbb{C} \mid |\zeta(k)| = 1 \} \quad (2.2b)$$

This problem is rather similar to the closed conformal field theory problem arising when we ask if, given a CFT on the Riemann sphere, this is sufficient to define uniquely the same CFT on higher genus surfaces. The answer to this question is known exactly: the theory on the sphere determines uniquely the CFT on higher genus surfaces because of locality of the Operator Product Expansions (OPE): their coefficients are uniquely defined on the sphere and do not “see” the different genus of the surrounding surface. However, asking for consistency on higher genus surfaces implies one more condition: the correlation functions must transform under the action of the modular group $SL(2, \mathbb{Z})$ at genus one.

An analogous general result holding for BCFT question is not really known. Once given a theory on the full complex plane, usually referred to as *the bulk theory*, what we know is the complete list of *sewing constraints* that have to be satisfied on if we introduce a boundary[73, 74]. However, it is not known *a priori* for which theory it is possible to find a complete set of solutions of these constraints. The set of examples in which a solution has been found shows that, even in this case, modular invariance plays a fundamental role: this is due to the similarity between the classification of modular invariant partition functions and the classification of non-negative integer matrix representation of

the fusion algebra (see [75] and references therein). To some extents, the theory of the compactified boson belongs to this set.

The OPE coefficient, which are fully determined by the bulk theory, define an algebra of fields which the boundary conditions must respect. We will show that each boundary condition can be described by a coherent state built on the states space of the bulk theory. However, not every such a boundary states describes an admissible extension of the bulk theory. The main point is that boundary states must preserve the bulk algebra of fields stating a precise gluing relation between their left and right modes.

Aiming to expose results in a self-contained way, we start the chapter with a brief introduction to the fundamental concepts in (boundary) conformal field theory. The expert reader can skip this part, and jump directly to section 2.2.

2.1 Bulk and Boundary Conformal Field Theory

To some extents, the CFT associated to the compactified free boson belongs to the set of conformal field theories whose extension on an arbitrary surface with boundary can be completely work out once we know properties of the same theory defined on the full complex plane, this latter being usually referred to as the *bulk theory*.

Let us recall that, given a vector space $\mathcal{H}$ and the vector space of linear endomorphism of $\mathcal{H}$ which are finite linear combination of homogeneous endomorphism, we can define a *field of conformal dimension* $(h, \bar{h}) \in \mathbb{Z}_+^2$ an $\text{End}V$ -valued formal power series in $\zeta, \bar{\zeta}$:

$$\phi(\zeta, \bar{\zeta}) = \sum_{m,n \in \mathbb{Z}^2} \phi_{m,n} \zeta^{-m-h} \bar{\zeta}^{-n-\bar{h}}, \quad (2.3)$$

i.e. the coefficient of this formal Laurent expansion are operators in $\text{End}V$

A bulk conformal theory is defined as an Hilbert space of states $\mathcal{H}^{(C)}$, endowed with the action of an Hamiltonian $H^{(C)}$ and of a vertex operation, *i.e.* a formal map

$$\Phi^{(C)}(\circ; \zeta, \bar{\zeta}) : \mathcal{H}^{(C)} \rightarrow \text{End}V[\zeta, \bar{\zeta}] \quad (2.4)$$

associating to each vector $|\phi\rangle \in \mathcal{H}^{(C)}$ a conformal field $\phi(\zeta, \bar{\zeta})$ of conformal dimension $h, \bar{h}$.

The bulk theory is completely worked out once they are known the coefficients of the Operator Product Expansion (OPE) for all fields in the theory. Actually, this task is tractable for most of the CFTs, since the states in $\mathcal{H}^{(C)}$ (and consequently the associated vertex operators) can be organized into irreducible representation of the observable algebra generated by the chiral fields.

Chiral fields depend on only one coordinate, ζ or $\bar{\zeta}$, thus they are either

holomorphic or antiholomorphic:

$$W^a(\zeta) = \sum_m W_m^a \zeta^{-m-h^a} \quad \bar{W}^a(\bar{\zeta}) = \sum_m \bar{W}_m^a \bar{\zeta}^{-m-\bar{h}^a} \quad (2.5)$$

where the index a labels the different chiral fields in the theory and h^a and $\bar{h}^a$ are respectively the conformal weights of $W^a(\zeta)$ and $\bar{W}^a(\bar{\zeta})$. Their Laurent modes W_m^a and $\bar{W}_m^a$ generate two commuting chiral algebras, $\mathcal{W}$ and $\bar{\mathcal{W}}$, which we will assume to be isomorphic.

The Virasoro fields T and $\bar{T}$ play a special role among the chiral fields of a CFT. Their modes L_n and $\bar{L}_n$ close two copies of the Virasoro algebra with central extension c :

$$\begin{aligned} [L_m, L_n] &= (m-n) L_{m+n} + \frac{c}{12} n(n^2-1) \delta_{m+n,0} \\ [\bar{L}_m, \bar{L}_n] &= (m-n) \bar{L}_{m+n} + \frac{c}{12} n(n^2-1) \delta_{m+n,0} \end{aligned}$$

The Virasoro algebra belongs to the enveloping algebra of the chiral algebra $\mathcal{W}$. As a matter of fact, since we can represent the holomorphic and antiholomorphic components of the stress-energy tensor as the ordered (creator-annihilator) product $T(z) = \gamma :W(\zeta)^a W(\zeta)^a:$, we get a representation of the their Laurent modes via the Sugawara construction:

$$L_0 \doteq \sum_{n>0} W_{-n} \cdot W_n + \frac{1}{2} (W_0)^2 \quad (2.6a)$$

$$L_n \doteq \frac{1}{2} \sum_{m \in \mathbb{Z}} :W_{n-m} \cdot W_m: \quad (2.6b)$$

The action of the chiral algebra modes W_m and $\bar{W}_m$ determine a decomposition of the Hilbert space into all the irreducible modules of the two commuting chiral algebra:

$$\mathcal{H}^{(C)} \doteq \bigoplus_{\lambda \bar{\lambda}} \mathcal{H}^\lambda \otimes \bar{\mathcal{H}}^{\bar{\lambda}}, \quad (2.7)$$

where the vectors (highest weights) $\{\lambda^a\}$ and $\{\bar{\lambda}^a\}$ are the charges (*i.e.* respectively the W_0^a and $\bar{W}_0^a$ eigenvalues) associated to all the possible ground-states (highest weight states) $|\lambda\rangle$ and $|\bar{\lambda}\rangle^1$, and where each chiral module is defined as:

$$\mathcal{H}_\lambda \doteq \text{Span} \{ W_{-m_1}^a \dots W_{-m_N}^a |\lambda\rangle \mid m_i > 0 \forall i = 1, \dots, N \}. \quad (2.8)$$

An equivalent definition holds for the antiholomorphic sector.

¹Let us recall that such ground-states are defined via the relations (holomorphic sector):

$$\begin{aligned} W_n^a |\lambda\rangle &= 0 & \forall a = 1, \dots, \dim \mathcal{W} \text{ and } \forall n > 0 \\ \bar{W}_n^a |\bar{\lambda}\rangle &= 0 & \forall a = 1, \dots, \dim \mathcal{W} \text{ and } \forall n > 0 \end{aligned}$$

From these definition we can introduce the vertex operator associated to a holomorphic ground-state $|\lambda\rangle$:

$$\varphi_\lambda(\zeta) \doteq \Phi^{(C)}(|\lambda\rangle; \zeta) = c_\lambda e^{\int \lambda_-(\zeta) d\zeta} e^{i\lambda W_0} z^{i\lambda q} e^{\int \lambda_+(\zeta) d\zeta} \quad (2.9)$$

where $\lambda_\pm(\zeta) = \sum_{m>0} \lambda W_{\pm m} z^{\mp m-1}$, c_λ is an arbitrary cocycle and q is a position operator.

The vertex operator associated with a single oscillator acting on the conformal vacuum is defined by:

$$\Phi^{(C)}(W_{-m}^a |0\rangle; \zeta) = \frac{1}{(m-1)!} \frac{d^{m-1}}{d\zeta} W^a(\zeta) \quad (2.10)$$

For a general homogeneous element $|\phi\rangle = W_{-m_1}^{a_1} \dots W_{-m_N}^{a_N} |\lambda\rangle$, linearity of $\text{End}\mathcal{H}$ ensures:

$$\Phi^{(C)}(|\phi\rangle; \zeta) = :\Phi^{(C)}(W_{-m_1}^{a_1} |0\rangle; \zeta) \dots \Phi^{(C)}(W_{-m_N}^{a_N} |0\rangle; \zeta) \Phi^{(C)}(|\lambda\rangle; \zeta): \quad (2.11)$$

In the definition (2.7), the module $\mathcal{H}^0$ refers to the vacuum representation. From the reconstruction equations (2.9) and (2.10), we see that it is mapped directly into $\mathcal{W}$ by the vertex operation defined by $\Phi^{(C)}$.

Each irreducible representation $\mathcal{H}^\lambda$ receive a $\mathbb{Z}$ -grading by the action of L_0 . As a matter of fact the L_0 eigenvalues (*i.e.* the conformal weights)

$$L_0(W_{-m_1} \dots W_{-m_N})|\lambda\rangle = m_1 + \dots + m_N + h|\lambda\rangle$$

define the level of a state as: $l = \sum_{i=1}^N m_i$ so that each irreducible representations of $\mathcal{W}$ can be decomposed as

$$\mathcal{H}^\lambda = \bigoplus_{l \geq 0} \mathcal{H}_l^\lambda. \quad (2.12)$$

With the above remarks, we see that each copy of the chiral algebra fulfills all the requirements of a vertex algebra²: for the holomorphic sector, the grading

²Let us recall that given a $\mathbb{Z}$ -graded vector space $V = \bigoplus_m V_m$, an endomorphism E of V is homogeneous of degree j if $E(V_m) \in V_{m+j}$. Thus we can define[76]:

Definition 1 A VERTEX ALGEBRA is a collection of data:

- a $\mathbb{Z}$ -graded vector space of states $V = \bigoplus_{m=0}^\infty V_m$;
- a vacuum vector $|0\rangle \in V_0$
- a translation operator $\omega : V \rightarrow V$ of degree one;
- a vertex operation, i.e. a linear map $\Phi(\circ, z) : V \rightarrow \text{End}V[[z, z^{-1}]]$ taking each state $|\phi\rangle \in V_n$ into a field of conformal dimension n

subjected to the following constraints:

- VACUUM AXIOM: $\Phi(|0\rangle, z) \doteq \mathbb{I}_V$
- TRANSLATIONAL AXIOM: For any $|\phi\rangle \in V$, $[\omega, \Phi(|\phi\rangle, z)] = \partial_z \Phi(|\phi\rangle, z)$.
- LOCALITY AXIOM: all fields $\Phi(|\phi\rangle, z)$ are mutually local with each other.

of the Hilbert space is the one induced by the action of L_0 described above. The conformal vacuum is identified with the ground state $|0\rangle = |\lambda = 0\rangle$, whose associated vertex operator is indeed $\mathbb{I}_{\mathcal{H}}$. Finally, the role of a traslation operator is played by L_{-1} , which satisfies $\Phi(L_{-1}|\phi\rangle, \zeta) = \frac{d}{d\zeta}\Phi(|\phi\rangle, \zeta)$ and $\Phi(|0\rangle, \zeta) = 0$, as we can see directly applying the Sugawara representation (2.6).

Let us consider, for each irreducible representations of the holomorphic³ chiral algebra $\mathcal{H}^\lambda$, the eigenspace of 0-level states V_0^λ , *i.e.*, in the case that $\mathcal{H}^\lambda$ are the highest weight representations of the chiral algebra, the subspace containing only the (eventually degenerate) highest weight state. It will carry an irreducible action of the horizontal subalgebra of $\mathcal{W}$ (*i.e.* the subalgebra closed by all the zero modes W_0^a) via the action of the associated linear map:

$$\mathbb{X}_{W^a}^\lambda \doteq W_0^a|_{V_0^\lambda} : V_0^\lambda \longrightarrow V_0^\lambda \quad (2.13)$$

As a consequence, $\varphi_\lambda(\zeta)$ defined in equation (2.9) is the holomorphic part of the (Virasoro primary) fields which arise as $\mathcal{W}$ -primary via the action of the horizontal subalgebra $\{W_0^a\}$. We can gather the holomorphic and antiholomorphic part in a single $\mathcal{W}$ -primary field $\varphi_{\lambda,\bar{\lambda}}(\zeta, \bar{\zeta})$ associated to states in $V_0^\lambda \otimes \bar{V}_0^\lambda$.

2.1.1 Glueing conditions and boundary fields

Dealing with open string theory naturally impose to study the consequences of conformal invariance on surfaces delimited by one or more boundaries, on which we have to define suitable field behaviour. The prototype example, mostly investigated in literature, is the BCFT defined on the Upper Half complex Plane (UHP from now on). Its results are easily extended to other restricted geometries via suitable local conformal transformations.

Thus, let us consider the complex coordinate $z = x + iy$ and let us restrict ourselves to $\Im z \geq 0$. In the interior part of the domain, the structure of the boundary CFT is fixed by the requiring local equivalence with the associated bulk theory: thus, locally, there is a one-to-one correspondence between fields of the boundary theory and fields of the parent bulk theory. Their local structures coincide in the sense that both theories have identical equal time commutators with chiral fields and identical operator product expansions. In particular, it exists a stress energy tensor $T_{x_1, x_2}(x, y)$ defined in $\Delta_{\varepsilon(k)}^*$ and, in this connection, the presence of a boundary for $\Im z = 0$ requires that no energy flows across the boundary itself. Introducing the holomorphic and antiholomorphic components, $T(z) = 2(T_{xx} + iT_{xy})$ and $\bar{T}(\bar{z}) = 2(T_{xx} - iT_{xy})$, this implies the condition

$$T(z) = \bar{T}(\bar{z})|_{y=0}. \quad (2.14)$$

When the bulk theory contains other chiral fields W and $\bar{W}$ of conformal dimension h_W and $\bar{h}_{\bar{W}}$, we have also to require the following continuity condition:

³Obviously, the same holds for the antiholomorphic sector

$$W(z) = \Omega \overline{W}(\bar{z})|_{y=0} \quad (2.15)$$

where Ω is an automorphism of the Dynkin diagram associated to the extended chiral algebra [77]. **To keep contact with the notation mostly used in literature, we use Ω to define the glueing automorphism on the boundary. It has not to be confused with the compactification radius $\Omega^\alpha(k)$.**

Since they are defined only on half complex plane, W and $\overline{W}$ are not sufficient to construct the action of two commuting copies of the chiral algebra. However, the assumption of the existence of continuity conditions in (2.14) and (2.15) has the powerful consequence to give rise to the action of only one copy of the chiral algebra on the state space $\mathcal{H}^{(H)}$ of the boundary theory. As a matter of fact, we can combine W and $\overline{W}$ into a single object $\mathbf{W}(z)$ which, considering the conformal field theory defined on the UHP domain can be chosen as:

$$\mathbf{W}(z) \doteq \begin{cases} W(z) & \text{when } \Im z \geq 0 \\ \Omega \overline{W}(\bar{z}) & \text{when } \Im z < 0 \end{cases} \quad (2.16)$$

Obviously, different domains allow for different choices of $\mathbf{W}(z)$. The key point is that, thanks to the gluing condition on the boundary, the field we are now dealing with is analytic on the whole complex plane; thus, it can be expanded in a Laurent series: $\mathbf{W}(z) = \sum_n W_n^{(H)} z^{-n-h}$, introducing the modes:

$$W_n^{(H)} = \frac{1}{2\pi i} \int_C dz z^{n+h} W(z) - \frac{1}{2\pi i} \int_C d\bar{z} \bar{z}^{n+\bar{h}} \Omega \overline{W}(\bar{z}) \quad (2.17)$$

Ward identities for correlators of the boundary theory can be retrieving from the singular part of the OPE of chiral fields with bulk fields $\varphi(z, \bar{z})$. OPE are fixed by requirement of local equivalence between the bulk theory and the bulk of the boundary theory. In we define the singular part of $\mathbf{W}$ with $\mathbf{W}_>(z) = \sum_{n>-h} W_n z^{-n-h}$, The singular part of the OPE is then given by

$$\begin{aligned} \mathbf{W}(w) \varphi(\zeta, \bar{z}) &\sim [\mathbf{W}_>(w), \varphi(\zeta, \bar{z})] = \\ &\sum_{n>-h} \left(\frac{1}{(w-z)^{n+h}} \Phi(W_n^{(C)} \varphi; z, \bar{z}) + \frac{1}{(w-\bar{z})^{n+h}} \Phi(\Omega \overline{W}_n^{(C)} \varphi; z, \bar{z}) \right) \end{aligned} \quad (2.18)$$

We have placed a superscript (C) on the modes $W_n^{(C)}, \overline{W}_n^{(C)}$ to display clearly that they act on the elements $\varphi \in \mathcal{H}^{(C)}$ labeling the bulk fields in the theory. Let us drop superscript (H) from now on: objects without superscripts are intended to belong to the boundary theory, while we will put the superscript (C) when we will have to consider objects belonging to the bulk theory.

The sum on the right hand side of equation (2.18) is always finite because φ is annihilated by all modes $W_m^{(C)}, \overline{W}_m^{(C)}$ with sufficiently large m . For $\Im w > 0$ only the first terms involving $W_n^{(C)}$ can become singular and the singularities

agree with the singular part of the OPE between $W(w)$ and $\varphi(z, \bar{z})$ in the bulk theory. In the same way, the singular part of the OPE between $\Omega \bar{W}(w)$ and $\varphi(z, \bar{z})$ in the bulk theory is reproduced by the terms which contain $\bar{W}_n^{(C)}$, if $\Im w < 0$.

For future utility, let us specialize formula (2.18) to the case of a chiral current $\mathbf{J}(w)$ (thus $h_J = 1$) acting on a primary field $\varphi_{\lambda, \bar{\lambda}} \in \mathcal{H}_\lambda \otimes \bar{\mathcal{H}}_{\bar{\lambda}}$. In this case, equation (2.18) reduces to

$$\mathbf{J}(w) \varphi_{\lambda, \bar{\lambda}}(z, \bar{z}) \sim \frac{X_J^\lambda}{w - z} \varphi_{\lambda, \bar{\lambda}}(z, \bar{z}) - \varphi_{\lambda, \bar{\lambda}}(z, \bar{z}) \frac{X_{\Omega \bar{J}}^{\bar{\lambda}}}{w - \bar{z}} \quad (2.19)$$

The linear maps X_W^λ and $X_{\Omega \bar{J}}^{\bar{\lambda}}$ has been introduced in equation (2.13); they act on $\varphi_{\lambda, \bar{\lambda}}$ by contraction in the first tensor component.

Ward identities for arbitrary n -point functions of fields φ_{ij} follow directly from equation (2.18). They have the same form as those for chiral conformal blocks in a bulk CFT with $2n$ insertions of chiral vertex operators with charges $i_1, \dots, i_n, \omega(j_1), \dots, \omega(j_n)$, where ω is the automorphism of the fusion rules algebra induced by Ω . In many concrete examples, one has rather explicit expressions for such chiral blocks. So we see that objects familiar from the construction of bulk CFT can be used as building blocks of correlators in the boundary theory (“doubling trick”). Note, however, that the Ward identities depend on the gluing map Ω .

Thanks to Ward identities and bulk fields OPE we can reduce the computation of correlators of n bulk fields to a product of 1-point functions $\langle \phi_{ij} \rangle_\alpha$. These no longer vanish close to a boundary because, traslation invariance towards the boundary itself is broken, and they can depend both on their distance from the boundary itself and, more in general, from the boundary condition. Transformation properties of bulk fields with respect to L_0 , $L_{\pm 1}$ and W_0 fix the form of the one-point function to

$$\langle \phi_{ij}(z, \bar{z}) \rangle_\alpha = \frac{A_{ij}^\alpha}{(z - \bar{z})^{h_i + h_j}} \quad (2.20)$$

where $A_{ij}^\alpha : \mathcal{H}_0^j \rightarrow \mathcal{H}_0^i$ obeys $\mathbf{X}_W^i A_{ij}^\alpha = A_{ij}^\alpha \mathbf{X}_{\Omega W}^j$. This last relation implies $j = \omega(i^+)$, thus $h_i = h_j$.

These parameters are not completely free. Sewing constraint has been worked out by several authors (see [75], [78] and references therein). In particular, if we consider the two point function of bulk primary fields $\langle \phi_{ij} \rangle_\alpha$, cluster properties of CFT and conformal Ward identities (see [78] and [74] for further details) allows to write the following relation for 1-point function coefficients:

$$A_i^\alpha A_j^\alpha = \sum_k \Xi_{ijk} A_0^\alpha A_k^\alpha, \quad (2.21)$$

where we have defined $A_i^\alpha \doteq A_{i\omega(i^+)}^\alpha$. The coefficients Ξ_{ijk} can be expressed as

combination of the coefficients of bulk OPE and of the fusing matrices relating the conformal blocks of the bulk theory.

The existence on the boundary of the gluing map Ω , defined in (2.15), give rise to the action of one chiral algebra $\mathcal{W}$ on the state space of the boundary theory and inducing a decomposition of the open CFT Hilbert Space $\mathcal{H}$ into irreducible representations of $\mathcal{W}$ [75]:

$$\mathcal{H} = \bigoplus_{\lambda} \mathcal{V}_{\lambda}. \quad (2.22)$$

This allows to introduce a new one-to-one state-field correspondence Φ between states $|\psi\rangle \in \mathcal{H}$ and so-called boundary fields $\psi(x)$, which are defined (at least) for x on the real line [79]. Let us suppose that boundary condition does not jump on the real axis (we will understand the meaning of this assertion later). Thus, it exists a unique sl_2 invariant vacuum state $|0\rangle$. Then, for any state $|\psi\rangle \in \mathcal{H}$, there exists a boundary operator $\psi(x) \doteq \Phi(\psi; x)$ such that:

$$\psi(x) |0\rangle = e^{xL_{-1}} |\psi\rangle \quad (2.23)$$

for all x in the real line. In particular, the operator $\psi(0)$ creates the state from the conformal vacuum state $|0\rangle$. Note also that L_{-1} generate translations along the boundary.

The conformal dimension of a boundary field $\psi(x)$ can be read off from the L_0 -eigenvalue of the corresponding state $|\psi\rangle \in \mathcal{H}$. If $|\psi\rangle$ is a primary state of conformal weight h , definition of conformal modes analogue to (2.17) leads to:

$$[L_n, \psi(x)] = x^n \left(x \frac{d}{dx} + h(n+1) \right) \psi(x) \quad (2.24)$$

The boundary fields assigned to elements in the vacuum sector $\mathcal{H}^0$ coincide with the chiral fields in the theory, i.e. $\mathbf{W}(x) = \Phi(w; x)$ for some $w \in \mathcal{H}^0$ and $\Im x = 0$. These fields can always be extended beyond the real line and coincide with either W or $\Omega \bar{W}$ in the bulk. If other boundary fields admit such an extension, this suggests an enlargement of the chiral algebra in the bulk theory.

Decomposition (2.22) also implies that the partition function may be expressed as a sum of characters $\chi_i(q)$ of the chiral algebra,

$$Z_{(\Omega, \alpha)}(q) := \text{Tr}_{\mathcal{H}}(q^{L_0 - c/24}) = \sum_i n_i^{\Omega \alpha} \chi_i(q), \quad (2.25)$$

where $n_i^{\Omega \alpha} \in \mathbb{N}$ are multiplicity coefficients.

The existence of boundary extra fields motivates to consider more general correlation functions in which the bulk-field on the interior of the UHP appear together with boundary fields $\psi(x)$. Following the standard reasoning in CFT, it is easy to conclude that the bulk fields $\varphi_{ij}(z, \bar{z})$ give singular contributions to the correlation functions whenever z approaches the real line [74]. This can be seen

from the fact that the Ward identities describe a mirror pair of chiral charges i and $\omega(j)$ placed on both sides of the boundary. Therefore, the singularities in an expansion of $\varphi(z, \bar{z}) \equiv \varphi_{ij}(z, \bar{z})$ around $x = \Re z$ are given by primary fields which are localized at x on the real line, i.e. the boundary fields $\psi(x)$. In other words, the observed singular behavior of bulk fields $\varphi(z, \bar{z})$ near the boundary with associated boundary condition α may be expressed in terms of a bulk-boundary OPE [74]

$$\varphi(z, \bar{z}) = \sum_k (2y)^{h_k - h - \bar{h}} C_{\varphi k}^{\alpha} \psi_k(x), \quad (2.26)$$

where $\psi_k(x)$ are primary fields of conformal weight h_k . Which ψ_k can possibly appear on the rhs. of (2.26) is determined by the chiral fusion of i and $\omega(j)$, but some of the coefficients $C_{\varphi k}^{\alpha}$ may vanish for some α .

2.2 The quantum amplitude on the annulus

To work out an explicit expression for an (open) string amplitude over M_{∂} , let us approach to the problem dealing first with building blocks of such an amplitude, *i.e.* the partition functions over the annular domains $\Delta_{\varepsilon(k)}^*$, $k = 1, \dots, N_0$.

As first remark, let us notice that the computation can be performed independently for each component X^{α} , $\alpha = 1, \dots, D$. As a matter of fact, in a stringy language the world-sheet action for the bosonic fields (string coordinates) is:

$$S = \frac{1}{4\pi} \int d\zeta(k) d\bar{\zeta}(k) G_{\alpha\beta}(k) \partial X^{\alpha}(k) \bar{\partial} \bar{X}^{\beta}(k) + B_{\alpha\beta}(k) \partial X^{\alpha}(k) \bar{\partial} \bar{X}^{\beta}(k) - \frac{1}{2} \Phi(k) R^{(2)} \quad (2.27)$$

where $G(k)$, $B(k)$ and $\Phi(k)$ are respectively the background metric, the background Kalb-Ramond field and the dilaton. Sometimes in string theory literature the D^2 components of G and B , which encode the geometrical data of the background, are gathered into a matrix E , whose symmetric part is G and whose antisymmetric part is B :

$$E = G + B. \quad (2.28)$$

E is usually called BACKGROUND MATRIX.

We will deal with flat toroidal backgrounds, *i.e.* we will consider a string moving in a background in which D dimensions are compactified and the metric G , the Kalb-Ramond field B and the dilaton are X independent. In particular, in the Polyakov background, we will put to zero the B -field and we will encode all data about the background metric in the compactification radius value, so that the background metric can be chosen diagonal and with $G_{\alpha\alpha}(k) = 1 \forall \alpha = 1, \dots, D$. Then, the world-sheet action on $\Delta_{\varepsilon(k)}^*$ is:

$$S = \frac{1}{8\pi} \int d\zeta(k) d\bar{\zeta}(k) \partial X^{\alpha}(k) \bar{\partial} \bar{X}^{\alpha}(k) \quad (2.29)$$

For the compactified boson, the dynamical degrees of freedom are defined via the modes of the bosonic fields $X^\alpha(\zeta(k), \bar{\zeta}(k))$ obeying the equation of motion $\partial\bar{\partial}X^\alpha(\zeta(k), \bar{\zeta}(k)) \doteq \Delta(k) = 0$, where $\Delta(k)$ is the Laplacian defined over $\Delta_{\varepsilon(k)}^*$.

The bulk theory sees a free boson as a closed string injection map propagating from the remote past ($y \rightarrow -\infty$) to the remote future ($y \rightarrow \infty$) and, after the conformal mapping in equation (1.46), we can write the following mode expansion:

$$X^\alpha(\zeta(k), \bar{\zeta}(k)) = X_0^\alpha(k) - ip_L^\alpha(k) \ln \zeta(k) - ip_R^\alpha(k) \ln \bar{\zeta}(k) + i \sum_{n \neq 0} \frac{1}{n} \left(\mathfrak{a}_n^\alpha(k) \zeta^{-n} + \bar{\mathfrak{a}}_n^\alpha(k) \bar{\zeta}^{-n} \right), \quad (2.30)$$

where we can identify the center of mass momentum components with the chiral zero modes:

$$p_L^\alpha(k) = \mathfrak{a}_0^\alpha(k) \quad p_R^\alpha(k) = \bar{\mathfrak{a}}_0^\alpha(k) \quad (2.31)$$

The only chiral fields are the holomorphic and antiholomorphic currents $J^\alpha(\tilde{\zeta}) = i \partial X^\alpha(\tilde{\zeta}) = \sum_n \mathfrak{a}_n^\alpha \tilde{\zeta}^{-n-1}$ and $\bar{J}^\alpha(\bar{\zeta}) = i \bar{\partial} X^\alpha(\bar{\zeta}) = \sum_n \bar{\mathfrak{a}}_n^\alpha \bar{\zeta}^{-n-1}$ which generate, for each field $\alpha = 1, \dots, D$, N_0 copies of the affine $\hat{u}(1)_L \otimes \hat{u}(1)_R$ algebra:

$$\begin{aligned} [\mathfrak{a}_n^\alpha(k), \mathfrak{a}_m^\beta(k)] &= n \delta_{n+m,0} \delta^{\alpha,\beta} & [\bar{\mathfrak{a}}_n^\alpha(k), \bar{\mathfrak{a}}_m^\beta(k)] &= n \delta_{n+m,0} \delta^{\alpha,\beta} \\ [\mathfrak{a}_n^\alpha(k), \bar{\mathfrak{a}}_m^\beta(k)] &= 0. \end{aligned} \quad (2.32)$$

At this level we can think that the theory on each cylindrical end decouples from the others. Thus, from now on, we will suppress the polytope index (k). We will assume to have N_0 copies of the same theory, and we will explain later how data propagate over the ribbon graph, *i.e.* how these different copies glue over it.

The bulk Hamiltonian is $H = \frac{2\pi}{L} \left(\frac{1}{2} \mathfrak{a}_0^2 + \frac{1}{2} \bar{\mathfrak{a}}_0^2 + \frac{1}{2} \sum_{m \neq 0} \mathfrak{a}_{-m} \cdot \mathfrak{a}_m + \bar{\mathfrak{a}}_{-m} \cdot \bar{\mathfrak{a}}_m \right)$, from which, replacing the expression of Virasoro generators obtained via the Sugawara construction in this particular case, eq. (2.6) with central extension $c = D$, we get:

$$H^{(C)} = \frac{2\pi}{L} \left(L_0 + \bar{L}_0 - \frac{D}{12} \right). \quad (2.33)$$

In the diagonal decomposition (2.7) of the Hilbert space into irreducible representations of the two chiral algebras, λ and $\bar{\lambda}$ are the (real numbers) $\hat{u}(1)_L$ and $\hat{u}(1)_R$ charges respectively.

Due to the winding condition (2.1), highest weight states carries different charges w.r.t the action of the zero modes $\mathfrak{a}_0$ and $\bar{\mathfrak{a}}_0$. The total North's momentum is $p^\alpha = \frac{1}{2} (\lambda^\alpha + \bar{\lambda}^\alpha)$, while the variation of α -th field along a closed cycle is $X^\alpha(k) (e^{2\pi i} \zeta, e^{-2\pi i} \bar{\zeta}) - X^\alpha(k) (\zeta, \bar{\zeta}) = 2\pi (\mathfrak{a}_0^\alpha - \bar{\mathfrak{a}}_0^\alpha)$. Thus, the holo-

morphic and antiholomorphic $u(1)$ charges

$$\lambda_{(\mu,\nu)}^\alpha = \mu^\alpha \frac{l(k)}{R^\alpha(k)} + \frac{1}{2} \nu^\alpha(k) \frac{R^\alpha(k)}{l(k)} \quad (2.34a)$$

$$\bar{\lambda}_{(\mu,\nu)}^\alpha = \mu^\alpha \frac{l(k)}{R^\alpha(k)} - \frac{1}{2} \nu^\alpha(k) \frac{R^\alpha(k)}{l(k)} \quad (2.34b)$$

We can collect the holomorphic and antiholomorphic ground-states into a single object $|\lambda^\alpha, \bar{\lambda}^\alpha\rangle = |\lambda_{(\mu,\nu)}^\alpha\rangle \otimes |\bar{\lambda}_{(\mu,\nu)}^\alpha\rangle$. The action of Heisenberg algebra on this vacuum is obvious:

$$\mathbf{a}_n^\alpha |(\mu^\alpha, \nu^\alpha)\rangle \doteq \mathbf{a}_n^\alpha \otimes \mathbb{I} |(\mu^\alpha, \nu^\alpha)\rangle = \mathbf{a}_n^\alpha |\lambda_{(\mu,\nu)}^\alpha\rangle \otimes |\bar{\lambda}_{(\mu,\nu)}^\alpha\rangle \quad (2.35a)$$

$$\bar{\mathbf{a}}_n^\alpha |(\mu^\alpha, \nu^\alpha)\rangle = \mathbb{I} \otimes \bar{\mathbf{a}}_n^\alpha |(\mu^\alpha, \nu^\alpha)\rangle = |\lambda_{(\mu,\nu)}^\alpha\rangle \otimes \bar{\mathbf{a}}_n^\alpha |\bar{\lambda}_{(\mu,\nu)}^\alpha\rangle \quad (2.35b)$$

and the associated chiral primary field is $:e^{i\lambda_\alpha X_L^\alpha(\zeta)} e^{i\bar{\lambda}_\alpha X_R^\alpha(\bar{\zeta})}:$.

Chiral Fock spaces are constructed by applying the rising operators on the chiral vacua. From the Sugawara construction of Virasoro generators, we see that $L_n |(\mu^\alpha, \nu^\alpha)\rangle = 0 \forall n > 0$, thus the vacua form a continuous set of highest weight states with holomorphic (resp. antiholomorphic) highest weights $h = \frac{1}{2}\lambda^2$ (resp. $\bar{h} = \frac{1}{2}\bar{\lambda}^2$).

The decomposition of the Fock spaces in term of Verma modules is straightforward: since the descendant states are obtained by the repeated application of $\mathbf{a}_{-n}$, $n > 0$, which is equivalent to the repeated application of L_{-n} , $n > 0$, because $\mathbf{a}_{-n}$ also raises the conformal dimension of n , the Fock spaces and the Verma modules coincide, $\mathcal{H}^\lambda = \mathcal{V}^h$. However, this happens only as long as $\lambda \neq \sqrt{2}m$, with $m \in \frac{1}{2}\mathbb{Z}$. On the other hand, when $\lambda = \sqrt{2}m$ with $m \in \frac{1}{2}\mathbb{Z}$, the presence of a null state⁴ at level $2|m| + 1$, causes each Fock space to be a reducible representation of the Virasoro algebra and to decompose into an infinite sum of Verma modules:

$$\mathcal{H}^{\sqrt{2}m} = \bigoplus_{l=1}^{\infty} \mathcal{V}^{(|m|+l)^2} \quad (2.36)$$

Let us stress that this decomposition has nothing to do with the grading induced by the zero modes of the Virasoro algebra and shown in equation (2.12): this last decomposition holds for each value of the left and right chiral algebra charges.

This decomposition shows an hidden $SU(2)$ symmetry. As a matter of fact, the Virasoro highest weight state with highest weight $h = j^2$ spans in the set

⁴Let us recall the definition and the main properties of a null state[80]:

Definition 2 Null state

Any descendant state $|\chi\rangle = L_{-k_1} \dots L_{-k_N}$, $k_1, \dots, k_N > 0$, which is annihilated by all the Virasoro generators L_n with $n > 0$ is called a SINGULAR VECTOR (or NULL VECTOR or NULL STATE).

Such a state generates its own Verma module $\mathcal{V}^\chi \in \mathcal{V}^\lambda$: as a matter of fact, singular vectors are orthogonal to the whole Verma module, and this property extends to all its own descendants.

$\{\mathcal{H}^{\sqrt{2}m}$ a $(2|m| + 1)$ -dimensional multiplet built with all the $\mathbf{u}_1$ highest weight defined by a value of m fulfilling the condition $(|m| + l) = j^2, l \in \mathbb{Z}_+$.

The same holds for the antiholomorphic sector, thus we can write[78]:

$$\begin{aligned} \mathcal{H}^{(C)} = & \int_{\substack{\lambda \neq \sqrt{2}m \\ \bar{\lambda} \neq \sqrt{2}n}} \mathcal{H}^\lambda \otimes \overline{\mathcal{H}}^{\bar{\lambda}} \oplus \bigoplus_{m,n \in \frac{\mathbb{Z}}{2}} \mathcal{H}^{\sqrt{2}m} \otimes \overline{\mathcal{H}}^{\sqrt{2}n} = \\ & \int_{\substack{\lambda \neq \sqrt{2}m \\ \bar{\lambda} \neq \sqrt{2}n}} \mathcal{V}_{\frac{\lambda^2}{2}} \otimes \overline{\mathcal{V}}_{\frac{\bar{\lambda}^2}{2}} \oplus \bigoplus_{j, \bar{j} \in \frac{\mathbb{Z}_+}{2}} \left(\mathcal{V}_{j^2}^{\oplus 2j+1} \otimes \overline{\mathcal{V}}_{\bar{j}^2}^{\oplus 2\bar{j}+1} \right) \quad (2.37) \end{aligned}$$

As outlined in the introduction, from a microscopic point of view, adding boundaries implies to define suitable boundary values assignments. In a stringy language, it means to work out which open string theory can be naturally related to closed string. As well known, closed string theory is naturally coupled to interacting open string. In this connection, our cylindrical end can be viewed both as an open string one-loop diagram or as a tree level closed string diagram, with a closed string propagating for a finite length path. In the open string (direct) channel, time flows around the cylinder. The associated quantization scheme defines functions of the modular parameter $\tau = i\theta(k) = i(2\pi - \varepsilon(k))$ (see eq. (1.51)). The close string (transverse) channel, sees time flowing along the cylinder, and the associated quantization scheme is related to the open string one via the modular transformation $\tau \rightarrow -\frac{1}{\tau}$. In the following, we will switch back and forth between these two points of view and the associated quantization schemes.

The amplitude associated to this diagram is a deep-investigated object when the boundary assignment are a priori defined as the usual specification of Neumann or Dirichlet boundary conditions (for a review see [81] and references therein). However, when we describe the metric geometry of the triangulation $|T_l| \rightarrow M$ with the Riemann surface M_∂ , we are actually unwrapping each conical two-cell of the dual polytope $|P_{T_l}| \rightarrow M$ into a finite cylindrical end, in such a way that the barycenter behavior of the eventually coupled matter gets “distributed” over the full outer boundary $S_{\varepsilon(k)}^{(-)}$. Thus, in our description, we will follow the procedure outlined by Charpentier and Gawedzky in [82], which allows to specify the amplitudes on an arbitrary Riemann surface with boundaries and with an arbitrary specification of fields on them. In a general framework, let us consider a two-dimensional compact (not necessary connected) Riemann surface Σ with boundary $\partial\Sigma = \bigsqcup_{I=1}^N S_I$. If we parametrize the boundary loops $S_I \in \Sigma$ by analytical real maps $p_I : S^1 \rightarrow S_I$, then the field theoretical propagation amplitude over the Euclidean surface Σ is formally given by the following functional integral over the injection maps $X : \Sigma \rightarrow \mathcal{T}$:

$$\int_{\{X \circ p_I = X_I\}} \mathcal{D}X e^{-S[X]} \quad (2.38)$$

where $S[X]$ is the euclidean action of the bulk CFT, $\mathcal{D}X$ is the formal measure on the target space and the field X is restricted to take prescribed values X_I over the boundary loops S_I .

The rather abstract and formal expression (2.38) acquires a precise sense when we deal with a bosonic field defining a flat toroidal background. As a matter of fact, if we call Ξ the full set of possible boundary assignments, it is always possible to fix some real map X_{cl}

$$X_{cl} : \Sigma \rightarrow \mathbb{R} \quad (2.39)$$

which maps $\partial\Sigma$ diffeomorphically and with the prescribed orientation in Ξ (*i.e.* it exists a real map X_{cl} such that $X_{cl} \circ p_I = X_I$) and such that X_{cl} is an harmonic function w.r.t Δ_Σ , the Laplacian operator defined over Σ . Then, for a generic field, we can write:

$$X = X_{cl} + \tilde{X} \quad (2.40)$$

where X_{cl} is the harmonic function described above, while $\tilde{X} : \Sigma \rightarrow \mathbb{R}$ is the collection of the off-shell modes of X satisfying

$$\tilde{X} \circ p_I = 0 \quad (2.41)$$

This last condition implies the diagonal decomposition of the bulk action, $S[X] = S[X_{cl}] + S[\tilde{X}]$. Moreover, expanding $\tilde{X} = \sum c_M \tilde{X}_M$, where $\tilde{X}_M$ are the non zero modes of the Laplacian ($\Delta_\Sigma \tilde{X}_M = \Lambda_M \tilde{X}_M$, $\Lambda_M \neq 0 \forall M$), we finally get:

$$\mathcal{A}_\Sigma = \sum_{X_{cl}} e^{-S[X_{cl}]} \times \det' \left[\frac{8\pi^2}{N\Delta_\Sigma} \right]^{\frac{1}{2}} \quad (2.42)$$

where the prime means the exclusion of the zero mode, while N is the normalization of the Laplacian eigenfunctions. This last expression clearly diverges and needs to be regularized. If $\Sigma \equiv \Delta_{\varepsilon(k)}^*$ and N is the cylindrical end area, the standard Riemann Zeta-function regularization technique gives

$$\det' \left[\frac{8\pi^2 \theta(k)}{L^2(k) \Delta_\Sigma} \right]^{\frac{1}{2}} = \frac{1}{4\pi} \frac{1}{\sqrt{\text{Im}\tau} \eta(q)} \quad (2.43)$$

where is the Dedekind- η function with argument $q = e^{2\pi i\tau}$.

To define a suitable parametrization of the on-shell mode, let us switch to the transverse channel. The modular transformation $\tau \rightarrow -\frac{1}{\tau}$ redefines

$$\frac{1}{\sqrt{\text{Im}\tau} \eta(q)} = \frac{1}{\eta(\tilde{q})}. \quad (2.44)$$

The Laurent expansion of the classical mode $X^\alpha(\zeta(k), \overline{\zeta(k)})_{cl}$, $\alpha = 1, \dots, D$

on the annular region $\Delta_{\varepsilon(k)}^*$ is:

$$\begin{aligned} X^\alpha(\zeta(k), \bar{\zeta}(k))_{cl} &= X^\alpha(\zeta(k))_{cl} + X^\alpha(\bar{\zeta}(k))_{cl} \\ X^\alpha(\zeta(k))_{cl} &= \frac{X_0^\alpha}{2} + a^\alpha(k) \ln \zeta(k) + \sum_{n \neq 0} X_n^\alpha(k) \zeta^n(k) \\ X^\alpha(\bar{\zeta}(k))_{cl} &= \frac{X_0^\alpha}{2} + b^\alpha(k) \ln \bar{\zeta}(k) + \sum_{n \neq 0} \bar{X}_n^\alpha(k) \bar{\zeta}^n(k) \end{aligned} \quad (2.45)$$

Following a prescription introduced in [82] and specialized to the compactified boson in [66], it is possible to parametrize the classical field (2.45) in term of its restrictions to the boundaries $S_{\varepsilon(k)}^{(+)}$ and $S_{\varepsilon(k)}^{(-)}$:

$$\begin{aligned} X^\alpha(k)_{cl}|_+ &\doteq X^\alpha(\zeta(k), \bar{\zeta}(k))|_{S_{\varepsilon(k)}^{(+)}} = X_0^\alpha - \frac{2\pi}{2\pi - \varepsilon(k)} (a^\alpha(k) + b^\alpha(k)) + \\ &+ i\theta(k) (a^\alpha(k) - b^\alpha(k)) + \sum_{n \neq 0} b_n^\alpha(k) e^{in\vartheta(k)} \quad b_{-n}^\alpha(k) = \bar{b}_n(k) \end{aligned} \quad (2.46)$$

$$\begin{aligned} X^\alpha(k)_{cl}|_- &\doteq X^\alpha(\zeta(k), \bar{\zeta}(k))|_{S_{\varepsilon(k)}^{(-)}} = X_0^\alpha + \\ &+ i\theta(k) (a^\alpha(k) - b^\alpha(k)) + \sum_{n \neq 0} a_n^\alpha(k) e^{in\vartheta(k)} \quad a_{-n}^\alpha(k) = \bar{a}_n(k) \end{aligned} \quad (2.47)$$

where we have introduced the two sets of coefficients $\{a_n^\alpha\}$ and $\{b_n^\alpha\}$ having the following expression in terms of the Laurent's modes $X_n^\alpha(k)$ and $\bar{X}_n^\alpha(k)$:

$$a_n^\alpha \doteq e^{-\frac{2\pi}{2\pi - \varepsilon(k)} n} X_n^\alpha(k) + e^{\frac{2\pi}{2\pi - \varepsilon(k)} n} \bar{X}_{-n}^\alpha(k) \quad (2.48a)$$

$$b_n^\alpha \doteq X_n^\alpha(k) + \bar{X}_{-n}^\alpha(k) \quad (2.48b)$$

Redefining the compactification radius as $\Omega^\alpha(k) = \frac{R^\alpha(k)}{l(k)}$, the winding boundary conditions impose the consequent parametrization of the difference between the coefficients $a^\alpha(k)$ and $b^\alpha(k)$

$$a^\alpha(k) - b^\alpha(k) = -i\nu(k)\Omega^\alpha(k), \quad \nu(k) \in \mathbb{Z}, \quad (2.49)$$

The last term to parametrize is the difference between $X^\alpha(k)_{cl}|_+$ and $X^\alpha(k)_{cl}|_-$. This can be achieved [66] via an integer number $\mu(k) \in \mathbb{Z}$ and a real parameter $t^\alpha(k) \in (0, 2\pi\Omega^\alpha(k))$ [66]

$$a^\alpha(k) + b^\alpha(k) = t^\alpha(k) + 2\pi\mu(k)\Omega^\alpha(k). \quad (2.50)$$

Then, the classical solution is fully parametrized by the two set of complex numbers $\{a_n^\alpha\}$ and $\{b_n^\alpha\}$, obeying the reality conditions $a_{-n}^\alpha(k) = \bar{a}_n(k)$ and $b_{-n}^\alpha(k) = \bar{b}_n(k)$, the couple of integers $(\mu(k), \nu(k)) \in \mathbb{Z}^2$ and the real number

$$t^\alpha(k) \in (0, 2\pi\Omega^\alpha(k)][66]:$$

$$\begin{aligned} X^\alpha(\zeta(k), \bar{\zeta}(k))_{cl} &= X_0^\alpha + \\ &\left[\frac{t^\alpha(k)}{2} + \pi\mu(k)\Omega^\alpha(k) - i\frac{\nu(k)}{2}\Omega^\alpha(k) \right] \ln \zeta(k) + \\ &\left[\frac{t^\alpha(k)}{2} + \pi\mu(k)\Omega^\alpha(k) + i\frac{\nu(k)}{2}\Omega^\alpha(k) \right] \ln \bar{\zeta}(k) + \\ &\sum_{n \neq 0} \frac{a_n^\alpha(k)}{\tilde{q}^n - \tilde{q}^{-n}} (\zeta^n - \bar{\zeta}^{-n}) + \sum_{n \neq 0} \frac{b_n^\alpha(k)}{\tilde{q}^{-n} - \tilde{q}^n} \left(\left(\frac{\zeta}{\tilde{q}} \right)^n - \left(\frac{\bar{\zeta}}{\tilde{q}} \right)^{-n} \right) \end{aligned} \quad (2.51)$$

With these parametrization the “classical term” of the full amplitude will be completely determined in terms of the boundary values of the Laurent expansion of the classical configuration, modulo the pair of integers $(\mu(k), \nu(k))$ which parametrizes the coefficients $a^\alpha(k)$ and $b^\alpha(k)$. Thus, the sum over all the independent “classical configurations” in equation (2.42) is worked out summing over all values of the couple of integers $(\mu(k), \nu(k))$. This leads to the further factorization[66]:

$$\sum_{(\mu, \nu)} e^{-S[X^\alpha(\zeta(k), \bar{\zeta}(k))_{cl}]} = \mathcal{A}_{cl}(\{a_n^\alpha(k)\} \{b_n^\alpha(k)\}) \sum_{(\mu, \nu)} \mathcal{A}_{cl}(\mu^\alpha(k), \nu^\alpha(k), t^\alpha), \quad (2.52)$$

where:

$$\begin{aligned} \mathcal{A}_{cl}(\{a_n^\alpha(k)\} \{b_n^\alpha(k)\}) &= \\ &\prod_{n > 0} \exp \left[-\frac{k}{8} \left(a_n^\alpha(k) a_{-n}^\alpha(k) \frac{1 + \tilde{q}^{2n}}{1 - \tilde{q}^{2n}} - a_{-n}^\alpha(k) b_n^\alpha(k) \frac{2\tilde{q}^n}{1 - \tilde{q}^{2n}} + \right. \right. \\ &\quad \left. \left. - a_n^\alpha(k) b_{-n}^\alpha(k) \frac{2\tilde{q}^n}{1 - \tilde{q}^{2n}} + b_n^\alpha(k) b_{-n}^\alpha(k) \frac{1 + \tilde{q}^{2n}}{1 - \tilde{q}^{2n}} \right) \right] \end{aligned} \quad (2.53)$$

and

$$\sum_{(\mu, \nu)} \mathcal{A}_{cl}(\mu(k), \nu(k), t^\alpha) = \sum_{(\mu, \nu)} e^{i\frac{\mu^\alpha}{\Omega^\alpha} t^\alpha} \tilde{q}^{\frac{1}{4} \left(\frac{\mu(k)^2}{\Omega(k)^2} + \frac{1}{2} \nu(k)^2 \Omega(k)^2 \right)} \quad (2.54)$$

The full amplitude is then

$$\begin{aligned} \mathcal{A}(\{a_n^\alpha(k)\} \{b_n^\alpha(k), t^\alpha(k)\}) &= \frac{1}{\eta(\tilde{q})} \mathcal{A}_{cl}(\{a_n^\alpha(k)\} \{b_n^\alpha(k)\}) \\ &\sum_{(\mu, \nu)(k)} e^{i\frac{\mu^\alpha}{\Omega^\alpha} t^\alpha} \tilde{q}^{\frac{1}{4} \left(\frac{\mu(k)^2}{\Omega(k)^2} + \frac{1}{2} \nu(k)^2 \Omega(k)^2 \right)} \end{aligned} \quad (2.55)$$

2.3 Generalized Langlands boundary states and their conformal properties

On the annular domain $\Delta_{\varepsilon(k)}^*$ boundaries are concentric circles centered on the origins. In radial quantization, boundary conditions are imposed by propagating states from an initial (coherent) state $|\mathbf{r}_{(\mu,\nu)}^\alpha(S_{\varepsilon(k)}^{(-)})\rangle$ residing on the inner boundary $S_{\varepsilon(k)}^{(-)}$ to a final (coherent) boundary state $|\mathbf{r}_{(\mu,\nu)}^\alpha(S_{\varepsilon(k)}^{(+)})\rangle$ naturally associated to the outer boundary $S_{\varepsilon(k)}^{(+)}$. The Hamiltonian generating this translation is $H^{(C)} = \frac{2\pi}{L(k)} (L_0 + \bar{L}_0 + \frac{c}{12})$, while the propagator is $\tilde{q}^{(L_0 + \bar{L}_0 + \frac{c}{12})}$. Therefore, the amplitude on the cylindrical end can be formally written as:

$$\mathcal{A}(\{a_n^\alpha(k)\} \{b_n^\alpha(k)\}, t^\alpha(k)) = \sum_{(\mu,\nu)(k)} \left\langle \mathbf{r}_{(\mu,\nu)}^\alpha(S_{\varepsilon(k)}^{(+)}) \left| \tilde{q}^{L_0 + \bar{L}_0 - \frac{c}{12}} \right| \mathbf{r}_{(\mu,\nu)}^\alpha(S_{\varepsilon(k)}^{(-)}) \right\rangle \quad (2.56)$$

The main goal of [66] was to find a formal map which associates to each boundary assignation $\{(\mu, \nu), \{a_n\}\}$ (resp. $\{(\mu, \nu), \{b_n\}\}$) a well defined coherent state $|\mathbf{r}_{(\mu,\nu)}^\alpha(S_{\varepsilon(k)}^{(-)})\rangle$ (resp. $|\mathbf{r}_{(\mu,\nu)}^\alpha(S_{\varepsilon(k)}^{(+)})\rangle \in \mathcal{H}_{\mu,\nu} \otimes \bar{\mathcal{H}}_{\mu,\nu}$.

The procedure, on one hand, exploits the exact correspondence between the couple of integer parameters of the classical part of the partition function and the integers which parametrize the left and right zero modes of the compactified bosonic field, while on the other hand it introduces a couple of functions $A_{m_1, m_2}^n(a_n, a_{-n})$ and $B_{m_1, m_2}^n(b_n, b_{-n})$ of only two parameters, whose explicit expression can be extracted rewriting $\mathcal{A}_{cl}(\{a_n(k)\} \{b_n(k)\})$ in the form[66]:

$$\mathcal{A}_{cl}(\{a_n(k)\} \{b_n(k)\}) = \prod_{n=1}^{\infty} \sum_{m_1, m_2} B_{m_1, m_2}^n \tilde{q}^{n(m_1+m_2)} A_{m_1, m_2}^n \quad (2.57)$$

Let us introduce the following generalization of Langlands boundary states:

$$\begin{aligned} |\mathbf{r}_{(\mu,\nu)}^\alpha(S_{\varepsilon(k)}^{(-)})\rangle &= e^{it_-^\alpha(\lambda_{(\mu,\nu)}^\alpha + \bar{\lambda}_{(\mu,\nu)}^\alpha)} \times \\ &\times \prod_{n=1}^{\infty} \sum_{m_1, m_2} A_{m_1, m_2}^n(a_n^\alpha, a_{-n}^\alpha) \frac{(\mathbf{a}_{-n}^\alpha)^{m_1} \otimes (\bar{\mathbf{a}}_{-n}^\alpha)^{m_2}}{\sqrt{n^{m_1+m_2} m_1! m_2!}} |(\mu^\alpha, \nu^\alpha)\rangle \end{aligned} \quad (2.58)$$

and

$$\begin{aligned} |\mathbf{r}_{(\mu,\nu)}^\alpha(S_{\varepsilon(k)}^{(+)})\rangle &= e^{it_+^\alpha(\lambda_{(\mu,\nu)}^\alpha + \bar{\lambda}_{(\mu,\nu)}^\alpha)} \times \\ &\times \prod_{n=1}^{\infty} \sum_{m_1, m_2} B_{m_1, m_2}^n(b_n^\alpha, b_{-n}^\alpha) \frac{(\mathbf{a}_{-n}^\alpha)^{m_1} \otimes (\bar{\mathbf{a}}_{-n}^\alpha)^{m_2}}{\sqrt{n^{m_1+m_2} m_1! m_2!}} |(\mu^\alpha, \nu^\alpha)\rangle \end{aligned} \quad (2.59)$$

with $t_-^\alpha - t_+^\alpha = t^\alpha$ and

$$\begin{aligned} A_{m_1, m_2}^n(a_n^\alpha, a_{-n}^\alpha) &= e^{i\pi s(m_1+m_2)} \times \\ &\times \begin{cases} (2i\sqrt{n}a_n^\alpha)^{m_1-m_2} \sqrt{\frac{m_2!}{m_1!}} e^{-2n|a_n^\alpha|^2} L_{m_2}^{(m_1-m_2)}(4n|a_n^\alpha|^2) & m_1 \geq m_2 \\ (2i\sqrt{n}a_n^\alpha)^{m_2-m_1} \sqrt{\frac{m_1!}{m_2!}} e^{-2n|a_n^\alpha|^2} L_{m_1}^{(m_2-m_1)}(4n|a_n^\alpha|^2) & m_2 \geq m_1 \end{cases} \end{aligned} \quad (2.60)$$

with $L_{m_2}^{(m_1-m_2)}(\circ)$ the m_2 -th Laguerre polynomial of the (m_1-m_2) -th kind. By replacing the a_n^α with the b_n^α , and by conjugation (induced by the orientation of the boundary), we get also $\mathcal{B}_{m_1, m_2}^n(b_n^\alpha, b_{-n}^\alpha) = \overline{\mathcal{A}_{m_1, m_2}^n(b_{-n}^\alpha, b_n^\alpha)}$.

Let us stress that the above formula is a generalization of the original Langland's boundary states. As a matter of fact, the explicit expression in [66] did not include the phase factor $e^{i\pi s(m_1+m_2)}$ for the $\mathcal{A}_{m_1, m_2}^n$ and $\mathcal{B}_{m_1, m_2}^n$ coefficients. The introduction of such a phase factor is natural since these coefficients enter in the partition function via the combination in equation (2.57). Moreover, its presence is crucial if we want the boundary states to describe the most general kind of boundary condition.

From a macroscopic point of view, we can assume the presence of a brane emitting and adsorbing the boundary states via non-perturbative processes. In the same way, in the open string scheme, branes are those objects open strings endpoints lay on. Boundary conformal field theory allows to define branes in a microscopic CFT language, without any referring to target space geometry: the boundary CFT describes the perturbative properties of the open string-brane system. In this connection, the role of boundary states is obvious: they contain the complete informations about the static D-branes. However, as stressed in the introduction to this chapter, not every linear map such that one defining generalized Langlands boundary state actually describes an admissible boundary state, *i.e.* a coherent static D-brane configuration. To restrict the set of boundary states to an admissible one, we have to implement the glueing relations (2.14) and (2.15) directly on (2.58) and (2.59). Through radial quantization, *i.e.* mapping the cylinder into the annulus via (1.46), the (2.14) and (2.15) translates into

$$\zeta^2 T(\zeta) = \bar{\zeta}^2 \bar{T}(\bar{\zeta})|_{|\zeta|=1, |\zeta|=\bar{q}} \quad \zeta^{h_w} W(\zeta) = \bar{\zeta}^{\bar{h}_w} \bar{W}(\bar{\zeta})|_{|\zeta|=1, |\zeta|=\bar{q}} \quad (2.61)$$

Introducing the Laurent expansion of involved fields, we get the following conditions applied to boundary states:

$$\left(W_n - (-1)^{\bar{h}_w} \Omega(\bar{W}_{-n}) \right) \|B\rangle\rangle = 0 \quad (L_n - \bar{L}_{-n}) \|B\rangle\rangle = 0 \quad (2.62)$$

where $\|s\rangle\rangle$ is a generic conformal boundary state.

In our connection, the only chiral fields involved besides Virasoro fields are the holomorphic and antiholomorphic currents $J(\zeta)$ and $\bar{J}(\bar{\zeta})$ generating the Heisenberg algebra (2.35). Being the current algebra abelian, the possible glueing maps reduces to:

$$J(\zeta) = \bar{J}(\bar{\zeta}) \quad (2.63a)$$

$$J(\zeta) = -\bar{J}(\bar{\zeta}) \quad (2.63b)$$

which are respectively mapped into

$$(\mathfrak{a}_n + \bar{\mathfrak{a}}_{-n}) \|B\rangle\rangle = 0, \quad (2.64a)$$

$$(\mathfrak{a}_n - \bar{\mathfrak{a}}_{-n}) \|B\rangle\rangle = 0. \quad (2.64b)$$

The Sugawara construction (2.6) ensures that this conditions are sufficient to enforce conformal invariance encoded in (2.14).

With a careful computation, which exploits recursion relation of Laguerre polynomials, and which is reported in appendix A, we reduced generalized Langlands boundary states to the Dirichlet and Neumann Ishibashi states associated with the free bosonic field $X^\alpha(k)$:

$$\left| \mathfrak{r}_\mu^\alpha(S_{\varepsilon(k)}^{(-)}) \right\rangle^{(D)} = e^{\sqrt{-1}t_+^\alpha \mu^\alpha \frac{L}{R^\alpha}} \exp \left(\sum_{n=1}^{\infty} \frac{1}{n} (\mathfrak{a}_{-n}^\alpha) (\bar{\mathfrak{a}}_{-n}^\alpha) \right) |(\mu^\alpha, 0)\rangle, \quad (2.65)$$

and

$$\left| \mathfrak{r}_\mu^\alpha(S_{\varepsilon(k)}^{(-)}) \right\rangle^{(N)} = \exp \left(\sum_{n=1}^{\infty} -\frac{1}{n} (\mathfrak{a}_{-n}^\alpha) (\bar{\mathfrak{a}}_{-n}^\alpha) \right) |(0, \nu^\alpha)\rangle. \quad (2.66)$$

The most general boundary state will be a superposition of these states, weighted by suitable coefficients:

$$\| \mathfrak{r}_\mu^\alpha(S_{\varepsilon(k)}^{(-)}) \rangle\rangle^{(D)} = \sum_{\mu \in \mathbb{Z}} \mathbf{C}_\mu \left| \mathfrak{r}_\mu^\alpha(S_{\varepsilon(k)}^{(-)}) \right\rangle^{(D)} \quad (2.67a)$$

$$\| \mathfrak{r}_\mu^\alpha(S_{\varepsilon(k)}^{(-)}) \rangle\rangle^{(N)} = \sum_{\nu \in \mathbb{Z}} \mathbf{C}_\nu \left| \mathfrak{r}_\nu^\alpha(S_{\varepsilon(k)}^{(-)}) \right\rangle^{(N)} \quad (2.67b)$$

The glueing automorphism Ω in (2.15) is sufficient to determine such coefficients uniquely. Strong constraints on their expression derive directly by the relation (2.21) relating coefficients of bulk-fields one-point function. To show how this works, let us introduce a more compact notation. Let us omit the vector index α . Moreover let us introduce:

- at fixed glueing map Ω , let $|I_{\lambda(p)}\rangle\rangle_\Omega$ be the collection of all the Ishibashi states which are built on the primary states of the Hilbert space of the BCFT defined over $\Delta_{\varepsilon(p)}^*$;
- let Y be the collection of all the indexes $\lambda(p)$ labeling such Ishibashi states: obviously, $Y \subseteq \mathcal{Y}$, where $\mathcal{Y} = \{(\mu, \nu)\}$ is the collection of labeling associated to the irreducible representations of $u_L(1) \times u_R(1)$;
- let us denote with $\mathcal{A} = \{A(p)\}$ the set of all possible boundary assignments, including both the glueing condition and the dependence from other parameters, which we include in a collective index α . Thus, each $A(p) \in \mathcal{A}$ will be a pair $A(p) = (\Omega, \alpha)^5$;

⁵We do not assign a polytope index to the sets of labels Y , $\mathcal{Y}$ and $\mathcal{A}$ since the BCFT constructed is the same over each cylindrical ends, thus these sets coincide over each $\Delta_{\varepsilon(p)}^*$

- let us indicate with $\|\mathfrak{r}_{(p)}(\alpha)\rangle\rangle^\Omega$ the collection of all the admissible boundary states on the boundary $S_{\varepsilon(p)}^{(-)}$ of $\Delta_{\varepsilon(p)}^*$.

With the statements above, let us write the relation between the boundary states and the Ishibashi states as:

$$\|\mathfrak{r}_{(p)}(\alpha)\rangle\rangle^\Omega = \sum_{\lambda(p) \in Y} B_{A(p)}^{\lambda(p)} |I_{\lambda(p)}\rangle\rangle^\Omega, \quad (2.68)$$

$B_{A(p)}^{l(p)}$ being the $|I_{l(p)}\rangle\rangle$ projection on $\|\mathfrak{r}_{(p)}(\alpha)\rangle\rangle^\Omega$.

Coefficients $B_{A(p)}^{\lambda(p)}$ are strictly related both to bulk fields one-point function coefficients $A_{\lambda(p)}^{A(p)}$ in (2.20) and to coefficients $C_{\phi(p)\lambda(p)}^{A(p)}$ entering into the bulk-to-boundary expansion (2.26) [74, 78, 75]. In particular, exploiting transformation properties of correlators under the conformal mapping relating the finite cylinder to the annulus, one can show [78]:

$$B_{A(p)}^{\lambda(p)} = A_{\lambda(p)}^{A(p)} \quad (2.69)$$

Replacing this identity into (2.21), we get the constraint over the boundary states' coefficients⁶:

$$B_{A(p)}^{\lambda(p)} \cdot B_{A(p)}^{\lambda'(p)} = B_{A(p)}^{\lambda(p)+\lambda'(p)} \quad (2.70)$$

from which we get the boundary states:

$$\|\mathfrak{r}^\alpha(S_{\varepsilon(k)}^{(-)})\rangle\rangle^{(D)} = \frac{1}{\sqrt{2\frac{L(k)}{R^\alpha(k)}}} \sum_{\mu \in \mathbb{Z}} e^{it_+^\alpha \mu^\alpha \frac{L}{R^\alpha}} \exp\left(\sum_{n=1}^{\infty} \frac{1}{n} (\mathfrak{a}_{-n}^\alpha) (\bar{\mathfrak{a}}_{-n}^\alpha)\right) |(\mu^\alpha, 0)\rangle, \quad (2.71)$$

i.e. the Dirichlet boundary state describing a boundary loop lying on a D-brane positioned at $X^\alpha|_+ = t_+$, and the states

$$\|\mathfrak{r}^\alpha(S_{\varepsilon(k)}^{(-)})\rangle\rangle^{(N)} = \sqrt{\frac{L(k)}{R^\alpha(k)}} \sum_{\nu \in \mathbb{Z}} e^{\frac{i}{2}\tilde{t}_+^\alpha \nu^\alpha \frac{R^\alpha}{L}} \exp\left(-\sum_{n=1}^{\infty} \frac{1}{n} (\mathfrak{a}_{-n}^\alpha) (\bar{\mathfrak{a}}_{-n}^\alpha)\right) |(0, \nu^\alpha)\rangle, \quad (2.72)$$

i.e. a Neumann boundary states describing a boundary loop moving in the associated direction. In the above formula, $\tilde{t}_+$ is the $U(1)$ holonomy naturally associated to the boundary loop.

⁶This is the specialized version of the sewing constraint holding for the projections of boundary states $\|\alpha\rangle\rangle$ of a Rational Conformal Field Theory on the associated Ishibashi states $|j\rangle\rangle$:

$$B_\alpha^i B_\alpha^j = \sum_k C_{ij}^k F_{k1} \begin{bmatrix} j & j \\ i & i \end{bmatrix} B_\alpha^k,$$

where C_{ij}^k and $F_{k1} \begin{bmatrix} j & j \\ i & i \end{bmatrix}$ are respectively the fusion coefficients and fusion matrices of the bulk theory

Chapter 3

Boundary Insertion Operators

To complete the description of the BCFT on the full Riemann Surface, we need to discuss how the N_0 distinct copies of the theory, each one living on a different cylindrical end $\Delta_{\varepsilon(k)}^*$, interact along the underlying polytope $|P_{T_l}| \rightarrow M$. In [67], a remarkable intuition proposed that this interaction could be mediated by boundary conditions changing operators, whose presence is predicted by boundary conformal field theory [79, 78] (see the general theory presented in the previous chapter).

In this chapter starting from this statement, we slightly modify it. We describe interaction on the ribbon graph introducing, beside ordinary boundary condition changing operators, a new class of operators which mediate the change in boundary conditions which actually take place non locally on the boundary shared by two adjacent cylindrical ends. As a matter of fact, while in ordinary boundary conformal field theory a jump in boundary conditions on the boundary is explained by the local action of boundary operators, this feature does not fit completely our model, both because, after the glueing process we do not deal with a true boundary, but with a “separation edge” between the two adjacent cylindrical ends, and because we do not have a jump in boundary conditions taking place in a precise point of this edge (which we will go on calling boundary, for the sake of simplicity) but with two different boundary conditions which, in the adjacency limit, coexist on this shared boundary. From these considerations, it follows the description of the dynamical glueing of the $N_0(T)$ copies of the bosonic BCFT which we present in this chapter.

3.1 Boundary Insertion Operators

In the background defined in section 2.1.1, it may happen that a boundary condition changes along the real line. In radial quantization, this situation is explained with the presence of a vacuum which is no longer invariant under




Figure 3.1: The insertion of boundary operators on the shared boundary between pairwise adjacent polytope allows to define the superposition of different boundary conditions[67]

the action of $L_{-1}^{(H)}$. In [79] it was proposed that such states were obtained by the action of a boundary operator on the true vacuum, *i.e.* it may happen that boundary operators induce transitions on boundary conditions along the boundary. These boundary condition changing operators are associated with vectors in the Hilbert space depending on both the boundary conditions, and they cannot be obtained from bulk fields through a bulk to boundary OPE.

In this section we will propose a slight modification of this picture, introducing a particular class of boundary condition changing operators which, living on the boundary shared between two adjacent polytopes and carrying an irreducible action of the chiral algebra, will mediate the change in boundary conditions between pairwise adjacent cylindrical ends.

To fully understand this statement, let us first analyze the geometry on the neighborhood of a shared ribbon graph edge. Ribbon graph's strips represent the uniformization of the singular structure of the RRT in the neighborhood of the 1-skeleton of the original triangulation. The ribbon graph arises as a consequence of the presence of a locally uniformizing coordinate $\zeta(k)$, which provides a counterclockwise orientation in the 2-cells of the original triangulation: such an orientation gives rise to a cyclic ordering on the set of the half edges $\{\rho^1(k)^\pm\}_{k=0}^{N_1(T)}$ incident on the vertexes $\{\rho^0(k)^\pm\}_{k=0}^{N_2(T)}$, stating a 1-to-1 correspondence between the 1-skeleton of the original triangulation and the set of trivalent metric ribbon graphs. In this connection, let us consider two adjacent cylindrical ends $\Delta_{\varepsilon(p)}^*$ and $\Delta_{\varepsilon(q)}^*$: these are dual to the adjacent cells $\rho^2(p)$ and $\rho^2(q)$. The cylindrical ends are glued to the oriented boundaries $\partial\Gamma_p$

and $\partial\Gamma_q$ of the ribbon graph, which are denoted respectively by $\partial\Gamma_p$ and $\partial\Gamma_q$. Let us consider the oriented strip associated with the edge $\rho^1(p, q)$ of the triangulation and its uniformized neighborhood $(U_{\rho^1(p, q)}, z(p, q))$ and let us take in $(U_{\rho^1(p, q)}, z(p, q))$ two infinitesimally neighboring points $z_1 = x_1 + iy_1$ and $z_2 = x_2 + iy_2$ such that $x_1 = x_2$. Thus, if $y_1 \rightarrow 0$ we are approaching the intersection between the boundary $\partial\Gamma_p$ of the ribbon graph and the oriented edge $\rho^1(p, q)$, while when $y_2 \rightarrow 0$ we are approaching the intersection between the boundary $\partial\Gamma_q$ of the ribbon graph and the oriented edge $\rho^1(q, p)$. From a BCFT point of view, this leads to the conclusion that, in the adjacency limit described above, the “effective boundary” between two adjacent cylindrical end is unique and shares the two different boundary conditions associated to $S_{\varepsilon(p)}^{(-)}$ and $S_{\varepsilon(q)}^{(-)}$.

In this connection, it is no longer correct to argue the presence of vacuum state which is not invariant under translations along the boundary. The shared boundary is obtained glueing two loops, each of them being part of a domain, on which we have defined a self-contained BCFT, with its associated Hilbert space and vacuum state invariant under translation along the boundary loop.

Thus, in order the glueing process to take place coherently, we have to require that each state of the Hilbert space melt without breaking the conformal and chiral symmetry of the model. This leads to the definition of Boundary Insertion Operators.

A CFT is consistently defined on each cylindrical end once we have imposed constraints in (2.61). In particular, specializing to the Virasoro fields, on $\Delta_{\varepsilon(p)}^*$ inner boundary we have

$$\zeta(p)^2 T(\zeta(p)) = \bar{\zeta}(p)^2 \bar{T}(\bar{\zeta}(p))|_{|\zeta(p)|=1} \quad (3.1)$$

while in the inner boundary of $\Delta_{\varepsilon(q)}^*$ it holds:

$$\zeta(q)^2 T(\zeta(q)) = \bar{\zeta}(q)^2 \bar{T}(\bar{\zeta}(q))|_{|\zeta(q)|=1} \quad (3.2)$$

This conditions allows to combine, on each cylindrical end, T and $\bar{T}$ in a unique object (well defined conformally mapping the cylinder into a semi annular domain in the UHP), thus allowing to associate to each cylindrical end a single copy of the Virasoro algebra (the same obviously holds for the other chiral fields, see eq. (2.17)).

In this connection, we can pursue further this last construction, and implement a non symmetry-breaking glueing of two adjacent cylindrical end by associating, to each pairwise adjacent couple of them, a unique copy of both Virasoro and chiral algebras. To achieve this, let us consider the neighborhood of the (p, q) edge defined in (1.26) and uniformized with the complex coordinate $z(p, q)$. Conformal mappings between $z(p, q)$ and the coordinates uniformizing




Figure 3.2: A small integration contour intersecting the (p, q) edge of the ribbon graph

each cylindrical end are define as:

$$\zeta(p) = e^{\frac{2\pi i}{L(p)}(L(p) - \hat{L}(p, q) + z(p, q))} \quad \zeta(q) = e^{\frac{2\pi i}{L(q)}(L(q) - \hat{L}(q, p) + z(q, p))} \quad (3.3)$$

where it holds $z(q, p) = -z(p, q)$. We can easily express the holomorphic and antiholomorphic components of the Virasoro fields defined on each cylindrical end in term of the strip coordinate. For the holomorphic components we have:

$$T_{(p)}(z(p, q)) = T_{(p)}(\zeta(p)) \left(\frac{dz(p, q)}{d\zeta(p)} \right)^{-2} \quad (3.4a)$$

$$T_{(q)}(z(q, p)) = T_{(q)}(\zeta(q)) \left(\frac{dz(q, p)}{d\zeta(q)} \right)^{-2}. \quad (3.4b)$$

Analogous relations hold for the antiholomorphic sector.

Rescaling $z(q, p) = -z(p, q)$, then we perform the glueing asking for the following relations to hold:

$$T_{(p)}(z(p, q)) = T_{(q)}(z(p, q))|_{y(p, q)=0} \quad (3.5a)$$

$$\overline{T}_{(p)}(\bar{z}(p, q)) = \overline{T}_{(q)}(\bar{z}(p, q))|_{y(p, q)=0} \quad (3.5b)$$

These relations allows to associate to each pairwise adjacent couples of theories a single copy of the Virasoro algebra. As a matter of fact, these glueing conditions allows to define a unique holomorphic component of the stress energy-tensor as:

$$T_{(p, q)} = \begin{cases} T_{(p)}(z(p, q)) & \text{if } z(p, q) \cup \zeta(p) \neq 0 \\ T_{(q)}(z(q, p)) & \text{if } z(q, p) \cup \zeta(q) \neq 0 \end{cases} \quad (3.6)$$

The same holds for the antiholomorphic sector, allowing to define a unique $\overline{T}(\bar{z}(p, q))$. Moreover, $T_{(p, q)}$ and $\overline{T}(\bar{z}(p, q))$ are not independent, because of relations (3.1) (3.2). Thus, to each pairwise adjacent couples of cylindrical ends, we can associate a unique copy of the Virasoro algebra. This can be defined considering a small integration contour crossing the (p, q) boundary, like that at the rhs of fig. 3.2

Thanks to the continuity condition along the boundary, we can actually

write:

$$\begin{aligned}
L_n^{(p,q)} = & \frac{1}{2\pi i} \oint_{C(p,q)} z(p,q)^{n+1} T_{(p,q)}(z(p,q)) = \\
& \frac{1}{2\pi i} \oint_{C(p)} dz(p,q) z(p,q)^{n+1} T_p(z(p,q)) + \\
& \frac{1}{2\pi i} \oint_{C(q)} dz(q,p) z(q,p)^{n+1} T_q(z(q,p)) \quad (3.7)
\end{aligned}$$

A similar reasoning can be performed for the other chiral fields of the model, but performing a particular attention at the gluing maps Ω in (2.16). As a matter of fact, we must keep in account the fact that this glueing process must relate the two glueing automorphisms $\Omega_{(p)}$ and $\Omega_{(q)}$ associated to the BCFTs defined respectively on $\Delta_{\varepsilon(p)}^*$ and $\Delta_{\varepsilon(q)}^*$.

With the above remarks, we can associate to each pairwise coupled BCFTs a unique Hilbert space carrying a representation of the $u(1)_{(p,q)}$ algebra. Consequently the associated collection of boundary operators will have a primary $u(1)$ (and Virasoro) index $\lambda(p,q)$. We ask the pair of polytope indexes (p,q) to be ordered, assuming that this means that the operator in question lives on the “ p -side” of the boundary of the strip connecting $\Delta_{\varepsilon(p)}^*$ with $\Delta_{\varepsilon(q)}^*$. Moreover, these operators carry the information about the conformal boundary condition over $S_{\varepsilon(p)}^{(-)}$, thus they are labeled by another index $A(p)$:

$$\psi_{\lambda(p,q)}^{A(p)}(x(p,q)), \quad (3.8)$$

where $x(p,q) = \Re z(p,q)$.

In the same way, we can define with

$$\psi_{\lambda(q,p)}^{B(q)}(x(q,p)) \quad (3.9)$$

(with $x(q,p) = \Re z(q,p)$) the associated operator, transformed under the conformal mapping $z(p,q) = -z(q,p)$. According to the above assumptions, operators in (3.9) live on the q -side of the (p,q) edge, thus carrying the information about the boundary condition on $S_{\varepsilon(q)}^{(-)}$.

However, the above distinction between p -side fields and q -side ones is merely artificial, since the spectrum of boundary operators associated to a given edge is unique. The natural consequence of this is that each of them must carry the information about the boundary conditions adopted on both $S_{\varepsilon(p)}^{(-)}$ and $S_{\varepsilon(q)}^{(-)}$.

Thus the complete labeling of Boundary Insertion Operators (BIO) which arises from the BCFT defined on the (p,q) -edge is


$$\psi_{\lambda(p,q)}^{B(q)A(p)}(x(p,q)); \quad (3.10)$$

In this connection, these objects naturally mediate the change in boundary conditions between pairwise adjacent boundaries. The labels ordering, indicates

the “direction” in which BIO acts, with the understanding that:

$$\psi_{A(p)B(q)}^{\lambda(p,q)}(x(q,p)) = \psi_{\lambda(p,q)}^{B(q)A(p)}(x(p,q)); \quad (3.11)$$

At this level of the discussion, the action of BIO is still purely formal. Since the switch in boundary conditions (from $A(p)$ to $B(q)$ and viceversa) will act both on the glueing automorphism and on the Wilson line/brane position (see remarks before equation (2.68)), we can relate their action to an automorphism of the algebra relating the two glueing maps $\Omega_{(p)}$ and $\Omega_{(q)}$ (see equation (2.16)). Moreover, BIOs are chiral primary operators of the $U(1)$ algebra: their expression suggests to adopt a description analogue to that one introduced in [83] for usual boundary operators in Rational Conformal Field Theory (RCFT). They can be represented as Chiral Vertex Operators (CVO) which, in every point of the boundary, map the coupling between states defined by the boundary condition $A(p)$ and states belonging to the $\lambda(p,q)$ representation of the chiral algebra to the spectrum of states which defines the boundary condition $B(q)$. We can introduce a pictorial description of the interaction as represented by the (non local, because the switch between boundary condition take place along the full boundary) interaction vertex:



This kind of interaction is equivalent to the usual chiral vertex which represent the map from the fusion of two representation of the (bulk) Virasoro algebra $\lambda(p)$ and $\lambda'(p)$ to the representation $\lambda''(p)$.

The analogy stated with the usual CVO formalism is purely formal, since the “fusion” process described above involve different objects like boundary states (which are superposition of Ishibashi states) and irreducible representations of the chiral algebra. However, this analogy allows to specify the conformal properties of these operators. It is possible, a priori, to assign a multiplicity index to BIOs, $t = 1, \dots, N_{\lambda(p,q)}^{A(p)B(q)}$ for each $\psi_{A(p)B(q)}^{\lambda(p,q)}(x)$ which takes into account the degeneration which can occur in the map process described before. Moreover, BIO are primary operators of the boundary chiral algebra, thus their conformal dimension coincide with the highest weight of the $V_{\lambda}(p,q)$ module of the $U(1)$ (Virasoro) algebra:

$$H(p,q) = \frac{1}{2}\lambda(p,q)^2 \quad (3.12)$$

BIOs live on the ribbon graph, thus their interactions are guided by the

trivalent structure of Γ . A careful and exhaustive analysis, which has been entirely reported in appendix B, shows that this structure allows to define all fundamental coefficients weighting the self-interaction of boundary insertion operators. Two points functions are well defined on the edges of the graph, while OPE coefficients naturally defines the fusion of different boundary insertion operators interacting in $N_2(T)$ tri-valent vertexes of Γ . Moreover, we have shown that thanks to the trivalent structure of the ribbon graph, we can define a set of sewing constraint these coefficients must satisfy. Remarkably, these constraints are perfectly analogues to the sewing constraint introduced in [73] for boundary conditions changing operator, thus enforcing the analogy between these latters and BIO, which merge dynamically the $N_0(T)$ copies of the BCFT which are defined on M_∂ .

3.2 Investigations at the T-duality self dual radius

The above description can be pursued further if we explore the rational limit of the compactified boson BCFT, *i.e.* the limit

$$\Omega(k) \doteq \Omega(k)_{s.d.} = \frac{R(k)_{s.d.}}{l(k)} = \sqrt{2} \quad (3.13)$$

where $\Omega(k)_{s.d.}$ is the self-dual radius, *i.e.* the fixed point of T -duality transformation $\Omega \rightarrow \frac{1}{\Omega}$. If we substitute $\Omega = \sqrt{2}$ into (2.34), we obtain $\lambda_{s.d} = \frac{(\mu+\nu)}{\sqrt{2}}$ and $\bar{\lambda}_{s.d} = \frac{(\mu-\nu)}{\sqrt{2}}$, thus we are falling exactly in the situation described in section 2.2. The holomorphic and antiholomorphic conformal dimensions of primary fields are respectively $h_{s.d} = \left(\frac{\mu+\nu}{2}\right)^2$ and $\bar{h}_{s.d} = \left(\frac{\mu-\nu}{2}\right)^2$, *i.e.* they are the square of an integer or semi-integer number. In full generality, we can write $h(p) = j(p)^2$, $j(p) \in \frac{1}{2}\mathbb{Z}$. This lead to the presence of a null state at the level $2j+1$ in the correspondent $U(1)$ module, which consequently decompose into into an infinite set of Virasoro ones. Consequences of this can be understood considering the torus partition function associated the compactified boson with $\Omega = \Omega_{s.d.}$ ¹:

$$Z(\sqrt{2}) = \frac{1}{|\eta(\tau)|^2} \sum_{\mu, \nu \in \mathbb{Z}} q^{\frac{1}{4}(\mu+\nu)^2} \bar{q}^{\frac{1}{4}(\mu-\nu)^2} \quad (3.14)$$

If we redefine $m = \mu + \nu$ and $n = \mu - \nu$ such that $m - n = 0 \bmod 2$, (3.14) reads

$$Z(\sqrt{2}) = \frac{1}{|\eta(\tau)|^2} \sum_{\substack{m, n \in \mathbb{Z} \\ m-n=0 \bmod 2}} q^{\frac{1}{4}m^2} \bar{q}^{\frac{1}{4}n^2}. \quad (3.15)$$

¹In this discussion we will omit the polytope index: it will be restored when we will deal again whit the interaction of the N_0 copies of the BCFT on the ribbon graph

We can introduce the extended characters[80]:

$$C_0(\tau) = \frac{1}{\eta(\tau)} \sum_{m \text{ even}} q^{\frac{1}{4}m^2} = \frac{1}{\eta(\tau)} \sum_{m \in \mathbb{Z}} q^{m^2} = \frac{\theta_3(2\tau)}{\eta(\tau)} \quad (3.16a)$$

$$C_1(\tau) = \frac{1}{\eta(\tau)} \sum_{m \text{ odd}} q^{\frac{1}{4}m^2} = \frac{1}{\eta(\tau)} \sum_{m \in \mathbb{Z}} q^{(m+\frac{1}{2})^2} = \frac{\theta_2(2\tau)}{\eta(\tau)} \quad (3.16b)$$

where $\theta_i(\tau)$ is the i -th Jacobi Theta function. The partition function now reads:

$$Z(\sqrt{2}) = |C_0(\tau)|^2 + |C_1(\tau)|^2. \quad (3.17)$$

We have actually reorganized the infinite Virasoro modules into a finite number of extended blocks which transforms linearly into each others by modular transformation. The extended $\mathcal{S}^{ext}$ matrix, which encodes the modular data via the relation $C_i(-\frac{1}{\tau}) = \mathcal{S}_{ij}^{ext} C_j(\tau)$, reads[80]:

$$\mathcal{S}^{ext} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \quad (3.18)$$

These extended blocks are identified with the two irreducible representations of the affine algebra $\hat{su}(2)_{k=1}$, namely the vacuum representation $\mathcal{H}_0$, generated from a state transforming in the singlet of the horizontal $su(2)$ algebra, and the representation $\mathcal{H}_{\frac{1}{2}}$, for which the highest weight state transform into the fundamental representation of the horizontal $su(2)$ algebra. Thus, when $\Omega(k) = \Omega(k)_{s.d.}$, the BCFT defined over each cylinder is equivalent to an $\hat{su}(2)_{k=1}$ -WZW model: it is an extended non-minimal model generating a Rational Conformal Field Theory (RCFT), a theory whose possibly infinite Verma modules can be reorganized into a finite number of extended blocks. These are irreducible representations of an extended symmetry algebra playing the role of chiral algebra for the underlining CFT.

In this connection, the set of labels associated to the irreducible representations of the chiral algebra is finite, $\mathcal{Y} = \{0, \frac{1}{2}\}$, and the Hilbert space is written as:

$$\mathcal{H}_{bulk} = \left(\mathcal{H}_0^{\hat{su}(2)} \otimes \overline{\mathcal{H}}_0^{\hat{su}(2)} \right) \oplus \left(\mathcal{H}_{\frac{1}{2}}^{\hat{su}(2)} \otimes \overline{\mathcal{H}}_{\frac{1}{2}}^{\hat{su}(2)} \right). \quad (3.19)$$

Since they will be useful in the following discussion, let us summarize the main features of the conformal field theory associated to the $\hat{su}(2)_{k=1}$ WZW model. $\hat{su}(2)_1$ (bulk) primary fields are[80]:

$$j = 0 \rightarrow \phi_{(0,0),(0,0)}(z, \bar{z}) = \mathbb{I} \quad (3.20a)$$

$$j = \frac{1}{2} \rightarrow \begin{cases} \phi_{(\frac{1}{2}, \frac{1}{2}), (\frac{1}{2}, \frac{1}{2})}(z, \bar{z}) = e^{\frac{i}{\sqrt{2}}X(z)} e^{\frac{i}{\sqrt{2}}X(\bar{z})} \\ \phi_{(\frac{1}{2}, -\frac{1}{2}), (\frac{1}{2}, -\frac{1}{2})}(z, \bar{z}) = e^{-\frac{i}{\sqrt{2}}X(z)} e^{-\frac{i}{\sqrt{2}}X(\bar{z})} \end{cases} \quad (3.20b)$$

The associated boundary theory is defined by a set of glueing conditions on the boundary (see formula (2.16)). Let us remember that, once we are given

a glueing automorphism, we can associate to each irreducible representation of the chiral algebra an Ishibashi state[84], *i.e.* a coherent state solution of the constraint (2.62). In this connection, let us denote the two Ishibashi states which are associated with the $\mathcal{H}_0^{\hat{su}(2)}$ and the $\mathcal{H}_{\frac{1}{2}}^{\hat{su}(2)}$ modules respectively with $|0\rangle\rangle$ and $|\frac{1}{2}\rangle\rangle$. A remarkable property defined by Cardy in [79] states that the set of boundary conditions of a rational boundary conformal field theory are labelled exactly by the modules of the chiral algebra, while the correspondent boundary states are given by[79]:

$$|A(p)\rangle\rangle = \sum_{l \in \mathcal{Y}(p)} \frac{\mathcal{S}_{Al}^{ext}}{\sqrt{\mathcal{S}_{0l}^{ext}}} |l(p)\rangle\rangle \quad (3.21)$$

In the case of $\hat{su}(2)_{k=1}$ -WZW model, we can write $\mathcal{Y} \equiv \mathcal{A} = \{0, \frac{1}{2}\}$, while formula (3.21) specializes to:

$$|J\rangle\rangle = 2^{-\frac{1}{4}} |0\rangle\rangle + (-1)^{2J} 2^{-\frac{1}{4}} |\frac{1}{2}\rangle\rangle, \quad \text{with } J = 0, \frac{1}{2}. \quad (3.22)$$

These properties allow for a complete definition of BIO of a rational conformal field theory. As a matter of fact, since both boundary states and boundary operators are identified by primary labels of the extended chiral algebra, in analogy to what happens for boundary conditions changing operators of rational minimal models, we can define the glueing process as the fusion between the representations associated to the two adjacent boundary states and the one BIO carries. Thus, let us consider the (p, q) edge $\rho^1(p, q)$. If $|J(p)\rangle\rangle$ and $|J(q)\rangle\rangle$ are the boundary conditions shared by $\rho^1(p, q)$, BIOs on $\rho^1(p, q)$ are defined as:

$$\psi_{j(p,q)}^{J(p)J(q)}(x(p, q)) = \mathcal{N}_{J(p)j(p,q)}^{J(q)} \psi_{j(p,q)}(x(p, q)) \quad (3.23)$$

i.e. they are the $\hat{su}(2)_{k=1}$ primary fields weighted by the fusion rule $N_{J(p)j(p,q)}^{J(q)}$. The latters are given by a combination of the $\mathcal{S}^{ext}$ matrix entries via the Verlinde formula

$$\mathcal{N}_{J(p)j(p,q)}^{J(q)} = \sum_{l \in \mathcal{Y}} \frac{\mathcal{S}_{J(p)l}^{ext} \mathcal{S}_{j(p,q)l}^{ext} \overline{\mathcal{S}}_{lJ(q)}^{ext}}{\mathcal{S}_{0l}^{ext}} \quad (3.24)$$

As it stands, this construction, valid for the $\hat{su}(2)_{k=1}$ model, does not apply to the compactified boson at the self-dual radius $\Omega_{s.d.}$. As a matter of fact, even if the Hilbert space of the two models is the same, the boundary theory associated to the latter one is quite more complicated. It will lead to the definition of BIOs of the compactified boson at the self dual radius as truly marginal deformations of the operators in (3.23) by the action of $SO(3)$ elements.

We will dedicate the remaining part of this section to the demonstration of this statement.

Due to the Sugawara construction, each highest weight representation of $\hat{su}(2)_{k=1}$ is also an irreducible representation of the $c = 1$ Virasoro algebra,

whose generators commute with the horizontal $su(2)$ ones. Thus, we can decompose each irreducible representation of $\hat{su}(2)_{k=1}$ entering in (3.19) with respect to $su(2) \otimes Vir$:

$$\mathcal{H}_j^{\hat{su}(2)} = \sum_{n=0}^{\infty} \mathcal{V}_{(n+j)^2} \otimes \mathcal{H}_{n+j}^{su(2)}, \quad (3.25)$$

where $\mathcal{V}_h$ is the conformal weight h irreducible representation of the Virasoro algebra, while $\mathcal{H}_j^{\hat{su}(2)}$ (resp. $\mathcal{H}_j^{su(2)}$) is the $(2j+1)$ -dimensional representation of $\hat{su}(2)$ (resp. $su(2)$). The bulk Hilbert space then decompose as:

$$\mathcal{H}_{bulk} = \sum_{\substack{n, \bar{n}=0 \\ n+\bar{n} \text{ even}}}^{\infty} \left(\mathcal{V}_{\frac{n^2}{4}} \otimes \bar{\mathcal{V}}_{\frac{\bar{n}^2}{4}} \right) \otimes \left(\mathcal{H}_{\frac{n}{2}}^{su(2)} \otimes \bar{\mathcal{H}}_{\frac{\bar{n}}{2}}^{su(2)} \right). \quad (3.26)$$

As we outlined before, the representations of the Virasoro algebra which occur in the decomposition are all degenerates. Due to this, the set of Virasoro primary fields is larger than in the case of generic compactification radius. The decomposition (3.26) shows that, for fixed (m, n) , the $U(1)_L \times U(1)_R$ representation of momenta $\lambda_{s,d}$ and $\bar{\lambda}_{s,d}$ contains the Virasoro representation provided that both $|m|, |n| \leq j$ and $j-m$ and $j-n$ are integers. Conversely, if m and n satisfies these constraints, the representation $\mathcal{V}_{\frac{n^2}{4}} \otimes \bar{\mathcal{V}}_{\frac{\bar{n}^2}{4}}$ of the Virasoro algebra enter exactly once into the representation of $U(1) \times U(1)$. Thus the Virasoro primary fields can be labelled by the triple of indexes $(j; m, n)$, with the constraint $|m|, |n| \leq j$. This is the celebrated DISCRETE SERIE OF STATES[85, 86, 87, 88] $\psi_{j,m} \bar{\psi}_{j,n}$. The associated Ishibashi states are defined in literature as $|j, m, n\rangle\rangle$ [87].

The arising of the discrete serie of states is strictly related to the deformation of an open bosonic string theory obtained by adding suitable boundary operators to the action. As a matter of fact, authors of [87] showed that the discrete spectrum generated by the free boson once it is compactified at the T -duality self dual radius can be equivalently explained with an open string model in which one end of string is subjected to Dirichlet boundary conditions while, on the other end, the boundary condition is dynamically defined by adding to the action the following boundary term:

$$S_b = \int dt \frac{1}{2} \left(g e^{i \frac{X(t,0)}{\sqrt{2}}} + \bar{g} e^{-i \frac{\bar{X}(t,0)}{\sqrt{2}}} \right) \quad (3.27)$$

They showed that the discrete serie of states cited above saturates the dynamic of the boundary problem, since Neumann boundary states has null momentum, and the period of the boundary momenta is such that it injects momenta which are integral multiples of $\frac{1}{\sqrt{2}}$, namely the values carried by the discrete states[89].

This effects is part of a wider connection in which, once we are given a boundary conformal field theory, we can associate it different models considering fluctuations in boundary condensate, where the condensate is defined by a

boundary term added to the bulk action:

$$S = S_{bulk} + g \int dx \psi(x) \quad (3.28)$$

If the operator $\psi(x)$ is truly marginal (for what this means, see appendix C), the deformation does not take the model away from the renormalization group fixed point[78]. Thus the bulk theory remains unvaried, and the perturbation effects involve only a redefinition of the boundary conditions, thus affecting boundary states and the boundary operators.

For the sake of completeness, we have included in appendix C a comprehensive introduction to the fundamental concepts and techniques in marginal deformations of a boundary conformal field theory. These techniques will be used extensively in the following.

In the connection of the compactified boson, the presence of the enhanced affine symmetry at special values of the compactification radius coincide with the presence of new massless OPEN string states which can be used to deform the theory[90]. When $\Omega(k) = \sqrt{2}$, the closed affine algebra generators can be represented in term of the boson field via the Frenkel-Kac-Siegel construction of the affine algebra. In the closed string channel, the left moving currents are:

$$H(\zeta(p)) = \partial X(\zeta(p)), \quad E^\pm = :e^{\pm i\sqrt{2}X(\zeta(p))}: \quad (3.29)$$

and the same construction holds for the antiholomorphic sector. The $\hat{su}(2)_1$ currents modes, J_m^a and $\bar{J}_m^a$ close 2 copies of the affine algebra:

$$[J_m^a, J_n^a] = \sum_c i f_{abc} J_{m+n}^c + km \delta_{a,b} \delta_{m+n,0} \quad (3.30)$$

The vertex operators for the massless closed string states are given in term of the currents above as:

$$V_P^a(p) = J^a(\zeta(p)) \bar{\partial} \bar{X}(\bar{\zeta}(p)) e^{iPX(\zeta(p)) + \bar{X}(\bar{\zeta}(p))} \quad (3.31)$$

$$\bar{V}_P^a(p) = \bar{J}^a(\bar{\zeta}(p)) \partial X(\zeta(p)) e^{iPX(\zeta(p)) + \bar{X}(\bar{\zeta}(p))} \quad (3.32)$$

where $P = p_L + p_R$ is the total center of mass momentum of the closed string. The vertex operators for the new open string scalar states can be written in the closed string channel as:

$$\begin{aligned} S_P^a &= J^a(\zeta(p)) e^{iPX(\zeta(p))} \\ &\equiv \frac{1}{2} \left[J^a(\zeta(p)) - \Omega(p)(\bar{J}^a(\bar{\zeta}(p))) e^{iPX(\zeta(p))} \right] \Big|_{|\zeta|=1, |\zeta|=\frac{2\pi}{2\pi-\varepsilon(k)}}, \end{aligned} \quad (3.33)$$

where $\Omega(p)$ is the glueing automorphism on the boundary (see equation (2.16)).

The occurrence of extra massless open string states in equation (3.33), indicates that, at this special point in compactification moduli space, also the chiral

algebra of the boundary theory is enhanced. The associated currents $\mathbf{J}^a(\zeta(p))$ (defined as in equation (2.16)) are truly marginal operators (see appendix C) and can actually be combined to deform the original theory with a boundary action of the form:

$$S_B = \int dx(p, q) \sum_a g_a \mathbf{J}^a(\zeta(p))|_{|\zeta|=\frac{2\pi}{2\pi-\epsilon(k)}} \quad (3.34)$$

In this connection, deformations in equation (3.27) are a particular case of (3.34), in which the deformation involves only generators associated to the simple roots of $su(2)$.

Since chiral marginal deformation are truly marginal [78] the deformed model will change only for a redefinition of boundary conditions (thus boundary states and boundary operators).

In particular, in [90, 78] it had been showed that the boundary condition induced by the presence of an action boundary term like that in (3.34) is represented by a boundary state defined as [90, 78]

$$||g\rangle\rangle = g ||N(0)\rangle\rangle_{s.d.} \quad \text{with} \quad g = e^{\sum_a i g_a J_0^a} \quad (3.35)$$

where J_0^a are the horizontal $SU(2)$ algebra generators. Thus, According to formula (3.35) boundary states are actually a rotation of a “generator” boundary state, which is that associated to Neumann boundary conditions with null Wilson line. This construction cover the full moduli space of boundary states. Naively, one can expect to obtain a second branch in the moduli space of boundary states via the same construction on a “Dirchlet-like” unperturbed boundary state $||D(0)\rangle\rangle_{s.d.}$. However, Dirichlet boundary states are included in the set (3.35). They are obtained via a perturbation of the form (3.34), with the particular choice $\Gamma = e^{-i\pi J_0^1}$ [87]:

$$||D(0)\rangle\rangle_{s.d.} = e^{-i\pi J_0^1} ||N(0)\rangle\rangle_{s.d.} \quad (3.36)$$

Boundary states in eq. (3.35) satisfy the rotated gluing condition:

$$(J_m^a + \Omega \circ \gamma_{\bar{J}}(\bar{J}_{-m}^a)) ||B(g)\rangle\rangle = 0 \quad (3.37)$$

with $\gamma_{\bar{J}}(\bar{J}^a) := e^{-i \sum_b g_b J_0^b} \bar{J}^a e^{i \sum_b g_b J_0^b}$

Cardy’s boundary states (3.22) can be easily retrieved in the set (3.35). As a matter of fact, when the unperturbed boundary state is a Neumann one with the Wilson line parametrized by $\tilde{t}_+$, the general gluing condition

$$E^\pm(\zeta) = e^{\pm i\sqrt{2}\tilde{t}_+} \bar{E}^\pm(\bar{\zeta}) \quad (3.38)$$

is invariant under the shift $\tilde{t}_+ \rightarrow \tilde{t}_+ + \pi\sqrt{2}$. The marginal perturbation implementing this shift, $\Gamma = \exp(i\frac{\pi}{\sqrt{2}}J_0^3)$ acts non trivially on a Neumann boundary state $||N(\tilde{t}_+)\rangle\rangle$, producing exactly a switch of the sign in front of the Ishibashi

state built on the $j = \frac{1}{2}$ $\hat{su}(2)_1$ module. Thus we can conclude that, in the description of eq. (3.35), Cardy's boundary corresponds to $\|N(0)\rangle\rangle_{s.d.}$ and $\|N(\pi/\sqrt{2})\rangle\rangle_{s.d.}$:

$$\|0\rangle\rangle = 2^{\frac{1}{4}} \left(|0\rangle\rangle + |\frac{1}{2}\rangle\rangle \right) \equiv \|N(0)\rangle\rangle_{s.d.} \quad (3.39a)$$

$$\|\frac{1}{2}\rangle\rangle = 2^{\frac{1}{4}} \left(|0\rangle\rangle - |\frac{1}{2}\rangle\rangle \right) \equiv \|N(\pi/\sqrt{2})\rangle\rangle_{s.d.} \quad (3.39b)$$

This result holds for any gluing condition we decide to fix on the boundary of the cylinder. The independent gluing conditions are then parametrized by $SO(3)$, because the elements which yield trivial gluing automorphism are those associated to the center of $SU(2)$. Cardy's boundary states are then those associated to the central elements of $SU(2)$.

Authors of [91] introduced an alternative description in which they showed that the boundary states generated by the perturbation (3.34) can be directly parametrized by means of the associated $SU(2)$ elements g and described via the following formula:

$$\|g\rangle\rangle = \sum_{j \in \mathcal{Y}} \sum_{\substack{-j \leq m \leq j \\ -j \leq n \leq j}} D_{m, -n}^j(g) |j; m, n\rangle\rangle \quad (3.40)$$

where $|j; m, n\rangle\rangle$ is the Ishibashi state associated to the discrete $h = j^2$ module of the $su(2)$ algebra and $D_{m, -n}^j(g)$ is the matrix elements of the j -representation of $g \in SU(2)$, parametrized as $g = \begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix}$. In this description, the generating boundary state $\|N(0)\rangle\rangle_{s.d.}$ is associated to the $SU(2)$ identity, $g = \mathbb{I}$ $N(0) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$.

We will not use this construction to define BIO, however we report it because it allows to write the annulus amplitude with different boundary conditions associate respectively to $S_{\varepsilon(k)}^{(-)}$ and $S_{\varepsilon(k)}^{(+)}$ in a quite simple way. Let us consider the transition amplitude between two boundary states $|\alpha\rangle$ and $|\beta\rangle$ which can be defined via the action of $g \in SU(2)$ on $|\alpha\rangle$, $|\beta\rangle = g|\alpha\rangle$. From the above definitions of boundary states, it can be shown that $A_{\alpha, g\alpha}$ depends only on the conjugacy classes of g (for a detailed demonstration see [78], or [91], section 4). Therefore, we can choose to deform the boundary state with an element in a given torus of $SU(2)$: $t = h^{-1}gh = \begin{pmatrix} e^{4\pi i \lambda} & 0 \\ 0 & e^{-4\pi i \lambda} \end{pmatrix}$ following the detailed analysis in [91], we can finally write the amplitude as

$$\mathcal{A}(p) = \frac{1}{\sqrt{2}} \sum_{j \in \frac{1}{2}\mathbb{Z}_+} \cos(8\pi j \lambda) \vartheta_{2\tilde{q}}(\tilde{q}) \quad (3.41)$$

where $\tilde{q} = e^{-\frac{2\pi i}{\tau}}$

A boundary perturbation will affect the boundary operators spectrum too. Generically, when the perturbing field is truly marginal, the study of the deformation of a correlator containing both bulk and boundary fields allows to

define the image $\tilde{\psi}_j$ of a boundary field ψ_j under a rotation generated by the perturbing field ψ as [78] (see appendix C):

$$\tilde{\psi}_j = \left[e^{\frac{1}{2}g\psi} \psi_j \right] (u) := \sum_{n=0}^{\infty} \frac{g^n}{2^n n!} \oint_{C_1} \frac{dx_1}{2\pi} \cdots \oint_{C_n} \frac{dx_n}{2\pi} \psi_j(u) \psi(x_n) \cdots \psi(x_1) \quad (3.42)$$

where C_i are small circles surrounding the insertion points of the operators $\psi(x_i)$ on the boundary.

As they stands, representations (3.35) and (3.40) of boundary conditions do not allow to successfully explain how the transition between pairwise adjacent boundary conditions take place. Thus, we have introduced a new representation for the infinite set of boundary conditions which can be applied to the compactified boson at the self-dual radius. This representation merges the infinite choice of boundary conditions with the the necessity to have a BIO acting “*a lá Cardy*”, ie mediating between the diffent boundary conditions exploiting the fusion rules of the associated chiral algebra.

We can exploit construction (3.35) to parametrize the generic boundary condition defined over the (inner or outer) boundary of the k -th cylindrical end, represented by the boundary state $\|g(k)\rangle\rangle$, with a couple of elements:

$$(\|J(k)\rangle\rangle, \Gamma(k)) \quad \text{with} \quad \begin{cases} J(k) & \in \mathcal{A} \\ \Gamma(k) & \in \frac{SU(2)}{\mathbb{Z}_2} \end{cases} \quad (3.43)$$

being $\|J(k)\rangle\rangle$ a Cardy’s boundary state (thus corresponding to an element in the center of $SU(2)$) and Γ an $\frac{SU(2)}{\mathbb{Z}_2}$ group element, such that:

$$\|g(k)\rangle\rangle = \Gamma \|J(k)\rangle\rangle. \quad (3.44)$$

In this connection, the model can be defined not as a truly marginal deformation of a open string theory by means of the action of elements of the affine chiral $SU(2)$, but as a truly marginal deformation of the $\hat{\mathfrak{su}}(2)_{k=1}$ by means of $SO(3)$ elements.

Thus, BIO for the compactified boson with $\Omega(p) = \sqrt{2}$ are actually given by the consequent deformation induced by the two adjacent boundary conditions on $\hat{\mathfrak{su}}(2)_{k=1}$ WZW model boundary insertion operators which we introduced in formula (3.23)

To understand how this deformation affects and define boundary insertion operators, let us consider the (p, q) -edge of the ribbon graph. Let the two adjacent boundary conditions be defined as:

$$\|g_1(p)\rangle\rangle = \Gamma_1(p) \|J_1(p)\rangle\rangle \quad (3.45a)$$

$$\|g_2(q)\rangle\rangle = \Gamma_2(q) \|J_2(q)\rangle\rangle \quad (3.45b)$$

According to the parametrization of boundary condition introduced above, BIO must mediate both between Cardy’s boundary states and between the

$\frac{SU(2)}{\mathbb{Z}_2}$ elements $\Gamma_1(p)$ and $\Gamma_2(q)$. While the former action is achieved through the fusion prefactor $\mathcal{N}_{J(p)j(p,q)}^{J(q)}$, the latter can be understood deforming BIO with the action of both the boundary potentials which we are adding on the (p, q) -edge. As a matter of fact, according to equation (3.45), the theory on the (p) -th polytope is deformed by the action of the boundary term $S_{B(p)} = \int dx(p, q) \mathbf{J}_1(\zeta(p))|_{|\zeta(p)|=\frac{2\pi}{2\pi-\varepsilon(p)}}$, while the theory on the (q) -th polytope is deformed by the boundary term $S_{B(q)} = \int dx(q, p) \mathbf{J}_2(\zeta(q))|_{|\zeta(q)|=\frac{2\pi}{2\pi-\varepsilon(q)}}$. Recalling that $x(q, p) = -x(p, q)$, (the functional part of) boundary insertion operators, we propose $\psi_{j(p,q)}$ to be deformed by a suitable combination of the $SO(3)$ operators which are associated to the above boundary terms. We ask this combination to cancel, on the boundary, the global effect of the boundary deformation, to let the two ends glue dynamically in such a way that this dynamic is actually governed by the fusion rules of the WZW model. This corresponds to a perform over $\psi_{j(p,q)}$ a rotation induced by the $\frac{SU(2)}{\mathbb{Z}_2}$ element $\bar{\Gamma} = \Gamma_2 \Gamma_1^{-1}$, with $\Gamma_i = e^{i\mathbf{J}_i}$. In the same way, $\psi_{j(q,p)}$ will be deformed by the action of by the action of $\bar{\Gamma}^{-1} = \Gamma_1 \Gamma_2^{-1}$.

To show how the above rotation alter the functional part of boundary insertion operators, let us consider the explicit expression of components of $\hat{\mathfrak{su}}(2)_1$ BIO which we introduced in (3.23). Let us drop for a while the dependence from the fusion rule factor $\mathcal{N}_{J_1(p)j(p,q)}^{J_2(q)}$. Such components are labelled by two (semi-)integers $j = 0, \frac{1}{2}$ and $-j < m < j$. For $j = 0$ the unique component is the identity operator $\psi_{0,0}x(p, q) = \mathbb{I}$, while for $j = \frac{1}{2}$ the two components are $\psi_{\frac{1}{2} \pm \frac{1}{2}} = e^{\pm \frac{i}{\sqrt{2}} X[\zeta(p)(z(p,q))]} \Big|_{y(p,q)=0}$.

Let us consider the action of the deformation on $\psi_{j(p,q)}$ generated by $\bar{\Gamma} = e^{i\bar{\mathcal{J}}}$ (since we are not moving to a definite representation, we can omit the quantum number m). According to (3.42), the rotated boundary operator will be:

$$\tilde{\psi}_{j(p,q)}(u(p, q)) = \left[e^{\frac{1}{2}\bar{\mathcal{J}}} \psi_{j(p,q)} \right] (u(p, q)) \quad (3.46)$$

We can compute explicitly the expression of $\tilde{\psi}_{j(p,q)}$ thanks to the self-locality of the boundary operators, and to the OPE between the truly marginal fields in the chiral algebra and a boundary operator:

$$\mathbf{J}(x)\psi_j(u) \sim \frac{\mathbf{X}_J^j}{x-u}\psi_j(u) \quad (3.47)$$

where $\mathbf{X}_J^j$ is the natural action of the chiral algebra on a state of the $h = j^2$ $\hat{\mathfrak{su}}(2)_1$ module (see formula (2.13)).

An order by order computation in (3.46) gives:

$$\tilde{\psi}_{j(p,q)}(u(p, q)) = e^{\frac{i}{2}\mathbf{X}_J^j}\psi_{j(p,q)} \quad (3.48)$$

i.e. the natural action of the chiral algebra on the vertex algebra fields translates into the natural action of an element of $SU(2)/\mathbb{Z}_2$ on the components of

the primary field associated to a given $\hat{su}(2)_1$'s module. Moving to a specific representation, equation (3.48) becomes:

$$\tilde{\psi}_{(j,m)(p,q)}(u(p,q)) = D_{mn}^j(\Gamma) \psi_{(j,n)(p,q)} \quad (3.49)$$

where D_{mn}^j are the Wigner functions associated to $\Gamma = \begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix}$:

$$D_{m,n}^j(\Gamma) = \sum_{l=\max(0,n-m)}^{\min(j-m,j-n)} \frac{[(j+m)!(j-m)!(j+n)!(j-n)!]^{\frac{1}{2}}}{(j-m-l)!(j+n-l)!l!(m-n+l)!} \times a^{j+n-l} (a^*)^{j-m-l} b^l (-b^*)^{m-n+l}. \quad (3.50)$$

Obviously, the spectrum of $\hat{su}(2)_1$ boundary primary fields must be invariant under the action of $\mathbb{Z}_2$. A first intuition about this comes once we remember the gluing condition (3.38) being invariant under the shift generating the $\|1/2\rangle$ Cardy's state: the spectrum of boundary operators is generated the action of a copy of the chiral algebra:

$$\mathbb{W}(\zeta) = \begin{cases} W(\zeta) & \Im \zeta \geq 0 \\ \Omega \circ \gamma_{\bar{\Gamma}}(\bar{W}(\bar{\zeta})) & \Im \zeta < 0 \end{cases}. \quad (3.51)$$

The gluing automorphism itself generates the boundary operators' spectrum, thus a boundary condensate which leaves invariant the gluing automorphism automatically leaves invariant the boundary operators' spectrum too. An explicit computation via formula (3.49) of Cardy's boundary operators rotated by the action of the boundary condensate correspondent to the $\|N(\pi/\sqrt{2})\rangle_{s.d.}$ boundary state confirms this statement: rotated boundary operators are obtained by multiplication by an inessential phase factor.

Finally, restoring the fusion multiplicative coefficient, we have the following expression for boundary insertion operators for the compactified boson at the self dual radius:

$$\psi_{[j,m](p,q)}^{[J_2, \Gamma_2](q) [J_1, \Gamma_1](p)} = \sum_{n=-j}^j D_{m,n}^{j(p,q)}(\Gamma_2 \Gamma_1^{-1}) \psi_{[j,n](p,q)}^{J_2(q) J_1(p)}. \quad (3.52)$$

where $\psi_{j(p,q)}^{J(p) J(q)}(x(p,q)) = \mathcal{N}_{J(p) j(p,q)}^{J(q)} \psi_{j(p,q)}(x(p,q))$

3.3 The algebra of rotated Boundary Insertion Operators

According to last section remarks, boundary insertion operators for the compactified boson at enhanced symmetry values of the self-dual radius have the following expression:

$$\psi_{[j,m](p,q)}^{[J_2, \Gamma_2](q) [J_1, \Gamma_1](p)} = \sum_{n=-j}^j D_{m,n}^{j(p,q)}(\Gamma_2 \Gamma_1^{-1}) \psi_{[j,n](p,q)}^{J_2(q) J_1(p)}. \quad (3.53)$$

The aim of this, quite technical, section, is to show that with this choice, effects of boundary perturbations do not affect the algebra of boundary operators, thus they do not change the dynamic of the model. This is a check of consistency of the above choice for boundary insertion operators, since actually all deformations we have considered are induced by truly marginal operators, thus they must not break the $su(2)$ chiral algebra.

Let us deal with the simplest case: the amplitude computed on the open surface associated with the sphere with three punctures. The amplitude over each cylindrical end will involve a sum over characters twisted by the action of conjugacy classes of $\frac{SU(2)}{\mathbb{Z}_2}$. The full amplitude will involve a sum over the intermediate channels associated with the three edges of the ribbon graph, each being associated to an automorphism by the operator $U = e^{i[g_1(q)-g_2(p)]}$, $\forall (p, q) = (1, 2), (1, 3), (2, 3)$. Such an automorphism acts at BIOs' level: the BIO associated to the (p, q) -th edge will be the Cardy's one rotated by the action of the $\frac{SU(2)}{\mathbb{Z}_2}$ element $\Gamma_2 \Gamma_1^{-1}$.

The algebra of rotated BIOs follows from their definition. Let us notice that rotated BIOs are just a superposition of the different components of Cardy's $\hat{su}(2)_1$ primary operators (with respect to the affine chiral algebra).

Let us focus our attention on the two points function. When we consider two BIOs both mediating a boundary condition changing in the p -to- q direction, we deal with the following expression:

$$\begin{aligned} & \left\langle \psi_{[j, m](p, q)}^{[J_2, \Gamma_2](q) [J_1, \Gamma_1](p)}(x_1(p, q)) \psi_{[j', m'](p, q)}^{[J_4, \Gamma_4](q) [J_3, \Gamma_3](p)}(x_2(p, q)) \right\rangle = \\ & \sum_{n, n'} D_{m, n(p, q)}^{j(p, q)} (\Gamma_2 \Gamma_1^{-1}) D_{m', n'(p, q)}^{j'(p, q)} (\Gamma_4 \Gamma_3^{-1}) \left\langle \psi_{[j, n](p, q)}^{J_2(q) J_1(p)}(x_1(p, q)) \psi_{[j', n'](p, q)}^{J_4(q) J_3(p)}(x_2(p, q)) \right\rangle \end{aligned} \quad (3.54)$$

First of all, we must notice that a coherent gluing impose the two operators to mediate between the same boundary conditions (see eq. (B.3)), thus the above expression reduces to:

$$\begin{aligned} & \left\langle \psi_{[j, m](p, q)}^{[J_2, \Gamma_2](q) [J_1, \Gamma_1](p)}(x_1(p, q)) \psi_{[j', m'](p, q)}^{[J_2, \Gamma_2](q) [J_1, \Gamma_1](p)}(x_2(p, q)) \right\rangle = \\ & \sum_{n, n'} D_{m, n(p, q)}^{j(p, q)} (\Gamma_2 \Gamma_1^{-1}) D_{m', n'(p, q)}^{j'(p, q)} (\Gamma_2 \Gamma_1^{-1}) \left\langle \psi_{[j, n](p, q)}^{J_2(q) J_1(p)} \psi_{[j', n'](p, q)}^{J_2(q) J_1(p)} \right\rangle \end{aligned} \quad (3.55)$$

Equation (3.55) shows that the net effect of the rotation on the two points function vanish, because we are actually implementing the same $SU(2)$ rotation on all boundary fields entering in the unperturbed correlator. Thus the local $SU(2)$ invariance ensures:

$$\begin{aligned} & \left\langle \psi_{[j, m](p, q)}^{[J_2, \Gamma_2](q) [J_1, \Gamma_1](p)}(x_1(p, q)) \psi_{[j', m'](p, q)}^{[J_2, \Gamma_2](q) [J_1, \Gamma_1](p)}(x_2(p, q)) \right\rangle = \\ & \left\langle \psi_{[j, m](p, q)}^{J_2(q) J_1(p)}(x_1(p, q)) \psi_{[j', m'](p, q)}^{J_2(q) J_1(p)}(x_2(p, q)) \right\rangle \end{aligned} \quad (3.56)$$




Figure 3.3: bla bla bla

Dealing with the two points function between a p -to- q and q -to- p mediating operators, the situation is slightly different. We have to compute:

$$\begin{aligned} \left\langle \psi_{[j, m](p, q)}^{[J_2, \Gamma_2](q) [J_1, \Gamma_1](p)}(x_1(p, q)) \psi_{[j', m'](q, p)}^{[J_1, \Gamma_1](p) [J_2, \Gamma_2](q)}(x_2(q, p)) \right\rangle = \\ \sum_{n, n'} D_{m, n(p, q)}^{j(p, q)} (\Gamma_2 \Gamma_1^{-1}) D_{m', n'(q, p)}^{j'(q, p)} (\Gamma_1 \Gamma_2^{-1}) \times \\ \left\langle \psi_{[j, n](p, q)}^{J_2(q) J_1(p)}(x_1(p, q)) \psi_{[j', n'](q, p)}^{J_1(p) J_2(q)}(x_2(q, p)) \right\rangle \quad (3.57) \end{aligned}$$

As a matter of fact, in the previous expression we are dealing with a representation of diagonal subgroup of the direct product $\frac{SU(2)}{\mathbb{Z}_2}(p, q) \times \frac{SU(2)}{\mathbb{Z}_2}(q, p)$, thus it holds (see eq. (D.1)):

$$D_{m, n(p, q)}^{j(p, q)} (\Gamma_2 \Gamma_1^{-1}) D_{m', n'(q, p)}^{j'(q, p)} (\Gamma_1 \Gamma_2^{-1}) = D_{m, n; m', n'}^{j \times j'}(\mathbb{I}) \quad (3.58)$$

The trivial Clebsh-Gordan expansion (eq. (D.2)) gives (we omit the polytope indices writing the Clebsh-Gordan coefficients):

$$\begin{aligned} \left\langle \psi_{[j, m](p, q)}^{[J_2, \Gamma_2](q) [J_1, \Gamma_1](p)}(x_1(p, q)) \psi_{[j', m'](q, p)}^{[J_1, \Gamma_1](p) [J_2, \Gamma_2](q)}(x_2(q, p)) \right\rangle = \\ \sum_{n, n'} \sum_{J, N} C_{j_1 m_1 j_2 m_2}^{J, N} C_{j_1' n_1' j_2' n_2'}^{J, N} \left\langle \psi_{[j, n](p, q)}^{J_2(q) J_1(p)}(x_1(p, q)) \psi_{[j', n'](q, p)}^{J_1(p) J_2(q)}(x_2(q, p)) \right\rangle = \\ \left\langle \psi_{[j, m](p, q)}^{J_2(q) J_1(p)}(x_1(p, q)) \psi_{[j', m'](q, p)}^{J_1(p) J_2(q)}(x_2(q, p)) \right\rangle. \quad (3.59) \end{aligned}$$

In the last equation we have used the unitarity of Clebsh-Gordan coefficients (see equation (D.3b)).

To calculate the OPE of rotated BIOs, let us notice that the rotation generated by the boundary condensate does not change them coordinate dependence. Let us consider the situation depicted in figure 3.3.

OPE between $\psi_{[j_1, m_1](r, p)}^{[J_1, \Gamma_1](p) [J_3, \Gamma_3](r)}$ and $\psi_{[j', m'](q, r)}^{[J_3, \Gamma_3](r) [J_2, \Gamma_2](q)}$ will mediate a change

in boundary conditions from $[J_2, \Gamma_2](q)$ to $[J_1, \Gamma_1](p)$. In particular,

$$\begin{aligned} & \psi_{[j_1, m_1](r, p)}^{[J_1, \Gamma_1](p) [J_3, \Gamma_3](r)}(\omega_r) \psi_{[j', m'](q, r)}^{[J_3, \Gamma_3](r) [J_2, \Gamma_2](q)}(\omega_q) = \\ & \sum_{n_1(r, p) n_2(q, r)} D_{m_1 n_1(p, q)}^{j_1(r, p)}(\Gamma_1 \Gamma_3^{-1}) D_{m_2 n_2(q, r)}^{j_2(q, r)}(\Gamma_3 \Gamma_2^{-1}) \psi_{[j_1, n_1](r, p)}^{J_1(p) J_3(r)}(\omega_r) \psi_{[j_2, m_2](q, r)}^{J_3(r) J_2(q)}(\omega_q) \end{aligned}$$

We are dealing again with a representation of the diagonal subgroup of the direct product $\frac{SU(2)}{\mathbb{Z}_2}(r, p) \times \frac{SU(2)}{\mathbb{Z}_2}(q, r)$, thus applying (D.1) and the Clebsh-Gordan series expansion (D.2) we are left with:

$$\begin{aligned} & \sum_{\substack{n_1(r, p) \\ n_2(q, r)}} \sum_{\substack{j=|j_1-j_2| \\ |m| \leq j \\ |n| \leq j}}^{j_1+j_2} C_{j_1(r, p) m_1(r, p) j_2(q, r) m_2(q, r)}^{j m} D_{m n}^j(\Gamma_1 \Gamma_2^{-1}) \\ & \times C_{j_1(r, p) n_1(r, p) j_2(q, r) n_2(q, r)}^{j n} \psi_{[j_1, n_1](r, p)}^{J_1(p) J_3(r)}(\omega_r) \psi_{[j_2, n_2](q, r)}^{J_3(r) J_2(q)}(\omega_q) \quad (3.60) \end{aligned}$$

The OPE between Cardy's boundary operators reads:

$$\begin{aligned} & \psi_{[j_1, n_1](r, p)}^{J_1(p) J_3(r)}(\omega_r) \psi_{[j_2, n_2](q, r)}^{J_3(r) J_2(q)}(\omega_q) = \sum_{j_3 n_3} |\omega_r - \omega_q|^{H(q, p) - H(r, p) - H(q, r)} \\ & C_{j_1 n_1 j_2 n_2}^{j_3 n_3} C_{j_1 j_2 j_3}^{J_1(p) J_3(r) J_2(q)} \psi_{[j_3, n_3](q, p)}^{J_1(p) J_2(q)}(\omega_q). \quad (3.61) \end{aligned}$$

The Clebsh-Gordan coefficients $C_{j_1 n_1 j_2 n_2}^{j_3 n_3}$ compensate the fact that the LHS and RHS terms have different transformation behavior under the action of the horizontal $su(2)$ algebra, while the coefficients $C_{j_1 j_2 j_3}^{J_1(p) J_3(r) J_2(q)}$ reflect the non trivial dynamic on each trivalent vertex of the ribbon graph.

The inclusion of this last OPE into (3.60) and the Clebsh-Gordan coefficients' unitarity (equation (D.3b)) leave us with:

$$\begin{aligned} & \psi_{[j_1, m_1](r, p)}^{[J_1, \Gamma_1](p) [J_3, \Gamma_3](r)}(\omega_r) \psi_{[j', m'](q, r)}^{[J_3, \Gamma_3](r) [J_2, \Gamma_2](q)}(\omega_q) = \\ & \sum_{j_3 m} C_{j_1 m_1 j_2 m_2}^{j_3 m} C_{j_1 j_2 j_3}^{J_1(p) J_3(r) J_2(q)} \psi_{[j_1, m_1](q, p)}^{[J_1, \Gamma_1](p) [J_2, \Gamma_2](q)}(\omega_p) \quad (3.62) \end{aligned}$$

We demonstrate that OPE between rotated BIOs is formally equal to OPE between unrotated BIOs. Thus, on the ribbon graph the non trivial dynamic is given by the fusion among the three representations entering in each trivalent vertex.

3.4 The action of BIOs at the self-dual radius

With the above remarks, we can investigate the properties of the four points functions of BIOs exploiting their crossing properties.

First of all let us consider the natural picture in which the computation of a four points function arises. Let us consider two near trivalent vertexes. Due




Figure 3.4: Four points function crossing symmetry

to the variable connectivity of the triangulation, the two configuration shown in figure 3.4 are both admissible. The transition from the situation depicted in the lhs and the one in the rhs of the pictorial identity of figure 3.4, corresponds exactly to the transition between the s -channel and the t -channel of the four point blocks of a single copy of the bulk theory, thus the two factorization of the four points function $\langle \psi_{j_1(s,p)}^{J_1(p) J_4(s)} \psi_{j_2(r,s)}^{J_4(s) J_3(r)} \psi_{j_3(q,r)}^{J_3(r) J_2(q)} \psi_{j_4(p,q)}^{J_2(q) J_1(p)} \rangle$, pictorially represented in 3.4, are related by the bulk crossing matrices:

$$F_{j_6(s,q) j_5(r,p)} \begin{bmatrix} j_4(p, s) & j_1(q, p) \\ j_3(s, r) & j_2(r, q) \end{bmatrix} \quad (3.63)$$

The explicit computation of the two factorization leads to the relation:

$$\begin{aligned} & \mathcal{C}_{j_2(r,s) j_3(q,r) j_5(q,s)}^{J_4(s) J_3(r) J_2(q)} \mathcal{C}_{j_1(s,p) j_5(q,s) j_1(s,p)}^{J_1(p) J_4(s) J_2(q)} \mathcal{C}_{j_1(s,p) j_1(s,p) 0}^{J_1(p) J_2(q) J_1(p)} = \\ & \sum_{j_5(r,p)} F_{j_6(s,q) j_5(r,p)} \begin{bmatrix} j_4(p, s) & j_1(q, p) \\ j_3(s, r) & j_2(r, q) \end{bmatrix} \times \\ & \mathcal{C}_{j_1(s,p) j_2(r,s) j_6(r,p)}^{J_1(p) J_4(s) J_3(r)} \mathcal{C}_{j_3(q,r) j_4(p,q) j_6(p,r)}^{J_3(r) J_2(q) J_1(p)} \mathcal{C}_{j_6(r,p) j_6(p,r) 0}^{J_1(p) J_3(r) J_1(p)}, \quad (3.64) \end{aligned}$$

i.e. the usual BCFT sewing relation among boundary operators' OPEs.

This complete our analysis of the conformal properties of the full theory arising by glueing together the BCFTs defined over each cylindrical ends: with the above construction, BIOs play exactly the role usual boundary operators play in BCFT.

This analogy allows us to apply to BIOs all boundary operators properties. In particular, we can identify their OPE coefficients with the fusion matrices (3.63) with the following entries assignation:

$$\mathcal{C}_{j_1(s,p) j_2(r,s) j_3(q,r)}^{J_1(p) J_2(q) J_3(s)} = F_{J_2(q) j_3(q,r)} \begin{bmatrix} J_1(p) & J_3(s) \\ j_1(s,p) & j_2(r,s) \end{bmatrix} \quad (3.65)$$

Relation (3.65), obtained first in [92] for the A -series minimal models exploiting the fact that both the primary and boundary conditions labels fall in

the same set, has been extended to all minimal models and extended rational conformal field theories in [83] and [93] noticing the full analogy between the equation (3.64) and the pentagon identity for the fusing matrices.

According to [94], WZW-models fusion matrices coincide with the $6j$ -symbols of the corresponding quantum group with deformation parameter given by the $(k + h^\vee)$ -th root of the identity, where k and $h^\vee$ are respectively the level and the dual Coxeter number of the extended algebra. Thus, with $k = 1$ and $h^\vee = 2$, the OPEs coefficients are the $SU(2)_{Q=e^{\frac{2}{3}\pi i}}$ $6j$ -symbols:

$$C_{j_1(s,p) j_2(r,s) j_3(q,r)}^{J_1(p) J_2(q) J_3(s)} = \left\{ \begin{matrix} j_1(s,p) & J_1(p) & J_2(q) \\ J_3(s) & j_2(r,s) & j_3(q,r) \end{matrix} \right\}_{Q=e^{\frac{2}{3}\pi i}} \quad (3.66)$$

3.5 Open string amplitude on a RRT

With the computation of OPE's coefficients we have all the building blocks to construct an open string amplitude on the domain defined by the open Riemann surface M_∂ .

As first step, let us extend results of the previous section to higher dimensional target spaces. To this end, let us consider D scalar fields $X^\alpha, \alpha = 1, \dots, D$ which, as stated in (2.1), wind ν^α times around the homology cycles of the compact target space manifold. Let us indicate with $E_{\alpha\beta} = G_{\alpha\beta} + B_{\alpha\beta}$ the background matrix on the compact target space manifold, where G is the metric and B the Kalb-Ramond field. The central charge of the model is $c = D$.

Let us find out which is the moduli space of inequivalent compactifications. This can be achieved by considering the torus partition function of the model: it factorizes into the product of contributions of each compact direction:

$$Z_{T_d} = \frac{1}{|\eta(\tau)|^2} \sum_{\Gamma_{D,D}} q^{\frac{p_L^2}{4}} \bar{q}^{\frac{p_R^2}{4}} \quad (3.67)$$

Asking for modular invariance let the total momentum $\hat{p} = \begin{pmatrix} p_L \\ p_R \end{pmatrix}$ take values into a self-dual, even-integer, Lorentzian lattice $\Gamma_{D,D}$ [95]. The space of such inequivalent lattices is locally isomorphic to:

$$\mathcal{M} = O(d, d, \mathbb{Z}) \backslash O(d, d) / [O(d) \times O(d)], \quad (3.68)$$

which thus is the moduli space of inequivalent toroidal compactifications in a D dimensional target space[96, 95]. The different orbits in this moduli space give rise to different theories in which the fundamental $U(1)_L \times U(1)_R$ current symmetry can be enhanced to different symmetry groups of rank at least D . This group plays the role of gauge group in the target space. Our choice is to compactify each direction at the self dual radius, because this allow to exploit the previous construction and define a coherent gluing of the conformal theory

along the ribbon graph. This means to choose a specific orbit into (3.68), *i.e.* to fix definitively the target space modular structure. Moreover, the background gauge group coming from the closed string sector defined on the cylindrical ends is $[SU(2)_L \times SU(2)_R]^D$. On the contrary, we could choose not to fix the compactification radius as the self-dual one. This would give us more freedom in choosing the target space modular and metrical structure, and consequently the enhanced symmetry group. However, in doing so we can loose information about the dynamic of the theory on the ribbon graph, because, with the exception of some particular cases which we will introduce in the following chapter, combinatorial factor of BIOs would be defined only as a formal map.

Thus, let us consider each direction compactified at the self dual radius. The amplitude on each $\Delta_{\varepsilon(p)}^*$ will receive a contribution from every direction:

$$\mathcal{A}_{\Delta_{\varepsilon(p)}^*} = \frac{1}{2^{\frac{D}{2}} \left[\eta \left(e^{-5\frac{4\pi}{\theta(p)}} \right) \right]^D} \prod_{\alpha=1}^D \sum_{j(p)=0, \frac{1}{2}} \cos(8\pi j(p) \lambda^\alpha(p)) e^{-\frac{4\pi}{\theta(p)} j(p)^2} \quad (3.69)$$

Moreover, Boundary Insertion Operators are primaries of the conformal theory, thus they also factorize into the contribution of each direction. This means that the full boundary theory factorizes into the contribution of each compact direction. In this connection, we can exploit a construction introduced in [67], which, exploiting a edge vertex factorization of the most general correlator we can write on the ribbon graph, allows to write the contribution to the amplitude from each compact direction as:

$$\begin{aligned} Z(|P_{T_l}|) = & \left(\frac{1}{\sqrt{2}} \right)^{N_0(T)} \sum_{\{j_p \in \frac{1}{2}\mathbb{Z}_+\}} \sum_{\{j_{(r,p)}\}} \prod_{\{\rho^0(p,q,r)\}}^{N_2(T)} \left\{ \begin{array}{ccc} j_{(r,p)} & j_p & j_r \\ j_q & j_{(q,r)} & j_{(p,q)} \end{array} \right\}_{Q=e^{\frac{\pi}{3}i}} \times \\ & \times \prod_{\{\rho^1(p,r)\}}^{N_1(T)} \left(b_{j_{(r,p)}}^{j_p j_r} \right)^2 L(p,r)^{-2H_{j_{(r,p)}}} \quad (3.70) \end{aligned}$$

Collecting the contribution of each direction and applying this results on the $N(0)$ channels defined by (3.69) we finally have:

$$\begin{aligned} Z(|P_{T_l}|, D) = & \frac{1}{2^{\frac{DN_0(T)}{2}}} \prod_{\alpha=1}^D \left[\sum_{\{j_p \in \frac{1}{2}\mathbb{Z}_+\}} \sum_{\{j_{(r,p)}\}} \prod_{\{\rho^0(p,q,r)\}}^{N_2(T)} \left\{ \begin{array}{ccc} j_{(r,p)} & j_p & j_r \\ j_q & j_{(q,r)} & j_{(p,q)} \end{array} \right\}_{Q=e^{\frac{\pi}{3}i}} \times \right. \\ & \times \left. \prod_{\{\rho^1(p,r)\}}^{N_1(T)} \left(b_{j_{(r,p)}}^{j_p j_r} \right)^2 L(p,r)^{-2H_{j_{(r,p)}}} \cos(8\pi j_p \lambda(i)) \frac{e^{-\frac{4\pi}{\theta(i)} j_p^2}}{\eta(e^{-5\frac{4\pi}{\theta(i)}})} \right]_{(\alpha)} \quad (3.71) \end{aligned}$$

where the subscript (α) indicates the contribution of the α -th direction.

Chapter 4

Inclusion of Open String gauge degrees of freedom: a proposal

A necessary step to rephrase our model in a gauge/gravity correspondence connection would be to include open string gauge degrees of freedom (propagating) along the boundaries of M_∂ . As a matter of fact the model we built in previous chapters presents an $SU(2)$ gauge symmetry at spacetime level. However, this symmetry is due to the particular geometry of the system, and it does not allow to define an appropriate gauge coloring neither of the ribbon graph underlying the triangulation, nor of the boundary components. Thus, it seems more appropriate to follow usual techniques in open string theory, where a non Abelian gauge theory can be naturally included into an open string model by a suitable assignation of non-Abelian Chan Paton factors at the open string endpoints [97]. This will define some modifications in our model. First of all, vertex algebra operators (and consequently BIOs) will be valued in the associated Lie algebra, thus ribbon graph amplitudes will be weighted by an appropriate group factor. The natural consequence will be the coupling of the conformal field theory on each cylindrical end with a background gauge field. In this connection, we will show that we can rephrase the arising of a target space non-Abelian gauge symmetry which arise with this process with a change in the background matrix, *i.e.* the process is equivalent to move to another point of moduli space of toroidal compactifications given in (3.68). According to [96, 90], when the chosen background is a fixed point under the extended D -dimensional T-duality group, the annulus BCFT is equivalent to a $\mathfrak{g}_{k=1}$ WZW model, where $\mathfrak{g}_{k=1}$ is a level-one untwisted affine Lie algebra associated to a semisimple product of simply laced Lie algebras of total rank D .

In the last section, we will try to understand how to extend the gluing technique described in the previous chapters to the current situation.

Conventions about $U(N)$ algebra

Let us represent $U(N) \sim SU(N) \times U(1)$. Thus we can choose a base of $U(N)$ generators as:

- The collection of $SU(N)$ generators:

$$T^a \quad a = 1, \dots, N^2 - 1 \quad (4.1)$$

satisfying the traceless condition $\text{Tr}(T^a) = 0$.

- $T^0 = C\mathbb{I}$, where C is arbitrary constant.

Let us impose the following normalization condition:

$$\text{Tr}(T^a T^b) = \frac{1}{2} \delta^{ab}. \quad (4.2)$$

Equation (4.2) fixes $C = \frac{1}{\sqrt{2N}}$.

The completeness relation for generators reads:

$$(T^a)_j^i (T^b)_l^k = \frac{1}{2} \delta_l^i \delta_j^k \quad \begin{array}{l} a = 1, \dots, N^2 - 1 \\ i, j, k, l = 1, \dots, N \end{array} \quad (4.3)$$

The $U(N)$ algebra is defined by:

$$[T^a, T^b] = i f^{abc} T^c \implies f^{abc} = \frac{2}{i} \text{Tr}([T^a, T^b] T^c) \quad (4.4a)$$

$$\{T^a, T^b\} = d^{abc} T^c \implies d^{abc} = 2 \text{Tr}(\{T^a, T^b\} T^c) \quad (4.4b)$$

where f^{abc} (resp. d^{abc}) are the antisymmetric (resp. symmetric) structure constants.

With the above conventions, a rapid computation shows:

$$\sum_{b,c} f^{abc} f^{dbc} = N (\delta^{ad} - \delta^{a0} \delta^{d0}) \quad (4.5a)$$

$$\sum_{b,c} d^{abc} d^{dbc} = N (\delta^{ad} + \delta^{a0} \delta^{d0}) \quad (4.5b)$$

4.1 Chan-Paton factors on a RRT

Chan-Paton factors are non-dynamical degrees of freedom which can be added to open string endpoints. To show in a few words how it works, let us consider an open string worldsheet (let us remember that the cylindrical ends $\Delta_{\varepsilon(k)}^*$ is an one loop open string worldsheet) with Neumann boundary conditions on each end, and let us suppose there is a quark q at one end of the string, and an antiquark

$\bar{q}$ at the other end, where $q^i, \bar{q}^i = 1, \dots, N$ are N -valued labels associated to the some representations of a gauge group G . Equations of motions for q_i simply require they to be independent from the coordinate parametrizing the boundary, thus they can be interpreted as charged, conserved under translations along the boundary. The target space action automatically has $SO(N)$ symmetry when the charges are real and $SO(2N)$ when they are complex. As a result, quantization produces an associated Hilbert space of states on which a spinor representation of respectively $SO(N)$ or $SO(2N)$ acts. To restrict to a subgroup of these two groups, (*i.e.* to a subrepresentation on which they act) it is necessary to reduce the Hilbert space. This can be achieved introducing a Lagrangian multiplier in the action. Unitarity of amplitudes actually restricts the admissible Lie algebras associated to such terms to belong to the classical series, thus admissible associated groups are $U(N)$, $SO(N)$ or $Sp(N)$.

Let us restrict to $G = U(N)$, with q^i in the defining representation N . Quantization of q^i generates N conserved charges attached to the propagating endpoints. The string for $U(N)$ should be oriented, since the charges at both ends of the string transform under inequivalent representations of the gauge group. The full string states now transform under the $N \otimes \bar{N}$ representation, namely the adjoint of $U(N)$. Thus, the generators T^a , $a = 1, \dots, \dim U(N) = N^2$ labels the string states belonging to the $N \times \bar{N}$ representation. The matrix elements $(T^a)_j^i$ specify which charges $\bar{q}^i$ and q_j are created at strings endpoints. The Fock space built over each ground state $|0, k; ij\rangle$ is now given by $\mathcal{H}^k \otimes N \otimes \bar{N}$.

For $G = U(N)$, $N \otimes \bar{N}$ transforms in the adjoint representation, thus massless open string states are natural candidates for Yang-Mills vector bosons with gauge group $U(N)$. The same construction might have been done for q^i in another representations of $U(N)$, but, in these cases, string states would have not rotated in the adjoint representation of the group, letting the above interpretation fail. Thus, the defining representation is singled out by its relevance in view of Yang-Mills particle interpretation of string states.

The string for $SO(N)$ or $Sp(N)$ should be unoriented, since charges at both ends transform under the same representation of the group. If we consider f to be the defining representation of $SO(N)$, we obtain string states in the adjoint of $SO(N)$ considering the only the antisymmetric part of the tensor product of the two representation, namely $(f \otimes f)_A$. On the other hand we obtain the string states in the adjoint of $Sp(N)$ by the symmetrized product of its defining representation with itself. In each case we recover exactly the the string states suitable for the Yang-Mills particles interpretation.

The effect of a general background gauge field is accounted for by including, for each boundary of the Polyakov path integral, a Wilson line term $\text{Tr } P \exp(-S_A)$, where S_A represents a boundary condensate of photon vertexes:

$$S_A = \int d\tau A_\alpha \partial_\tau X^\alpha \quad (4.6)$$




Figure 4.1: The ribbon graph associated to the open surface dual to three-punctured sphere

Preservation of conformal invariance impose the associate β -functions to vanish. Equation $\beta_A = 0$ reduce, to the leading order in the σ -model expansion, to the Yang-Mills equation.

In our connection, let us suppose that each cylindrical end's open string is decorated with a suitable assignation of Chan Paton factors. Thus, we can repeat the previous construction for the theory defined over each cylindrical end. Let us assign to each open string running one loop a suitable decoration. In this way, each vertex operator of the open string spectrum will get as a prefactor a $U(N)$ generator.

Now let us think to glue together the various cylindrical ends along the pattern defined by the ribbon graph. Some considerations naturally arise. Let us look again at figure 3.1 on page 62: since the two open string worldsheet boundaries which get glued along one edge of the ribbon graph pattern have opposite orientation, it seems natural to let the associated quarks to fall into opposite representations of the gauge group. The same holds for each edge of the ribbon graph, which thus acquire a well defined gauge coloring. Thus, we can think to the ribbon graph underling the triangulation as a sort of image of the gauge coloring of the boundaries of the surface ∂M .

About the spectrum, we can think to define the gluing process once again introducing a unique spectra of the common boundary theory: the (p, q) open string will be describe by a unique object being got by a sort of "fusion process" of the Chan-Paton factors of the original strings and which retains the open string data associated to the the free ends of the open unglued strings. However, since the fact that string states rotate into the adjoint representation of the gauge group is fundamental to their limiting description as particles, we want an object obtained by the product of the two original Chan-Paton factors and belonging to the original algebra. The only products obeying this constraint are the symmetric and antisymmetric products of generators, thus each BIO belonging to the (p, q) BCFT spectrum will be decored by a $U(N)$ generator T_{il}^a

defines as:

$$T_{il}^a(p, q) = f^{abc} [T_{ij}^b(p), T_{jl}^c(q)] + d^{abc} \{T_{ij}^b(p), T_{jl}^c(q)\} \quad (4.7)$$

Denoting the conformal structure of BIOs in formula (3.52) with a collective index $\Xi(p, q)$, non-Abelian BIOs will be $\psi_{\Xi(p, q)}^a = T^a \psi_{\Xi(p, q)}$. As a consequence, two and three-point functions of BIOs get a prefactor given by a trace of generators.

Let us notice that the above construction allows to reissue the description of the open string gauge data transmitting all the informations on the associated ribbon graph. Naively speaking, to each gauge coloring of ∂M boundaries we can uniquely associate a gauge coloring of the underlying ribbon graph “displaying” the $U(N)$ charges left after the gluing procedure from the outer to the inner boundary of each cylindrical end (see fig. 4.1).

In this connection, it is easy to understand that only the two points function between opposite directed BIOs (in the sense defined in appendix B) makes sense. In this case, the two point function will be simply weighted by the $U(N)$ generators normalization constant. The modified BIOs’ algebra can be easily retrieved by (anti)symmetrizing the product of generators:

$$\begin{aligned} \psi_{\Xi_1(r, p)}^a(\omega_r) \psi_{\Xi_2(q, r)}^b(\omega_q) &\sim \frac{1}{2} \sum_{\Xi_3, c} |\omega_r - \omega_q|^{H(q, p) - H(r, p) - H(q, r)} \times \\ &\quad (if^{abc} + d^{abc}) \mathfrak{C}(\Xi_1(r, p), \Xi_2(p, q), \Xi_3(q, r)) \psi_{\Xi_3(q, p)}^c(\omega_q), \end{aligned} \quad (4.8)$$

Thus, three-point functions will be decorated by the sum of the antisymmetric plus symmetric structure constants.

4.2 Rephrasing the geometrical data of the triangulation in term of string quantities

The overall picture we are dealing with sees a stack of N coincident D- p -branes winding around some (or all) of cycles of the toroidal background. In this connection, we are considering cylindrical ends as one-loop open string worldsheets. As prototype picture, we can consider the first drawing in fig. 4.2. Red circles are the worldvolumes of N coincident D-0-branes, while the green strip is the pictorial representation of a cylindrical ends injected by the maps X and Y on the two dimensional torus. In particular, looking at the cylindrical end as a one loop open string worldsheet, the associated open string winds one time around the X direction, while it does not wind around Y . Thus, the injection maps are characterized by the following values of winding numbers and center of mass




Figure 4.2: A cylindrical end winding in a two dimensional background

momentum:

$$\mu_{(O)}^X = 0 \quad \nu_{(O)}^X = 1 \quad (4.9a)$$

$$\mu_{(O)}^Y = 1 \quad \nu_{(O)}^Y = 0 \quad (4.9b)$$

Performing a worldsheet-duality transformation, thus looking at the closed string which is emitted by the brane and it is reabsorbed after having run around the X direction once, actually we exchange the role of total momentum and winding, thus the Hilbert space states are characterized by the following values of quantum numbers:

$$\mu_{(C)}^X = 1 \quad \nu_{(C)}^X = 0 \quad (4.10a)$$

$$\mu_{(C)}^Y = 0 \quad \nu_{(C)}^Y = 1 \quad (4.10b)$$

In this connection, computation of the target space spectrum shows that, when we consider an open string whose endpoint lay on a $D-p$ -brane, the background fields depends only upon coordinates on the brane world-volume (*i.e.* on coordinate ξ^μ , $\mu = 0, \dots, p$ parametrizing the brane embedding in the target space). Moreover, massless states vertex operators spacetime components actually divide in two set: those of the first set are components of a Yang-Mills field living on the $D-p$ -brane world-volume, with $p+1$ components tangent to the hyperplane, $A^i(\xi^\mu)$, $\mu, i = 0, \dots, p$, while the others $\phi^m(\xi^\mu) = \phi_b^m T^b$ are $N \times N$ matrices, whose entries are scalars field from the brane world volume point of view. They describe the specific shape of the brane in the target space.

In ordinary (bosonic) string theory, D -branes are dynamic objects, and such that, they must be able to respond to the values of the various background fields in the theory. This is obvious if one consider that actually brane's location and shape is controlled by the various open strings ending on them, and which indeed interact with background fields. A world-volume action describing this dynamic has been derived exploiting T -duality in target space (for a comprehensive review, see [95], chapter 5). If we consider a $D-p$ -brane in ordinary

D -dimensional open bosonic string theory, it turns out to be, in the Abelian case,

$$S_p = -T_p \int d^{p+1}\xi \epsilon^{-\Phi} \det^{\frac{1}{2}} [G_{ij} + B_{ij} + 2\pi\alpha' F_{ij}] \quad (4.11)$$

In equation 4.11, G_{ij} and B_{ij} are the pull-back of background fields to the brane, α' is the Regge slope and T_p is the D-brane tension defined as:

$$T_p = \frac{\sqrt{\pi}}{16\kappa_0} (4\pi^2 \alpha')^{\frac{11-p}{2}} \quad (4.12)$$

The non-Abelian case is quite more subtle, because the various background fields typically depends on the transverse coordinates, which, as stated before, are actually matrix valued in the gauge algebra. Thus, introduction of non-Abelian quantities, implies to perform a trace on the integrand, in order to get a gauge invariant object as action. It has been shown that considering the symmetrized trace over the gauge indexes, we get a result consistent with various studies of both scattering amplitudes and non-Abelian soliton solutions (see [95], section 5.5, and references therein). Thus, according to these remark, it has been shown that the worldvolume action associated to N coincident D- p -branes is:

$$S_p = -T_p \int d^{p+1}\xi \text{STr} \left\{ \det^{\frac{1}{2}} [E_{ij} E_{im} (Q^{-1} - \delta)^{mn} E_{nj} + 2\pi\alpha' F_{ij}] \det^{\frac{1}{2}} [Q_m^n] \right\} \quad (4.13)$$

where E is the background matrix introduced in (2.28), and $Q_m^n = \delta_m^n + i2\pi\alpha' [\phi^n, \phi^p] E_{pm}$.

If we expand this Lagrangian to the second order in the gauge field, and noting that

$$\det[Q_m^n] = 1 - \frac{(2\pi\alpha')^2}{4} [\phi^n, \phi^m] [\phi^n, \phi^m] + \dots \quad (4.14)$$

we can write the leading order of action (4.13) as

$$S_p = -\frac{T_p(2\pi\alpha')^2}{4} \int d^{p+1}\xi e^{-\Phi} \text{Tr} \left[F_{ab} F^{ab} + 2\mathcal{D}_a \phi^m \mathcal{D}^a \phi^m + [\phi^m, \phi^n]^2 \right]. \quad (4.15)$$

This is exactly the dimensional reduction of a D -dimensional Yang-Mills Lagrangian, displaying the non trivial scalar potential involving commutator for the adjoint scalars. This expansion allows to write the $(p+1)$ -dimensional Yang-Mills coupling for the theory on the brane:

$$g_{YM,p}^2 = g_s T_p^{-1} (2\pi\alpha')^{-2} \quad (4.16)$$

where $g_s = e^\Phi$ is the usual worldsheet string coupling.

Let us turn to the Polyakov string defined on each cylindrical end via the action in (2.29). We can easily rewrite it in a stringy shape by introducing




Figure 4.3: Some simple configurations with three cylindrical ends winding in a two dimensional flat toroidal background

dimensionful coordinates via the mapping $X^\alpha(\zeta(k), \bar{\zeta}(k)) \rightarrow \frac{l(k)}{2} X^\alpha(\zeta(k), \bar{\zeta}(k))$. Polyakov action becomes:

$$S(k) = \frac{1}{2\pi l(k)^2} \int d\zeta(k) d\bar{\zeta}(k) \partial X^\alpha \bar{\partial} \bar{X}_\alpha \quad (4.17)$$

thus suggesting the parameter $l^2(k)$ to play a role analogue to that of the Regge slope in the discrete (k) -sector. However, from a classical point of view, since the above analogy has been derived from a fixed triangulation of the worldsheet, we are not able to associate it a real physical meaning. Moreover, if we consider the collection of all the discrete sectors, the above identification seems to lead to a strange non-sense. As a matter of fact, if we consider the full surface M_∂ , we deal with cylindrical ends having their inner boundaries glued together along the pattern defined by the ribbon graph, while outer ones lay on (at least) one stack of N tiled D-branes.

In particular, let us assume exactly the situation depicted in fig. 4.3. We are considering again the open surface dual to the three-punctured sphere and which has been sketched in fig. 4.1. In the left drawing each string has zero winding number in both direction, while, in the right one, each string winds once around X and does not wind around Y . Moreover, let us assume that the outer boundaries of each cylindrical end lays on the same stack of D-branes. According to the above remarks about the role played by the parameter $l(k)$, it may seem strange that each copy of the model, each being defined over one cylindrical end, has associated an independent value of the “ α' -analogue” parameter. It may seem meaningless especially if we consider the cylindrical ends boundaries laying on the same D-brane, whose tension is defined in term of α' via the relation (4.12).

An elegant solution of this puzzle can be achieved if we consider the problem from a “quantistic” point of view: the construction of the partition function for the bosonic string involve a sum over all possible configurations *i.e.* over all possible ribbon graphs. In this framework clearly we are considering an

average over all possible parameters associated to the triangulation, and thus the Regge slope analogue is not a fixed value associated to each cylindrical end but it arises as a statistical average over all possible configurations. A further development arises if we perform the continuous limit of such a discrete model. For discrete planar surfaces this is usually performed as limit in which the lengths of boundary components go to infinity, while a suitable cutoff a goes to zero, in order to maintain the product $L(k)a$ finite. Thus, if we identify on each cylindrical end the (statistical average) of the characteristic length $l(k)$ with:

$$l(k) = \sqrt{L(k)a} \quad (4.18)$$

and we perform the continuum limit requiring that all the (statistical averages) of finite values of the above products converge to a given value, we can actually interpret such a result as an analogue of the Regge slope which, according to relation (4.12), determine the tension of the brane naturally included in our model.

Moreover, this defines a precise map between the geometrical data associated to the triangulation and string quantities.

4.3 Coupling with background gauge potential: extending the gluing procedure

Let us consider a D -dimensional background, in which each direction X^α , $\alpha = 0, \dots, D-1$ is compactified at the T-duality self dual radius $\Omega^\alpha = \sqrt{2}$. As explained in the last chapter, the theory in each direction decouples and the model presents a $(SU(2)_L \times SU(2)_R)^D$ gauge symmetry. In each direction we can apply previous chapter's constructions. Thus, let us consider the k -th cylindrical end, $\Delta_{\varepsilon(k)}^*$, and let us interpret it as worldsheet associated to one-loop open string diagram. Consistently with the gluing process, let us assume the injection maps (target space coordinates) to obey simple Neumann boundary conditions on the inner boundary. On the other hand, on the outer boundary, let us assume to have $p+1$ directions satisfying Neumann boundary conditions and $D-p-1$ directions obeying Dirichlet's ones:

$$\{X^\alpha\} \doteq \{X^i, X^m\}, \text{ with } i = 0, \dots, p, m = p+1, \dots, D-1 \quad (4.19)$$

According to formulae (3.36) and (3.43), these can be characterized by:

$$\text{Neumann directions : } (\|N(0)_{(k)}\rangle\rangle_{sd}, \mathbb{I}) \quad (4.20)$$

$$\text{Dirichlet directions : } (\|N(0)_{(k)}\rangle\rangle_{sd}, e^{-i\pi J_0^1(k)}) \quad (4.21)$$

In particular, with the above choice, we are dealing with a D - p -brane laying along the $X^0, \dots, X_p$ directions, thus, if we introduce the D -brane world-volume

parameters $\xi_0, \dots, \xi_p$, we have chosen to parametrize the brane world-volume with $\xi_0 = X_0, \xi_1 = X_1, \dots, \xi_p = X_p$.

As described in the previous section, in order to endow the brane with an interesting dynamic, we have to couple the model to a background gauge field living on its worldvolume: this can be worked out introducing non chiral truly marginal deformation defined by the following boundary action ([98, 99]):

$$S_A = \int d\tau \left[\sum_{i=0}^p A_i(X^i) \partial_\tau X^i + \sum_{m=p+1}^{D-1} \phi_m(X^i) \partial_\sigma X^m \right], \quad (4.22)$$

where we have taken the boundary to lay at constant σ . The $A_i = A_i^a T^a$ are Lie algebra valued gauge fields on the D-brane, while the entries of the $N \times N$ matrices $\phi_m = \phi_m^a T^a$ are scalar from the world volume point of view: they describe the motion of the brane in the transverse space. Such a boundary term will act as a non chiral deformation of the original quantum model by means of a truly marginal operator.

For the sake of simplicity, let us assume the brane static in the transverse space imposing $\phi_m = \mathbf{0}_{N \times N} \forall m = p+1 \dots D-1$. Moreover, let us consider constant electric and magnetic fields along the brane worldvolume, so that the boundary term reads:

$$S_A = F_{ij} \int d\tau X^j \partial_\tau X^i. \quad (4.23)$$

Let us move to the Abelian subsector. Such a limiting situation can be achieved including in each Neumann direction of the T-dual theory a Wilson line $A_i = \frac{\text{diag}\{\theta_1, \dots, \theta_N\}}{\Omega_i}$ which breaks $U(N)$ to $U(1)^N$. At spacetime level, the global effect will be a displacement of the position of the N D-branes. Thus, now we are dealing with N separated D-branes, and we can assume that open strings generating the cylindrical ends have their outer endpoint attached over different brane.

Thus, in each (k) -subsector, we can couple the open string with a different electromagnetic potential $A_i(k)$, allowing the boundary term to have a simple closed string interpretation. As a matter of fact, the Lagrangian in (4.23) is the integral of a total derivative, thus, after a short computation, S_A can be rewritten as:

$$S_{A(k)} = F_{ij}(k) \int d\zeta(k) d\bar{\zeta}(k) \partial X^i(k) \bar{\partial} \bar{X}^j(k). \quad (4.24)$$

Comparing last formula with formula (2.27), we can state that, in the Abelian subsector, the inclusion of such a boundary term is equivalent to moving to a different point in flat toroidal background moduli space (see formula (3.68)):

Config. $\mathcal{A}$	Config. $\mathcal{B}$
$G_{\alpha\beta} = \mathbb{I}_{D \times D}$	$G_{\alpha\beta} = \mathbb{I}_{D \times D}$
$B_{\alpha\beta} = 0 \forall \alpha, \beta$	$B_{\alpha\beta} = 4\pi \Lambda_{\alpha\beta}$
$F_{\alpha\beta} = \Lambda_{\alpha\beta}$	$F_{\alpha\beta} = 0 \forall \alpha, \beta$

On this wise, we can directly choose a peculiar flat toroidal background, with associated suitable values of the background matrix E entries (see formula (2.28)), and rephrase the B -field dependence in term of a gauge field strength associated to a potential living on the D-brane world-volume.

As explained in the previous chapter, the full moduli space of flat toroidal backgrounds is parametrized by the coset space:

$$O(D, D, \mathbb{Z}) / O(D, D, \mathbb{R}) \setminus O(D) \times O(D). \quad (4.25)$$

In the previous formula $O(D, D, \mathbb{Z})$ is the generalized T-duality group.

Thus, higher dimensional toroidal compactifications are described by non-trivial background fields B and G and, in such a given background, the maximally enhanced symmetry points are those fixed under the action of $O(D, D, \mathbb{Z})$, which elements are generated by a combination of:

- a conjugations with a matrix $M \in SL(d, \mathbb{Z})$;
- an integer shift of B by means of antisymmetric, integer valued, matrix Θ ;

i.e. the rotated background matrix will be

$$E' = M^t (E + \Theta) M \quad (4.26)$$

In those special points of the moduli space in which $E' = E$, we can obtain extended target space symmetries with respect to semi-simple products of simply laced Lie algebras (thus belonging to the A-D-E decomposition) of total rank D .

In particular, the maximally enhanced symmetry background can be chosen in the following way[96]. If $C_{\alpha\beta}$, $\alpha, \beta = 1, \dots, D$, is the Cartan matrix of the simply laced Lie algebra of total rank D , then the background fields are chosen as:

$$G_{\alpha\beta} = \frac{1}{2} C_{\alpha\beta} \quad (4.27a)$$

$$B_{\alpha\beta} = G_{\alpha\beta} \quad \forall \alpha > \beta \quad B_{\alpha\beta} = -G_{\alpha\beta} \quad \forall \alpha < \beta \quad B_{\alpha\alpha} = 0 \quad \forall \alpha \quad (4.27b)$$

With these choices, the background matrix $E = G + B$ is an element of $SL(d, \mathbb{Z})$, thus it is fixed under the action of $O(D, D, \mathbb{Z})$.

Let us consider, as an example, an $O(D, D, \mathbb{Z})$ transformation generated by $M = E^{-1}$ and $\Theta = E^\dagger + E$. The duality map states (4.26) tells: $E' = E^{-1}$ and, if $G = \mathbb{I}_{D \times D}$ and $B = \mathbf{0}_{D \times D}$, this is exactly the case of $(SU(2)_L \times SU(2)_R)^D$.

With the above choice of static gauge, this is exactly the situation for the $D - p - 1$ Dirichlet directions. Thus, the extended symmetry group associated to the action (4.24) will be:

$$\mathbb{G}_D = (G_{p+1} \times G_{p+1}) \times (SU(2)_L \times SU(2)_R)^{D-p-1} \quad (4.28)$$

4.3.1 Chiral currents and boundary states

As we did for one compactified boson at the self dual radius, the level-one untwisted Kac-Moody algebra arising in this connection can be represented in term of the bosonic coordinate by means of the usual vertex operators construction[80]. In the cylinder transverse channel (*i.e.* in terms of coordinate $\zeta(k)$ and $\bar{\zeta}(k)$ on the k -th cylindrical end), the left moving and right-moving currents are, respectively (let us omit the polytopal index k)

$$H^\alpha(\zeta) = \partial X^\alpha, \quad E^\lambda(\zeta) = c(\lambda) :e^{i\lambda_\alpha X^\alpha}: \quad (4.29a)$$

$$\bar{H}^\alpha(\bar{\zeta}) = M_\beta^\alpha \bar{\partial} \bar{X}^\beta, \quad \bar{E}^\lambda(\bar{\zeta}) = -\bar{c}(\lambda) :e^{i\lambda_\alpha M_\beta^\alpha \bar{X}^\beta}:. \quad (4.29b)$$

Here $H^\alpha(\zeta)$ and $\bar{H}^\alpha(\bar{\zeta})$ denotes the elements in the Cartan subalgebra of $\mathfrak{g}_{Dk=1}$, while $\{\lambda\}$ is the set of roots (positive plus negative) of the associated semi-simple product of Lie algebras. The objects $c(\lambda)$ and $\bar{c}(\lambda)$ are $\mathbb{Z}_2$ values cocycles. They are operators acting on the Fock spaces, and they depend only upon the momentum part of the free-boson zero modes. Their inclusion let the combinations of the above currents, $J_L^\alpha(\zeta)$, to satisfy the correct OPE:

$$J^a(\zeta)J^b(\zeta') \sim \frac{\delta^{ab}}{(\zeta - \zeta')^2} + \sum_c i f^{abc} \frac{J^c(\zeta')}{\zeta - \zeta'} \quad (4.30)$$

If we introduce the Laurent expansion, $J^a(\zeta) = \sum_{n \in \mathbb{Z}} \zeta^{-n-1} J_n^a$, we get the commutation relation for the affine algebra $\mathfrak{g}_{Dk=1}$:

$$[J_m^a, J_n^b] = \sum_c i f^{abc} J_{m+n}^c + m \delta^{ab} \delta_{m+n,0}. \quad (4.31)$$

Obviously, the same holds for the antiholomorphic sector.

Limiting to the trivial automorphism in equation (2.15), we get the following constraints on the boundary states:

$$(J_n^a + \bar{J}_{-n}^a) \|B\rangle\rangle \quad (4.32)$$

Vertex operators for the closed string massless vector states are the D-dimensional extensions of formula (3.31):

$$V_P^{a\alpha} = J^a(\zeta) \bar{\partial} \bar{X}^\alpha(\bar{\zeta}) e^{iP \cdot X(\zeta) + \bar{X}(\bar{\zeta})} \quad (4.33)$$

$$\bar{V}_P^{a\alpha} = \bar{J}^a(\bar{\zeta}) \partial X^\alpha(\zeta) e^{iP \cdot X(\zeta) + \bar{X}(\bar{\zeta})} \quad (4.34)$$

where $P = p_L + p_R$ is the total center of mass momentum of the closed string. Once again, vertex operators for new open string scalar states attached to the boundaries $|\zeta| = 1, |\bar{\zeta}| = \frac{2\pi}{2\pi - \varepsilon(k)}$ can be written, in the closed string channel, as:

$$S_P^a = J^a(x) e^{iP X(x)}, \quad (4.35)$$

where, as usual, we have parametrized the boundary with $x \doteq \Re z(k, \circ)[\zeta(k)]$.

4.3.2 Gluing along the Ribbon Graph: a proposal

The occurrence of extra massless boundary operators (4.35) at the enhanced symmetry point can be obviously rephrased as the arising of new truly-marginal perturbing boundary operators, which give rise to the enhanced symmetry also in the open string spectrum. Such operators come associates with perturbing terms in the action of the form:

$$S'_g = \int dx \left(\sum_{\lambda} g_{\lambda} e^{i\lambda_{\alpha} X^{\alpha}} + \sum_{\alpha} g_{\alpha} \partial_x X^{\alpha} \right) |_{|\zeta(k)|=1}, \quad (4.36)$$

where $(g_{\lambda}, g_{\alpha})$ are real coupling constant, which give rise to the enhanced symmetry in the open sector when λ_i takes special values. In [100] it was shown that these particular values arise when we can define a set of vectors:

$$\hat{\lambda}_{\alpha} = (\delta_{\beta}^{\alpha} + M_{\beta}^{\alpha}) \lambda^{\beta} \quad (4.37)$$

(where $M = \frac{G+B}{G-B}$), such that the $\hat{\lambda}$ are roots of the simply-laced algebra $\mathfrak{g}_D$. According to deformation rules of previous chapter, this lead to a modified gluing condition on the boundary:

$$J^a(\zeta) = \gamma_{g_a \bar{J}_0^a}(\bar{J}^a)(\bar{\zeta}), \quad (4.38)$$

while the effect on the boundary state is a rotation with respect to the left-moving zero modes of the currents:

$$\|B\rangle\rangle_g = e^{g_{\hat{\lambda}} E_0^{\hat{\lambda}} + g_{\alpha} H_0^{\alpha}} \|B\rangle\rangle_g \implies \|B\rangle\rangle_g = e^{\sum_a g_a J_0^a} \|B\rangle\rangle \quad (4.39)$$

This construction, perfectly equivalent to the one arising when we deal with only one injection map, allows us to make a concrete hypothesis about the possible parametrization of boundary states in this higher dimensional model. If we are able to identify a generating boundary state $\|B_0\rangle\rangle$ equivalent to $\|N(0)\rangle\rangle_{sd}$, we can hypothesize to identify Cardy's boundary states associated to the level one $\mathfrak{g}_D$ WZW-model with the deformations of $\|B_0\rangle\rangle$ with central elements of the group $\mathbb{G}$. Once proved this statement, we could apply the same construction of previous chapters to write an open string amplitude over ∂M , with the particular choice of background fields in each cylindrical end given by (4.27).

If we consider the simpler case of $\mathbb{G}$ being a simple Lie group, strong indications of the above come if we notice that the number of elements of the (discrete) center of $\mathbb{G}$, $B(\mathbb{G})$, coincide with the number of level one modules associated to the (simply-laced) simple Lie algebra $\mathfrak{g}_D$. Moreover the center of the group is isomorphic to the group of outer automorphisms associated to the

affine simply laced Lie algebra $\hat{\mathfrak{g}}_D$ ¹:

$$\mathcal{O}(\hat{\mathfrak{g}}_D) = B(\mathbb{G}). \quad (4.41)$$

Remembering that $\mathcal{O}(\hat{\mathfrak{g}}_D)$ maps the set of dominant weights which label the irreducible representations of the affine algebra, this suggests the possibility to generate the full set of Cardy boundary states associated to the (level one) $\hat{\mathfrak{g}}_D$ WZW model via the action of elements of $B(\mathbb{G})$ on the boundary state associated to the identity element of $\mathbb{G}$.

4.4 Conclusions and perspectives

In spirit of understanding the (mathematical and physical) origins of open/closed string dualities, in previous sections we have investigated some aspects of coupling of a bosonic string theory with the peculiar geometry arising when we uniformize the singular Euclidean structure naturally associated with the Regge polytope barycentrically dual to a Random Regge Triangulation.

The worldsheet process guiding the holes glueing has been made precise only at a topological level, while in the more complicated *AdS/CFT* framework we are still far from understanding the origin of such a behavior pattern.

In this sense, a remarkable hint has been recently given by Gopakumar, who intuited that such a process must take place as a change of variable at level of integrand in the sum over moduli space parametrizing the free field $\mathcal{N} = 4$ SYM correlators[63] and suggested a concrete way of associating a closed Riemann surface to a gauge theory correlator. However, the argument introduced by Gopakumar is quite general, since up to now it has been applied only to the free field theory sector, thus leaving open the question about a possible application to interacting theories.

In this connection, it is straightforward to notice that the key object is represented by the underlying geometrical structure, namely the skeleton graph naturally associated to gauge correlator and which can be considered as dual to a triangulation of the Riemann surface defining the closed string worldsheet. In this connection, in view of the two dual uniformizations of the Euclidean structure naturally associated to a RRT and introduced in [61, 60] which provide a concrete algorithm to switch from an open to a closed Riemann surface, it seems that a better understanding of the interplay between geometrical quantities associated to a discrete realization of the worldsheet and the analytic

¹As a matter of fact, the relation between $\mathcal{O}(\hat{\mathfrak{g}}_D)$ and $B(\mathbb{G})$ is even stronger because if we consider an element $A \in \mathcal{O}(\hat{\mathfrak{g}}_D)$, we can define the associated element $b \in B(\mathbb{G})$ as the $\mathcal{S}^{ext}$ dual to A :

$$b = \mathcal{S}^{ext\dagger} A \mathcal{S}^{ext} \quad (4.40)$$

quantities of string theory, may provide a key step towards the understanding of mathematical aspects of string dualities.

The main goal of previous section has been the description of a very reliable way of coupling an ordinary bosonic open string theory with the discrete geometry defined by the uniformization of M_{∂} . In particular, exploiting BCFT techniques, we have been able to furnish a concrete proposal about how to include open string gauge degrees of freedom in our model. Not less important, we have been able to define a natural interplay between the geometrical quantities parametrizing the uniformization defined over M_{∂} and the Regge slope α' .

Wide investigation areas are still open. The inclusion of open string gauge degrees of freedom itself is not completely fixed. In the abelian case the statement sketched at the end of section 4.3.2 needs some further investigations. Moreover, nevertheless the nice picture presented in section 4.2, the extension of the above proposal to the non-Abelian case is quite far from being understood.

Furthermore, a very interesting point to address would be investigating the possible extending of the above glueing procedure to minimal string theories[101, 102], which, despite of their simplicity, they are interesting laboratories for the study of string theories.

Part II

COSMOLOGICAL BACKGROUNDS OF SUPERSTRING THEORY: OXIDATION AND BRANES

Chapter 5

Geodesic on maximally non compact cosets U/H and differential equations

We have recalled how both maximally extended supergravities (of type A and B) reduce, stepping down from $D = 10$ to $D = 3$ to the following non linear sigma model coupled to $D = 3$ gravity:

$$\mathcal{L}^{\sigma-model} = \sqrt{-\det g} \left[2 R[g] + \frac{1}{2} h_{IJ}(\phi) \partial_\mu \phi^I \partial_\nu \phi^J g^{\mu\nu} \right] \quad (5.1)$$

where h_{IJ} , $I, J = 1, \dots, 128$, is the metric of the 128-dimensional homogeneous coset manifold

$$\mathcal{M}_{128} = \frac{E_{8(8)}}{SO(16)} \quad (5.2)$$

The above manifold falls in the general category of manifolds U/H such that $\mathbb{U}$ (the Lie algebra of U) is the maximally non-compact real section of a simple Lie algebra $\mathbb{U}_C$ and the subgroup H is generated by the maximal compact subalgebra $\mathbb{H} \subset \mathbb{U}$. In this case the solvable Lie algebra description of the target manifold U/H is universal. The manifold U/H is isometrical to the solvable group manifold:

$$\mathcal{M}_d = \exp [Solv(U/H)] \quad (5.3)$$

where the solvable algebra $Solv(U/H)$ is spanned by all the Cartan generators $\mathcal{H}_i$ and by the step operators E^α associated with all positive roots $\alpha > 0$ (on the solvable Lie algebra parametrization of supergravity scalar manifolds see [47, 48]). On the other hand the maximal compact subalgebra $\mathbb{H}$ is spanned by all operators of the form $E_\alpha - E_{-\alpha}$ for all positive roots $\alpha > 0$. So the dimension of the coset d , the rank r of $\mathbb{U}$ and the number of positive roots p are generally

related as follows:

$$\dim[U/H] \equiv d = r + p \quad ; \quad p \equiv \# \text{positive roots} = \dim \mathbb{H} \quad ; \quad r \equiv \text{rank } \mathbb{U} \quad (5.4)$$

In the present chapter we concentrate on studying solutions of a bosonic field theory of type (5.1) that are only time-dependent. In so doing we consider the case of a generic manifold U/H and we show how the previously recalled algebraic structure allows to retrieve a complete generating solution of the field equations depending on as many essential parameters as the rank of the Lie algebra $\mathbb{U}$. These parameters label the orbits of solutions with respect to the action of the two symmetries present in (5.1), namely U global symmetry and H local symmetry. The essential observation is that, as long as we are interested in solutions depending only on time, the field equations of (5.1) can be organized as follows. First we write the field equations of the matter fields ϕ^I which supposedly depend only on time. At this level the coupling of the sigma-model to three dimensional gravity can be disregarded. Indeed the effect of the metric g_{00} is simply that the field equations for the scalars ϕ^I have the same form as they would have in a rigid sigma model with just the following proviso. The parameter we use is proper time rather than coordinate time. Next in the variation with respect to the metric we can use the essential feature of three-dimensional gravity, namely the fact that the Ricci tensor completely determines also the Riemann tensor. This means that from the stress energy tensor of the sigma model solution we reconstruct, via Einstein field equations, also the corresponding three dimensional metric.

5.1 Decoupling the sigma model from gravity

Since we are just interested in configurations where the fields depend only on time, we take the following ansatz for the three dimensional metric:

$$ds_{3D}^2 = A^2(t) dt^2 - B^2(t) (dr^2 + r^2 d\phi^2) \quad (5.5)$$

where $A(t)$ and $B(t)$ are undetermined functions of time. Then we observe that one of these functions can always be reabsorbed into a redefinition of the time variable. We fix such a coordinate gauge by requiring that the matter field equations for the sigma model should be decoupled from gravity, namely should have the same form as in a flat metric. This will occur for a special choice of the time variable. Let us see how.

In general, the sigma model equations, coupled to gravity, have the following form:

$$\square_{cov} \phi^I + \Gamma_{JK}^I \partial_\mu \phi^J \partial_\nu \phi^K g^{\mu\nu} = 0 \quad (5.6)$$

In the case we restrict dependence only on time the above equations reduce to:

$$\frac{1}{\sqrt{-\det g}} \frac{d}{dt} \left(\sqrt{-\det g} g^{00} \frac{d}{dt} \phi^I \right) + \Gamma_{JK}^I \frac{d}{dt} \phi^J \frac{d}{dt} \phi^K g^{00} = 0 \quad (5.7)$$

We want to choose a new time $\tau = \tau(t)$ such that with respect to this new variable equations (5.7) take the same form as they would have in a sigma model in flat space, namely:

$$\ddot{\phi}^I + \Gamma_{JK}^I \dot{\phi}^J \dot{\phi}^K = 0 \quad I, J, K = 1, \dots, \dim \mathcal{M} \quad (5.8)$$

where Γ_{JK}^I are the Christoffel symbols for the metric h_{IJ} . The last equations are immediately interpreted as geodesic equations in the target scalar manifold.

In order for equations (5.7) to reduce to (5.8) the following condition must be imposed:

$$\sqrt{-\det g} g^{00} \frac{d}{dt} = \frac{d}{d\tau} \Rightarrow dt = \sqrt{-\det g} g^{00} d\tau \quad (5.9)$$

Inserting the metric (5.5) into the above condition we obtain an equation for the coefficient $A(t)$ in terms of the coefficient $B(t)$. Indeed in the new coordinate τ the metric (5.5) becomes:

$$ds_{3D}^2 = B^4(\tau) d\tau^2 - B^2(\tau) (dr^2 + r^2 d\phi^2) \quad (5.10)$$

The choice (5.10) corresponds to the following choice of the *dreibein*:

$$e^0 = B^2(\tau) d\tau \quad ; \quad e^1 = B(\tau) dr \quad ; \quad e^2 = B(\tau) r d\phi \quad (5.11)$$

For such a metric the curvature 2-form is as follows:

$$\begin{aligned} R^{01} &= \frac{2\dot{B}^2(\tau) - B(\tau)\ddot{B}(\tau)}{B^6(\tau)} e^0 \wedge e^1 \\ R^{02} &= \frac{2\dot{B}^2(\tau) - B(\tau)\ddot{B}(\tau)}{B^6(\tau)} e^0 \wedge e^2 \\ R^{12} &= -\frac{\dot{B}^2(\tau)}{B^6(\tau)} e^1 \wedge e^2 \end{aligned} \quad (5.12)$$

The Einstein equations, following from our lagrangian (5.1) are the following ones, in flat indices:

$$2G_{ab} = T_{ab} \quad ; \quad G_{ab} \equiv \text{Ric}_{ab} - \frac{1}{2} R \eta_{ab} \quad ; \quad a, b = 0, 1, 2 \quad (5.13)$$

With the above choice of the vielbein, the flat index Einstein tensor is easily calculated and has the following form:

$$G_{00} = \frac{\dot{B}^2(\tau)}{2B^6(\tau)} ; \quad G_{0i} = 0 ; \quad G_{ij} = \frac{2\dot{B}^2(\tau) - B(\tau)\ddot{B}(\tau)}{2B^6(\tau)} \delta_{ij} ; \quad i, j = 1, 2 \quad (5.14)$$

On the other hand, calculating the stress energy tensor of the scalar matter in the background of the metric (5.10) we obtain (also in flat indices):

$$T_{00} = \frac{1}{2B^4(\tau)} \left(\dot{\phi}^I \dot{\phi}^J h_{IJ} \right) \quad ; \quad T_{0i} = 0 \quad ; \quad T_{ij} = \frac{1}{2B^4(\tau)} \left(\dot{\phi}^I \dot{\phi}^J h_{IJ} \right) \delta_{ij} \quad ; \quad i, j = 1, 2 \quad (5.15)$$

where

$$\dot{\phi}^I \dot{\phi}^J h_{IJ} = \varpi^2 \quad (5.16)$$

is a constant independent from time as a consequence of the geodesic equations (5.8). To prove this it suffices to take a derivative in τ of ϖ^2 and verify that it is zero upon use of eq.s (5.8).

Hence in order to satisfy the coupled equations of gravity and matter fields it is necessary that:

$$2 \frac{\dot{B}^2(\tau)}{2B^6(\tau)} = 2 \frac{2\dot{B}^2(\tau) - B(\tau)\ddot{B}(\tau)}{2B^6(\tau)} = \frac{1}{2B^4(\tau)} \varpi^2 \quad (5.17)$$

The first of the above equalities implies:

$$\dot{B} = k B \quad \Rightarrow \quad B(\tau) = \exp[k \tau] \quad (5.18)$$

where k is some constant. The second equality is satisfied if:

$$k = \pm \frac{1}{\sqrt{2}} |\varpi| = \pm \frac{1}{\sqrt{2}} \sqrt{\dot{\phi}^I \dot{\phi}^J h_{IJ}} \quad (5.19)$$

In this way we have completely fixed the metric of the three-dimensional space as determined by the solution of the geodesic equations for the scalar matter:

$$ds_{3D}^2 = \exp[4k\tau] d\tau^2 - \exp[2k\tau] (dx_1^2 + dx_2^2) \quad (5.20)$$

with the parameter k given by eq.(5.19).

5.2 Geodesic equations in target space and the Nomizu operator

Having clarified how the three dimensional metric is determined in terms of the solutions of the sigma model, we concentrate on this latter. We focus on the geodesic equations (5.8) and in order to study them, we rely on the solvable Lie group description of the target manifold going to an anholonomic basis for the tangent vectors to the geodesic. Since gravity is decoupled from the scalars, we deal with a rigid sigma-model where the fields depend only on time

$$\mathcal{L}^{\sigma-model} \propto h_{IJ}(\phi) \dot{\phi}^I \dot{\phi}^J \quad (5.21)$$

As was mentioned before, the equations of motion in this case reduce to the geodesic equations for the metric $h_{IJ}(\phi)$ and time plays the role of a parameter

along the geodesics (see eq.(5.8)). Since $h_{IJ}(\phi)$ is the metric of a scalar manifold which is a maximally non-compact coset $\mathcal{M} = \mathbb{U}/\mathbb{H}$, we can derive this metric from a coset representative $\mathbb{L}(\phi) \in U$

$$h_{IJ}(\phi) = \text{Tr}(\mathbb{P}_K \mathbb{L}^{-1} \partial_I \mathbb{L} \mathbb{L}^{-1} \partial_J \mathbb{L}) \quad (5.22)$$

$\mathbb{P}_K$ being a projection operator on the coset directions of the Lie algebra $\mathbb{U}$ to be discussed in a moment. To this effect we introduce the following general notation. We make the orthogonal split of the $\mathbb{U}$ Lie algebra:

$$\mathbb{U} = \mathbb{H} \oplus \mathbb{K} \quad (5.23)$$

where $\mathbb{H} \subset \mathbb{U}$ is the maximal compact subalgebra and $\mathbb{K}$ its orthogonal complement. We adopt the following normalizations for the generators in each subspace:

$$\begin{aligned} \mathbb{U} &= \text{Span}\{H_i, E_\alpha, E_{-\alpha}\} \\ \mathbb{K} &= \text{Span}\{K_A\} = \text{Span}\{H_i, \frac{1}{\sqrt{2}}(E_\alpha + E_{-\alpha})\} \\ \mathbb{H} &= \text{Span}\{t_\alpha\} = \text{Span}\{(E_\alpha - E_{-\alpha})\} \end{aligned} \quad (5.24)$$

The $\mathbb{U}$ Lie-algebra valued left invariant one-form

$$\Omega = \mathbb{L}^{-1} d\mathbb{L} = V^A K_A + \omega^\alpha t_\alpha \quad (5.25)$$

is in general expanded along all the generators of $\mathbb{U}$ (not only along $\mathbb{K}$) and $V = V^A K_A$ corresponds to the coset manifold vielbein while $\omega = \omega^\alpha t_\alpha$ corresponds to the coset manifold $\mathbb{H}$ -connection.

As it is well known neither the coset representative $\mathbb{L}(\phi)$, nor the one-form Ω are unique. Indeed $\mathbb{L}$ is defined up to multiplication on the right by an element of the compact subgroup $h \in H$. This is a gauge invariance which can be fixed in such a way that the coset representative lies in the solvable group $\text{Solv}(\mathbb{U}/\mathbb{H})$ obtained by exponentiating the solvable subalgebra $\text{Solv}(\mathbb{U}/\mathbb{H})$

$$\mathbb{L}(\phi) = \exp(\text{Solv}(\mathbb{U}/\mathbb{H})) \cdot \phi \quad (5.26)$$

In the case of $\mathbb{U}/\mathbb{H}$ being maximally non compact $\text{Solv}(\mathbb{U}/\mathbb{H})$ coincides with the Borel subalgebra and therefore it is spanned by the collection of all Cartan generators and step-operators associated with positive roots, as we already stated, namely:

$$\text{Solv}(\mathbb{U}/\mathbb{H}) = \text{Span}\{T_A\} = \text{Span}\{H_i, E_\alpha\} \quad (5.27)$$

If the coset representative $\mathbb{L}$ is chosen to be a solvable group element, as in eq.(5.26), namely if we are in the solvable parametrization of the coset, we can also write:

$$\Omega = \mathbb{L}^{-1} d\mathbb{L} = \tilde{V}^i \mathcal{H}_i + \tilde{V}^\alpha E_\alpha = \tilde{V}^A T_A \quad (5.28)$$

since Ω is contained in the solvable subalgebra $Solv(\mathbb{U}/\mathbb{H}) \subset \mathbb{U}$. Eq.s (5.25) and (5.28) are compatible if and only if:

$$V^\alpha = \sqrt{2} \omega^\alpha \quad (5.29)$$

In this case we can identify $\tilde{V}^i = V^i$ and $\tilde{V}^\alpha = \sqrt{2} V^\alpha$; eq.(5.29) is the solvability condition for a coset representative.

Hence we can just rewrite the metric of our maximally non-compact manifold $\mathbb{U}/\mathbb{H}$ as follows:

$$ds_{\mathbb{U}/\mathbb{H}}^2 = \sum_{A=1}^{\dim \mathbb{U}} V^A \otimes V^A = \tilde{V}^i \oplus \tilde{V}^i + \frac{1}{2} \tilde{V}^\alpha \oplus \tilde{V}^\alpha \quad (5.30)$$

It is interesting to discuss what are the residual $\mathbb{H}$ -gauge transformations that remain available after the solvable gauge condition (5.29) has been imposed. To this effect we consider the multiplication

$$\mathbb{L} \mapsto \mathbb{L} h = \overline{\mathbb{L}} \quad (5.31)$$

where

$$h = \exp [\theta^\alpha t_\alpha] \quad (5.32)$$

is a finite element of the $\mathbb{H}$ subgroup singled out by generic parameters θ . For any such element we can always write:

$$\begin{aligned} h^{-1} t_\alpha h &= A(\theta)_\alpha^\beta t_\beta \\ h^{-1} K_A h &= D(\theta)_A^B K_B \end{aligned} \quad (5.33)$$

where the matrix $A(\theta)$ is the adjoint representation of h and $D(\theta)$ is the D -representation of the same group element. We obtain:

$$\begin{aligned} \overline{\Omega} \equiv \overline{\mathbb{L}}^{-1} d\overline{\mathbb{L}} &= h^{-1} dh + h^{-1} \Omega h \\ &= \underbrace{h^{-1} dh + h^{-1} \omega h}_{=\overline{\omega}} + \underbrace{h^{-1} V h}_{=\overline{V}} \end{aligned} \quad (5.34)$$

where

$$\begin{aligned} \overline{\omega}^\alpha &= \frac{1}{\text{tr}(t_\alpha^2)} \text{tr} (h^{-1} dh t_\alpha) + \omega^\beta A(\theta)_\beta^\alpha \\ \overline{V}^\alpha &= V^\beta D(\theta)_\beta^\alpha + V^i D(\theta)_i^\alpha \end{aligned} \quad (5.35)$$

Suppose now that the coset representative $\mathbb{L}$ is solvable, namely it satisfies eq.(5.29). The coset representative $\overline{\mathbb{L}}$ will still satisfy the same condition if the h -compensator satisfies the following condition:

$$\frac{\sqrt{2}}{\text{tr}(t_\alpha^2)} \text{tr} (h^{-1}(\theta) dh(\theta) t_\alpha) = V^\beta \left(A(\theta)_\beta^\alpha - D(\theta)_\beta^\alpha \right) + V^i D(\theta)_i^\alpha \quad (5.36)$$

The above equations are a set of $n = \#roots = \dim H$ differential equations on the parameters θ^α of the h -subgroup element (*compensator*). In the following we will use such set of equations as the basis of an algorithm to produce solutions of the geodesic equations (5.8).

Given these preliminaries, we can establish a new notation. We introduce tangent vectors to the geodesics in the anholonomic basis:

$$\Phi^A = \tilde{V}_I^A(\phi) \dot{\phi}^I \quad (5.37)$$

which are functions only of time: $\Phi^A = \Phi^A(t)$. In this basis the field equations reduce to

$$\dot{\Phi}^A + \Gamma_{BC}^A \Phi^B \Phi^C = 0 \quad (5.38)$$

where now Γ_{BC}^A are the components of the Levi-Civita connection in the chosen anholonomic basis. Explicitly they are related to the components of the Levi Civita connection in an arbitrary holonomic basis by:

$$\Gamma_{BC}^A = \Gamma_{JK}^I V_I^A V_B^J V_C^K - \partial_K(V_J^A) V_B^J V_C^K \quad (5.39)$$

where the inverse vielbein is defined in the usual way:

$$V_I^A V_B^I = \delta_B^A \quad (5.40)$$

The most important point here is that, the connection Γ_{BC}^A can be identified with the *Nomizu connection* defined on a solvable Lie algebra, if the coset representative $\mathbb{L}$ from which we construct the vielbein via eq.(5.25), is solvable, *namely if and only if* the solvability condition (5.29) is satisfied. In fact, as we can read in [103], once we have defined over *Solv* a non degenerate, positive definite and symmetric form:

$$\begin{aligned} \langle , \rangle & : Solv \otimes Solv \longrightarrow \mathbb{R} \\ \langle X, Y \rangle & = \langle Y, X \rangle \end{aligned} \quad (5.41)$$

whose lifting to the manifold produces the metric, the covariant derivative is defined through the **Nomizu operator**:

$$\forall X \in Solv : \mathbb{L}_X : Solv \longrightarrow Solv \quad (5.42)$$

so that

$$\forall X, Y, Z \in Solv : 2\langle Z, \mathbb{L}_X Y \rangle = \langle Z, [X, Y] \rangle - \langle X, [Y, Z] \rangle - \langle Y, [X, Z] \rangle \quad (5.43)$$

while the Riemann curvature 2-form is given by the commutator of two Nomizu operators:

$$R^W_Z(X, Y) = \langle W, \{[\mathbb{L}_X, \mathbb{L}_Y] - \mathbb{L}_{[X, Y]}\} Z \rangle \quad (5.44)$$

This implies that the covariant derivative explicitly reads:

$$\mathbb{L}_X Y = \Gamma_{XY}^Z Z \quad (5.45)$$

where

$$\Gamma_{XY}^Z = \frac{(\langle Z, [X, Y] \rangle - \langle X, [Y, Z] \rangle - \langle Y, [X, Z] \rangle)}{2 \langle Z, Z \rangle} \quad \forall X, Y, Z \in \text{Solv} \quad (5.46)$$

In concrete, the non degenerate, positive definite, symmetric form on the solvable Lie algebra which agrees with equation (5.30) is defined by setting:

$$\begin{aligned} \langle \mathcal{H}_i, \mathcal{H}_j \rangle &= 2 \delta_{ij} \\ \langle \mathcal{H}_i, E_\alpha \rangle &= 0 \\ \langle E_\alpha, E_\beta \rangle &= \delta_{\alpha, \beta} \end{aligned} \quad (5.47)$$

$\forall \mathcal{H}_i, \mathcal{H}_j \in \text{CSA}_{E_{8(8)}}$ and $\forall E_\alpha$, step operator associated to a positive root α of $E_{8(8)}$. Then the Nomizu connection (which is constant) is very easy to calculate. We have:

$$\begin{aligned} \Gamma_{jk}^i &= 0 \\ \Gamma_{\alpha\beta}^i &= \frac{1}{2} (-\langle E_\alpha, [E_\beta, H^i] \rangle - \langle E_\beta, [E_\alpha, H^i] \rangle) = \frac{1}{2} \alpha^i \delta_{\alpha\beta} \\ \Gamma_{ij}^\alpha &= \Gamma_{i\beta}^\alpha = \Gamma_{j\alpha}^i = 0 \\ \Gamma_{\beta i}^\alpha &= \frac{1}{2} (\langle E^\alpha, [E_\beta, H_i] \rangle - \langle E_\beta, [H_i, E^\alpha] \rangle) = -\alpha_i \delta_\beta^\alpha \\ \Gamma_{\alpha\beta}^{\alpha+\beta} &= -\Gamma_{\beta\alpha}^{\alpha+\beta} = \frac{1}{2} N_{\alpha\beta} \\ \Gamma_{\alpha+\beta\beta}^\alpha &= \Gamma_{\beta\alpha+\beta}^\alpha = \frac{1}{2} N_{\alpha\beta} \end{aligned}$$

where $N_{\alpha\beta}$ is defined by the commutator:

$$[E_\alpha, E_\beta] = N_{\alpha\beta} E_{\alpha+\beta} \quad (5.48)$$

which has to be worked out in the algebra.¹ Notice that $\Gamma_{XY}^Z \neq \Gamma_{YX}^Z$ since its expression consists of the first term which is antisymmetric in (X, Y) and the sum of the last two which is symmetric. The component $\Gamma_{\beta i}^\alpha$ consists of the sum of two equal contributions from the antisymmetric and symmetric part, the same contributions cancel in $\Gamma_{i\beta}^\alpha$ which indeed vanishes. By substituting the explicit expression of the Nomizu connection in (5.38) and introducing for the further convenience new names for the tangent vectors along the Cartan generators $\chi^i \equiv \Phi^i$ we have the equations:

$$\begin{aligned} \dot{\chi}^i + \frac{1}{2} \sum_{\alpha \in \Delta_+} \alpha^i \Phi_\alpha^2 &= 0 \\ \dot{\Phi}^\alpha + \sum_{\beta \in \Delta_+} N_{\alpha\beta} \Phi^\beta \Phi^{\alpha+\beta} - \alpha_i \chi^i \Phi^\alpha &= 0 \end{aligned} \quad (5.49)$$

¹The values of the constants $N_{\alpha\beta}$, that enable to construct explicitly the representation of $E_{8(8)}$, used in this paper, are given in the hidden appendix. To see it, download the source file, delete the tag `end{document}` after the bibliography and *LaTeX*.

Eq.s (5.49) encode all the algebraic structure of the $D = 3$ sigma model and due to our oxidation algorithm of the original supergravity in ten dimensions.

All this means that, thanks to the solvability of the algebra (and also to the fact that we know the explicit form of the connection via the Nomizu operator), we have reduced the entire problem of finding time dependent backgrounds for either type IIA or type IIB superstrings or M-theory to the integration of a system of differential equations firmly based on the algebraic structure of $E_{8(8)}$. This is a system of non-linear differential equations, and from this point of view it might seem hopeless to be solved. Yet, due to its underlying algebraic structure, one can use its isometries to generate the complete integral depending on as many integration constants as the number of equations in the system. This is the compensator algorithm we alluded to above, which we shortly outline. To this effect we discuss the role of initial conditions for the tangent vectors to the geodesics. There exist a number of possibilities for such conditions that can truncate the whole system to smaller and simpler ones. The simplest choice is to put all root-vectors to zero in the origin. This will ensure that root-vectors will remain zero at all later times and the system will reduce to

$$\dot{\chi}^i = 0 \quad (5.50)$$

The solution of such a reduced system is trivial and consists of a constant vector $\tilde{V}^A = (\chi^i, 0)$. If we apply an H-rotation to this tangent vector

$$\bar{V}^A = V^B D(\theta)_B{}^A \quad (5.51)$$

we produce a new one, yet, for generic H-rotations we will break the solvable gauge, so that the result no longer produces a solution of eq.s(5.49). However, if we restrict the θ^α parameters of the rotation to satisfy condition (5.36), then the solvable gauge is preserved and the rotated tangent vector $\bar{V}^A$ is still a solution of eq.s(5.49). Hence a general algorithm to solve the differential system (5.49) has been outlined. One starts from the trivial solution in eq.(5.50) and then tries to solve the differential equation for the theta parameter corresponding to one particular H-generator $t_\alpha = E_\alpha - E_{-\alpha}$. Applying this rotation to the trivial solution we obtain a new non trivial one. Then starting from such a new solution we can repeat the procedure and try to solve again the differential equation for the theta parameter relative to a new generator. If we succeed we obtain a further new solution of the original system and we can repeat the procedure a third time for a third generator, iteratively. Indeed, considering eq.(5.36) we see that if $h(t)$ is just a general element of the subgroup H, the system is rather difficult to solve, yet if we choose a rotation around a single axis $h_{\alpha_0} = e^{\theta^{\alpha_0}(t)t_{\alpha_0}}$, then $\frac{1}{\text{Tr}(t_{\alpha_0}^2)}\text{Tr}(h^{-1}dh t_{\alpha_0}) = \dot{\theta}^{\alpha_0}$ and, if all the other equations for $\alpha \neq \alpha_0$ are identically satisfied, as it will turn out to be the case in the examples we consider, then the system reduces to only one first order differential equation on the angle $\theta^{\alpha_0}(t)$.

We name such an algorithm *the compensator method* and we will illustrate it in the next chapter with specific examples.

Chapter 6

The A_2 toy model as a paradigm

In this chapter we consider explicit examples of solutions of the geodesic problem in the case of an A_2 simple algebra. Later we will consider the possible embeddings of such an algebra into the E_8 algebra, so that the solutions we construct here will be promoted to particular solutions of the full $E_{8(8)}/SO(16)$ sigma model. The diverse embeddings will correspond to diverse oxidations of the same three dimensional configuration to $D = 10$ configurations. In other words there exist various non abelian solvable subalgebras $S_5 \subset Solv(E_8/SO(16))$ of dimension 5 which by means of a linear transformation can be identified as the solvable Lie algebra of the simple Lie algebra A_2 , namely the solvable Lie algebra description of the coset manifold:

$$\mathcal{M}_5 \equiv \exp[Solv(A_2)] \cong \frac{SL(3, \mathbb{R})}{SO(3)} \quad (6.1)$$

The detailed study of this model provides our paradigm for the general solution of the complete theory based on the coset manifold $E_{8(8)}/SO(16)$. We emphasize that the possibility of choosing a *normal form* for the initial tangent vector to the geodesic allows to reduce the system of first order equations to a much simpler set, as we started to discuss in the previous chapter in general terms. Such a normal form can be chosen in different ways. In particular it can always be chosen so that it contains only Cartan generators. When this is done the system is always exactly solvable and in terms of pure exponentials. The solution obtained in this way provides a representative for the orbit of geodesics modulo isometries. We can then generate new solutions of the differential system (5.49) by the compensator method we described in the previous chapter. In this chapter we illustrate such an algorithm in the case of the toy A_2 model. The resulting solutions have not only a tuitional interest, rather they provide

examples of solutions of the full $E_{8(8)}$ system and hence of full supergravity. It suffices to embed the A_2 Lie algebra in the full algebra $E_{8(8)}$. We will discuss such embeddings and the corresponding oxidations of our sigma model solutions in later chapters.

6.1 Structure of the A_2 system

Our model consists of 5 scalar fields, which parametrize a coset manifold $\mathcal{M}_5 = \text{SL}(3)/\text{SO}(3)$. Our chosen conventions are as follows. The two simple roots of $\text{SL}(3)$ are:

$$\beta_1 = \left\{ \sqrt{2}, 0 \right\}, \quad \beta_2 = \left\{ -\frac{1}{\sqrt{2}}, \sqrt{\frac{3}{2}} \right\} \quad (6.2)$$

and the third positive root, which is the highest is:

$$\beta_3 = \beta_1 + \beta_2 = \left\{ \frac{1}{\sqrt{2}}, \sqrt{\frac{3}{2}} \right\} \quad (6.3)$$

Furthermore the step operator E_{β_3} is defined through the commutator:

$$E_{\beta_3} = [E_{\beta_1}, E_{\beta_2}] \quad (6.4)$$

and this completely fixes all conventions for the Lie algebra structure constants.

The three generators of the maximally compact subgroup are defined as:

$$t_1 = E_{\beta_1} - E_{-\beta_1}, \quad t_2 = E_{\beta_2} - E_{-\beta_2}, \quad t_3 = E_{\beta_3} - E_{-\beta_3} \quad (6.5)$$

and they satisfy the standard commutation relations:

$$[t_i, t_j] = \epsilon_{ijk} t_k \quad (6.6)$$

In the orthogonal decomposition of the Lie algebra:

$$A_2 = \text{SO}(3) \oplus \mathbb{K}_5 \quad (6.7)$$

the 5-dimensional subspace $\mathbb{K}_5$ is identified with the tangent space to $\mathcal{M}_5$ and corresponds to the $j = 2$ representation of $\text{SO}(3)$

$$[t_\beta, K_A] = Y_{\beta A}^B K_B \quad (6.8)$$

This subspace is spanned by the following generators:

$$\mathbb{K}_5 = \text{Span} \left\{ H_1, H_2, \frac{E_{\beta_1} + E_{-\beta_1}}{\sqrt{2}}, \frac{E_{\beta_2} + E_{-\beta_2}}{\sqrt{2}}, \frac{E_{\beta_3} + E_{-\beta_3}}{\sqrt{2}} \right\} \quad (6.9)$$

Applying to this case the general formulae (5.49) based on the Nomizu connection (5.48) we obtain the differential system:

$$\begin{aligned}
\dot{\chi}_1(t) + \frac{1}{\sqrt{2}}\Phi_1^2(t) - \frac{1}{2\sqrt{2}}\Phi_2^2(t) + \frac{1}{2\sqrt{2}}\Phi_3^2(t) &= 0 \\
\dot{\chi}_2(t) + \frac{\sqrt{3}}{2\sqrt{2}}\Phi_2^2(t) + \frac{\sqrt{3}}{2\sqrt{2}}\Phi_3^2(t) &= 0 \\
\dot{\Phi}_1(t) + \Phi_2(t)\Phi_3(t) - \sqrt{2}\Phi_1(t)\chi_1(t) &= 0 \\
\dot{\Phi}_2(t) - \Phi_1(t)\Phi_3(t) + \frac{1}{\sqrt{2}}\Phi_2(t)\chi_1(t) - \sqrt{\frac{3}{2}}\Phi_2(t)\chi_2(t) &= 0 \\
\dot{\Phi}_3(t) - \frac{1}{\sqrt{2}}\Phi_3(t)\chi_1(t) - \sqrt{\frac{3}{2}}\Phi_3(t)\chi_2(t) &= 0
\end{aligned} \tag{6.10}$$

In order to solve this differential system of equations we recall their geometrical meaning. They are the geodesics equations for the manifold (6.1) written in flat indices, namely in an anholonomic frame. Any geodesics is completely determined by two data: the initial point $p_0 \in M_5$ and the initial tangent vector $\vec{t}_0 \in T(M_5)$ at time $t = 0$. Since our manifold is homogeneous, all points are equivalent and we can just choose the origin of the coset manifold. Since we are interested in determining the orbits of geodesics modulo the action of the isometry group, the relevant question is the following: *in how many irreducible representations* of the tangent group $SO(3)$ does the tangent space decompose? The answer is simple: the 5 dimensional tangent space is irreducible and corresponds to the $j = 2$ representation of $SO(3)$. The next question is: *what is the normal form of such a representation* and how many parameters does it contain. The answer is again simple. A spin two representation is just a symmetric traceless tensor g_{ij} in three dimensions. By means of $SO(3)$ rotations we can reduce it to a diagonal form and the essential parameters are its eigenvalues, namely two parameters, since the third eigenvalue is minus the sum of the other two, being the matrix traceless. So by means of $SO(3)$ rotations a generic 5-dimensional tangent vector can be brought to contain only two parameters. This argument is also evident from the consideration that $5 - 3 = 2$, namely by means of the three $SO(3)$ parameters we can set three components of the 5-dimensional vector to zero.

We can also analyze the normal form of the 5-dimensional representation from the point of view of eigenstates of the angular momentum third component t_3 . This latter has skew eigenvalues $\pm 2, \pm 1$ and 0. The transformation of the matrix $g = \{g_{ij}\}$ under any generator t of the $SO(3)$ Lie algebra is

$$\delta g = [t, g] \tag{6.11}$$

so that the pair of skew eigenstates of the generator t_3 , as given in eq. (6.31), pertaining to the skew eigenvalues ± 2 is provided by the symmetric matrices of the form:

$$g_{(\pm 2)} = \begin{pmatrix} a & 0 & b \\ 0 & 0 & 0 \\ b & 0 & -a \end{pmatrix} \tag{6.12}$$

which can be diagonalized through $SO(3)$ rotations (actually $SO(2)$ in this case) and brought to the normal form:

$$g_2 = \begin{pmatrix} \sqrt{a^2 + b^2} & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -\sqrt{a^2 + b^2} \end{pmatrix} \quad (6.13)$$

which is just one of the two in the pair of skew eigenstates. On the other hand the symmetric traceless matrix that corresponds to the null eigenstate of t_3 is:

$$g_{(0)} = \begin{pmatrix} s & 0 & 0 \\ 0 & -2s & 0 \\ 0 & 0 & s \end{pmatrix} \quad (6.14)$$

A superposition $g_2 + g_{(0)}$ provides the most general diagonal traceless symmetric matrix, namely the *normal form* to which any state in the $j = 2$ irreducible representation can be brought by means of $SO(3)$ rotations.

Alternatively, since the $j = 2$ representation is provided by the tangent space to the $\mathcal{M}_5$ manifold, spanned by the coset generators of $SL(3, \mathbb{R})$ not lying in the compact $SO(3)$ subalgebra, we can identify the normal form of a 5-dimensional vector as one with non vanishing components only in the directions of the Cartan generators. Indeed, by means of $SO(3)$ rotations any vector can be brought to such a form and the counting of independent parameters coincides, namely two. This is a completely general statement for maximally non compact coset manifolds. The rank of the coset is equal to the number of independent parameters in the normal form of the $\mathbb{H}$ representation provided by the coset subspace $\mathbb{K}$.

Relying on these considerations, let us consider the explicit representation of the group $SO(3)$ on the tangent space to our manifold $\mathcal{M}_5$ and how, by means of its transformation we can bring the initial tangent vector to our geodesic to our desired normal form. Indeed our aim is to solve the geodesic equations (6.10) fixing initial conditions:

$$\{\chi_1(0), \chi_2(0), \Phi_1(0), \Phi_2(0), \Phi_3(0)\} = \tilde{V} = \{\tilde{V}_1, \tilde{V}_2, \dots, \tilde{V}_5\} \quad (6.15)$$

where $\tilde{V}$ is the normal form of the 5 vector. To this effect it is convenient to inspect the representative matrices of $SO(3)$ on the tangent space. The three generators of the maximally compact subgroup were defined in (6.5) and in the basis of $\mathbb{K}_5$ provided by the generators (6.9) the 5×5 matrices representing

$SO(3)$ are:

$$\begin{aligned}
 t_1^{[5]} &= \begin{pmatrix} 0 & 0 & -2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 2 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & -1 & 0 \end{pmatrix}, \quad t_2^{[5]} = \begin{pmatrix} 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -\sqrt{3} & 0 \\ 0 & 0 & 0 & 0 & -1 \\ -1 & \sqrt{3} & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \end{pmatrix} \\
 t_3^{[5]} &= \begin{pmatrix} 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & -\sqrt{3} \\ 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 1 & \sqrt{3} & 0 & 0 & 0 \end{pmatrix}
 \end{aligned} \tag{6.16}$$

These matrices have the expected skew eigenvalues:

$$(\pm 2, \pm 1, 0) \tag{6.17}$$

For the generator $t_3^{[5]}$ the corresponding eigenvectors are:

$$\begin{aligned}
 \text{eigenvalue } 0 &\Rightarrow \{-\sqrt{3}, 1, 0, 0, 0\} \\
 \text{eigenvalues } \pm 1 &\Rightarrow \begin{cases} \{0, 0, 1, 1, 0\} \\ \{0, 0, -1, 1, 0\} \end{cases} \\
 \text{eigenvalues } \pm 2 &\Rightarrow \begin{cases} \{\frac{1}{\sqrt{2}}, \sqrt{\frac{3}{2}}, 0, 0, \sqrt{2}\} \\ \{-\frac{1}{\sqrt{2}}, -\sqrt{\frac{3}{2}}, 0, 0, \sqrt{2}\} \end{cases}
 \end{aligned} \tag{6.18}$$

So reduced to normal form the 5-vector of initial condition is a linear combination of the vectors $\vec{g}_{\pm 2} = \{\pm \frac{1}{\sqrt{2}}, \pm \sqrt{\frac{3}{2}}, 0, 0, \sqrt{2}\}$ with the vector $\vec{g}_0 = \{-\sqrt{3}, 1, 0, 0, 0\}$. In particular writing:

$$\begin{aligned}
 \tilde{V}_{\text{normal form}} &= a \vec{g}_0 + b (\vec{g}_{+2} - \vec{g}_{-2}) \\
 &= (-\sqrt{3}a + \sqrt{2}b, a + \sqrt{6}b, 0, 0, 0)
 \end{aligned} \tag{6.19}$$

we obtain an initial tangent vector that has non vanishing components only in the directions of the Cartan generators.¹ For reasons of later convenience we parametrize the initial normal tangent vector as follows:

$$\tilde{V}_{\text{normal form}} = \left(\frac{\omega - \kappa}{4\sqrt{2}}, \frac{3\omega + \kappa}{4\sqrt{6}}, 0, 0, 0 \right) \tag{6.20}$$

and we conclude that we can find a generating solution of the geodesic equations if we solve the first order system for the tangent vectors (eq.s (6.10)) with the initial conditions given by eq.(6.20). With such conditions the differential system

¹Indeed, starting from the Cartan subalgebra, we can generate the whole $\mathbb{K}$ space by applying the adjoint action of the $\mathbb{H}$ subalgebra $Ad_h H_i = h^\alpha [H_i, t_\alpha] = \sqrt{2}\alpha_i h^\alpha K_\alpha$.

(6.10) is immediately solved by:

$$\begin{aligned}\Phi_1^{(gen)}(t) &= 0, & \Phi_2^{(gen)}(t) &= 0, & \Phi_3^{(gen)}(t) &= 0 \\ \chi_1^{(gen)}(t) &= \frac{\omega - \kappa}{4\sqrt{2}}, & \chi_2^{(gen)}(t) &= \frac{3\omega + \kappa}{4\sqrt{6}}\end{aligned}\quad (6.21)$$

From this generating solution we can obtain new ones by performing $\text{SO}(3)$ rotations such that they keep the solvable parametrization of the coset stable. In particular by rotating along the three possible rotation axes we can switch on the root fields $\Phi_\beta(t)$, one by one. This procedure is discussed in chapter 6.3.

6.2 Scalar fields of the A_2 model

In order to find the solutions for the scalar fields ϕ^I , we have to construct explicitly the $\text{SL}(3, \mathbb{R})/\text{SO}(3)$ coset representative $\mathbb{L}$. First, we fix the parametrization of the coset representative as follows

$$\mathbb{L} = \exp[\varphi^3(t) E_3] \exp[\varphi^1(t) E_1 + \varphi^2(t) E_2] \exp[h^1(t) H_1 + h^2(t) H_2] \quad (6.22)$$

Note that here we have ordered the exponentials by height grading, first the highest root of level two, then the simple roots of level one, finally the Cartan generators of level zero. As we will appreciate in eq.s (8.21), this is crucial in order to interpret the scalar fields φ_i as the components of the corresponding p -forms, in oxidation. Choosing the following normalizations for the generators of the fundamental defining representation of the group $\text{SL}(3, \mathbb{R})$:

$$H_1 = \begin{pmatrix} \frac{1}{\sqrt{2}} & 0 & 0 \\ 0 & -\frac{1}{\sqrt{2}} & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad H_2 = \begin{pmatrix} \frac{1}{\sqrt{6}} & 0 & 0 \\ 0 & \frac{1}{\sqrt{6}} & 0 \\ 0 & 0 & -\sqrt{\frac{2}{3}} \end{pmatrix} \quad (6.23)$$

and

$$E^1 = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad E^2 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}, \quad E^3 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad (6.24)$$

we construct a coset representative $\mathbb{L} \in \text{SL}(3, \mathbb{R})/\text{SO}(3)$ explicitly as the following upper triangular matrix:

$$\mathbb{L} = \begin{pmatrix} e^{\frac{h_1(t)}{\sqrt{2}} + \frac{h_2(t)}{\sqrt{6}}} & e^{-\frac{h_1(t)}{\sqrt{2}} + \frac{h_2(t)}{\sqrt{6}}} \varphi_1(t) & e^{-\sqrt{\frac{2}{3}} h_2(t)} (\frac{1}{2} \varphi_1(t) \varphi_2(t) + \varphi_3(t)) \\ 0 & e^{-\frac{h_1(t)}{\sqrt{2}} + \frac{h_2(t)}{\sqrt{6}}} & e^{-\sqrt{\frac{2}{3}} h_2(t)} \varphi_2(t) \\ 0 & 0 & e^{-\sqrt{\frac{2}{3}} h_2(t)} \end{pmatrix} \quad (6.25)$$

Then we calculate the vielbein components through the formula:

$$V^I = \text{Tr} \left[\mathbb{L}^{-1} \frac{d}{dt} \mathbb{L} \mathbf{K}_5^I \right] \quad (6.26)$$

where $\mathbf{K}_5^I$ are the generators of the coset defined in eq.(6.9). The vielbein V^A can be found explicitly as a function of time, recalling that in the solvable gauge it is connected with the solutions of the eq.s (6.10) by the formula $\tilde{V}^i = V^i$, $\tilde{V}^\beta = \Phi^\beta = \sqrt{2}V^\beta$. We obtain the following equations:

$$\begin{aligned} V^1 &= \dot{h}_1(t) &= \chi_1(t) \\ V^2 &= \dot{h}_2(t) &= \chi_2(t) \\ V^3 &= e^{-\sqrt{2}h_1(t)} \frac{1}{\sqrt{2}} \dot{\varphi}_1(t) &= \frac{1}{\sqrt{2}} \Phi_1(t) \\ V^4 &= e^{\frac{h_1(t)-\sqrt{3}h_2(t)}{\sqrt{2}}} \frac{1}{\sqrt{2}} \dot{\varphi}_2(t) &= \frac{1}{\sqrt{2}} \Phi_2(t) \\ V^5 &= e^{-\frac{h_1(t)+\sqrt{3}h_2(t)}{\sqrt{2}}} \frac{(\varphi_2(t)\dot{\varphi}_1(t) - \varphi_1(t)\dot{\varphi}_2(t) + 2\dot{\varphi}_3(t))}{2\sqrt{2}} &= \frac{1}{\sqrt{2}} \Phi_3(t) \end{aligned} \quad (6.27)$$

where in the last column we are supposed to write whatever functions of the time t we have found as solutions of the differential equations (6.10) for the tangent vectors. For future use in the oxidation procedure it is convenient to give a name to the following combination of derivatives:

$$W(t) = \varphi_2(t) \varphi_1'(t) - \varphi_1(t) \varphi_2'(t) + 2\varphi_3'(t) \quad (6.28)$$

and rewrite the last of equations (6.27) as follows:

$$\Phi_3(t) = \frac{1}{2} e^{-\frac{h_1(t)+\sqrt{3}h_2(t)}{\sqrt{2}}} W(t) \quad (6.29)$$

In particular, the generating solution for the tangent vectors (inserting $\chi^1 = \frac{\omega-\kappa}{4\sqrt{2}}$, $\chi^2 = \frac{3\omega+\kappa}{4\sqrt{6}}$, $\Phi^1 = 0$, $\Phi^2 = 0$, $\Phi^3 = 0$) gives, up to irrelevant integration constants, the following scalar fields:

$$\begin{aligned} h_1(t) &= \frac{(\omega - \kappa)t}{4\sqrt{2}}, & h_2(t) &= \frac{(3\omega + \kappa)t}{4\sqrt{6}}, \\ \varphi_1(t) &= 0, & \varphi_2(t) &= 0, & \varphi_3(t) &= 0 \end{aligned} \quad (6.30)$$

6.3 Differential equations for the H-compensators and the generation of new solutions

Non trivial solutions of the system (6.10) can now be obtained from the generating solution (6.21) by means of a suitable H-subgroup compensating transformation, applying to the present case the general procedure of the compensator method outlined at the end of chapter 5. In previous paragraphs we have already collected all the ingredients which are necessary to construct the explicit form of eq.s (5.36). Indeed from eq.s (6.23), (6.24), by recalling the definition (6.5), we immediately obtain the three generators t_i of the compact subgroup SO(3) in the 3-dimensional representation which is also the adjoint:

$$t_1^{[3]} = \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad t_2^{[3]} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix}, \quad t_3^{[3]} = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix} \quad (6.31)$$

On the other hand in eq. (6.1), we constructed the generators t_i in the 5-dimensional $j = 2$ representation, spanned by the vielbein. Hence introducing a compensating group element $h \in \text{SO}(3)$, parametrized by three time dependent angles in the following way:

$$h = \exp [\theta_3(t) t_3] \exp [\theta_2(t) t_2] \exp [\theta_1(t) t_1] \quad (6.32)$$

we immediately obtain the explicit form of the adjoint matrix $A(\theta)$ and of the matrix $D(\theta)$, by setting:

$$\begin{aligned} A(\theta) &= \exp [\theta_3(t) t_3^{[3]}] \exp [\theta_2(t) t_2^{[3]}] \exp [\theta_1(t) t_1^{[3]}] \\ D(\theta) &= \exp [\theta_3(t) t_3^{[5]}] \exp [\theta_2(t) t_2^{[5]}] \exp [\theta_1(t) t_1^{[5]}] \end{aligned} \quad (6.33)$$

Inserting the normal form vector (6.20) and the above defined matrices $A(\theta)$ and $D(\theta)$ into the differential system (5.36) we obtain the following explicit differential equations for the three time dependent θ -parameters:

$$\begin{aligned} \dot{\theta}_3(t) &= \frac{1}{4} \omega \sin 2 \theta_3(t) \\ \dot{\theta}_2(t) &= \frac{1}{8} [\kappa + \omega \cos 2 \theta_3(t)] \sin 2 \theta_2(t) \\ \dot{\theta}_1(t) &= -\frac{1}{16} [\kappa + \kappa \cos 2 \theta_2(t) + \omega [\cos 2 \theta_2(t) - 3] \cos 2 \theta_3(t)] \sin 2 \theta_1(t) + \\ &\quad + \frac{1}{2} \omega \sin^2 \theta_1(t) \sin \theta_2(t) \sin 2 \theta_3(t) \end{aligned} \quad (6.34)$$

At the same time the rotated tangent vector reads as follows in terms of the chosen angles:

$$\begin{aligned} V_{\text{rot}} &\equiv \vec{v}_{\text{n.f.}} D(\theta) \\ V_{\text{rot}}^1 &= \frac{1}{16\sqrt{2}} \{ -\cos 2\theta_1 [2\kappa + 2\kappa \cos 2\theta_2 + \omega \cos 2(\theta_2 - \theta_3)] \\ &\quad - 6\omega \cos 2\theta_3 + \omega \cos 2(\theta_2 + \theta_3) \} + 8\omega \sin 2\theta_1 \sin \theta_2 \sin 2\theta_3 \} \\ V_{\text{rot}}^2 &= \frac{1}{16\sqrt{6}} \{ -2\kappa + 6\kappa \cos 2\theta_2 + 3\omega \cos 2(\theta_2 - \theta_3) \\ &\quad + 6\omega \cos 2\theta_3 + 3\omega \cos 2(\theta_2 + \theta_3) \} \\ V_{\text{rot}}^3 &= \frac{1}{16\sqrt{2}} \{ -[2\kappa + 2\kappa \cos 2\theta_2 + \omega \cos 2(\theta_2 - \theta_3) - 6\omega \cos 2\theta_3 \\ &\quad + \omega \cos 2(\theta_2 + \theta_3)] \sin 2\theta_1 - 8\omega \cos 2\theta_1 \sin \theta_2 \sin 2\theta_3 \} \\ V_{\text{rot}}^4 &= \frac{1}{4\sqrt{2}} \{ \cos \theta_1 (\kappa + \omega \cos 2\theta_3) \sin 2\theta_2 + 2\omega \cos \theta_2 \sin \theta_1 \sin 2\theta_3 \} \\ V_{\text{rot}}^5 &= \frac{1}{4\sqrt{2}} \{ -(\kappa + \omega \cos 2\theta_3) \sin \theta_1 \sin 2\theta_2 \\ &\quad + 2\omega \cos \theta_1 \cos \theta_2 \sin 2\theta_3 \} \end{aligned} \quad (6.35)$$

In this way finding solutions of the original differential system for tangent vectors is reduced to the problem of finding solutions of the differential system for the compensating angles (6.34). The main property of this latter system is that it can be solved iteratively. By inspection we see that the first of eq.s (6.34) is a single differential equation in separable variables for the angle θ_3 .

Inserting the resulting solution into the second of eq.s (6.34) produces a new differential equation in separable variables for θ_2 which can also be solved by direct integration. Inserting these results into the last equation produces instead a non-linear differential equation for θ_1 which is not with separable variables and reads as follows:

$$p_1(t) \sin 2\theta_1(t) + p_2(t) \sin^2 \theta_1(t) + \theta_1'(t) = 0 \quad (6.36)$$

In eq.(6.36) $p_i(t)$ are two functions of time determined by the previous solutions for $\theta_{2,3}(t)$. Explicitly they read:

$$\begin{aligned} p_1(t) &= \frac{1}{32} \{ 2\kappa + 2\kappa \cos 2\theta_2(t) + \omega \cos 2[\theta_2(t) - \theta_3(t)] \\ &\quad - 6\omega \cos 2\theta_3(t) + \omega \cos 2[\theta_2(t) + \theta_3(t)] \} \\ p_2(t) &= -\frac{1}{2}\omega \sin \theta_2(t) \sin 2\theta_3(t) \end{aligned} \quad (6.37)$$

and we can evaluate them using the general solutions of the first two equations in (6.34), namely:

$$\begin{aligned} \theta_3(t) &= -\arcsin \left[\frac{e^{\frac{t\omega}{2}}}{\sqrt{e^{t\omega} + e^{\omega\lambda_3}}} \right] \\ \theta_2(t) &= -\arcsin \frac{e^{\frac{t(\kappa+\omega)+\lambda_2}{4}}}{\sqrt{e^{t\omega} + e^{\omega\lambda_3} + e^{\frac{t(\kappa+\omega)+\lambda_2}{2}}}} \end{aligned} \quad (6.38)$$

where $\lambda_{2,3}$ are two integration constants. Equation (6.36) is actually an integrable differential equation. Indeed multiplying (6.36) by $1/\sin^2 \theta_1$ and introducing the new depending variable $y(t) = \cot \theta_1$, (6.36) becomes actually the following linear differential equation for $y(t)$

$$2y(t)p_1(t) + p_2(t) - y(t)' = 0 \quad (6.39)$$

which can easily be solved. Hence the general integral of (6.36) reads as follows:

$$\theta_1(t) \rightarrow -\operatorname{arccot} \left[e^{2 \int p_1(t) dt} \left(- \int \frac{p_2(t)}{e^{2 \int p_1(t) dt}} dt + \lambda_1 \right) \right] \quad (6.40)$$

where λ_1 is a third integration constant.

In this way the system of eq.s (6.34) has obtained a fully general solution containing three integration constants. By inserting this general solution into equation (6.35) one also obtains a complete general solution of the original differential system for the tangent vectors containing five integration constants $\omega, \kappa, \lambda_1, \lambda_2, \lambda_3$, as many as the first order equations in the system.

Let us consider for instance the choice $\lambda_2 = \lambda_3 = 0$. In this case the solution

(6.38) for the rotation angles $\theta_{3,2}$ reduces to:

$$\begin{aligned}\theta_3(t) &\hookrightarrow -\arcsin \frac{e^{\frac{t\omega}{2}}}{\sqrt{1+e^{t\omega}}} \\ \theta_2(t) &\hookrightarrow -\arcsin \frac{e^{\frac{t(\kappa+\omega)}{4}}}{\sqrt{1+e^{t\omega}+e^{\frac{t(\kappa+\omega)}{2}}}}\end{aligned}\quad (6.41)$$

and by replacing this result into the integrals we get:

$$2 \int p_1(t) dt = \frac{t(\kappa-\omega) + 4 \log(1+e^{t\omega}) - 2 \log(1+e^{t\omega}+e^{\frac{t(\kappa+\omega)}{2}})}{4} \quad (6.42)$$

and

$$-\int \frac{p_2(t)}{e^{2 \int p_1(t) dt}} dt = \frac{1}{1+e^{t\omega}} \quad (6.43)$$

Substituting the above explicit integrations into eq.(6.40) we obtain:

$$\theta_1 \hookrightarrow \operatorname{arccot} \left[\frac{e^{\frac{t(\kappa-\omega)}{4}} (1 + (1+e^{t\omega}) \lambda_1)}{\sqrt{1+e^{t\omega}+e^{\frac{t(\kappa+\omega)}{2}}}} \right] \quad (6.44)$$

that together with eq.s (6.41) provides an explicit solution of equations (6.34). We can replace such a result in eq.(6.35) and obtain the tangent vectors after three rotations. Yet as it is evident from eq.(6.35) the first two rotations are already sufficient to obtain a solution where all the entries of the 5-dimensional tangent vector are non vanishing and hence all the root fields are excited. In the sequel we will consider the two solutions obtained by means of the first rotation and by means of the first plus the second. They will constitute our paradigm of how the full system can be eventually solved. These solutions however, as we discuss in later chapters, are not only interesting as toy models and examples. Indeed through oxidation they can be promoted to very interesting backgrounds of ten dimensional supergravity that make contact with the physics of S -branes.

6.3.1 Solution of the differential equations for the tangent vectors with two Cartan and one nilpotent field

Let us consider the system (6.34) and put

$$\theta_1 = \theta_2 = \text{const} = 0 \quad (6.45)$$

This identically solves the last two equations and we are left with the first whose general integral was already given in eq.(6.38). By choosing the integration constant $\lambda_3 = 0$ we can also write:

$$\theta_3(t) = \arccos \frac{1}{\sqrt{1+e^{t\omega}}} \quad (6.46)$$

By inserting (6.46) and (6.45) into (6.35) we obtain the desired solution for the tangent vectors:

$$\begin{aligned}\chi_1(t) &= -\frac{\kappa + \omega \tanh \frac{t\omega}{2}}{4\sqrt{2}}, & \chi_2(t) &= \frac{\kappa - 3\omega \tanh \frac{t\omega}{2}}{4\sqrt{6}}, \\ \Phi_1(t) &= 0, & \Phi_2(t) &= 0, & \Phi_3(t) &= \frac{\omega}{\sqrt{(1+e^{-t\omega})(1+e^{t\omega})}}\end{aligned}\quad (6.47)$$

where one root field is excited.

Next we address the problem of solving the equations for the scalar fields, namely eq.s(6.27), which are immediately integrated, obtaining:

$$\begin{aligned}h_1(t) &= -\frac{t\kappa + 2\log(\cosh \frac{t\omega}{2})}{4\sqrt{2}}, & h_2(t) &= \frac{t\kappa - 6\log(\cosh \frac{t\omega}{2})}{4\sqrt{6}}, \\ \varphi_1(t) &= 0, & \varphi_2(t) &= 0, & \varphi_3(t) &= \frac{\sqrt{1 + \tanh(\frac{t\omega}{2})}}{e^{\frac{t\omega}{2}}\sqrt{1 + e^{t\omega}}}\end{aligned}\quad (6.48)$$

We can now insert eq.s (6.48) into the form of the coset representative (6.25) and we obtain the geodesic as a map of the time line into the solvable group manifold and hence into the coset manifold depending on your taste for interpretation:

$$\mathbb{R}_t \hookrightarrow \exp[Solv(A_2)] \simeq \frac{SL(3, \mathbb{R})}{SO(3)} \quad (6.49)$$

In chapter (8.2.1) the oxidation of this sigma-model solution to a full fledged supergravity background in $D = 10$ is studied.

6.3.2 Solution of the differential equations for the tangent vectors with two Cartan and three nilpotent fields

Then we continue the hierarchical solution of the (6.34) differential system by considering the next rotation θ_2 . We set $\theta_1 = \text{const} = 0$ and we replace in eq.s (6.34) the solution (6.46) for θ_3 , with $\lambda_3 = 0$. The first and the last differential equations are identically satisfied. The second equation was already solved in eq.(6.38). By choice of the irrelevant integration variable $\lambda_2 = 0$ a convenient solution of the above equation is provided by the following time dependent angle:

$$\theta_2(t) = -\arcsin \frac{e^{\frac{t(\kappa+\omega)}{4}}}{\sqrt{1 + e^{t\omega} + e^{\frac{t(\kappa+\omega)}{2}}}} \quad (6.50)$$

Inserting (6.46) and (6.50) into (6.35) we obtain :

$$\begin{aligned}
\chi_1(t) &= \frac{-(1+e^{t\omega})^2 \kappa - (-1+e^{t\omega}) \left(1+e^{t\omega} + 2e^{\frac{t(\kappa+\omega)}{2}}\right) \omega}{4\sqrt{2} (1+e^{t\omega}) \left(1+e^{t\omega} + e^{\frac{t(\kappa+\omega)}{2}}\right)} \\
\chi_2(t) &= \frac{\left(1+e^{t\omega} - 2e^{\frac{t(\kappa+\omega)}{2}}\right) \kappa - 3(-1+e^{t\omega}) \omega}{4\sqrt{6} \left(1+e^{t\omega} + e^{\frac{t(\kappa+\omega)}{2}}\right)} \\
\Phi_1(t) &= -\frac{e^{\frac{t(\kappa+3\omega)}{4}} \omega}{(1+e^{t\omega}) \sqrt{1+e^{t\omega} + e^{\frac{t(\kappa+\omega)}{2}}}} \\
\Phi_2(t) &= \frac{e^{\frac{t(\kappa+\omega)}{4}} (\kappa + e^{t\omega} (\kappa - \omega) + \omega)}{2\sqrt{1+e^{t\omega}} \left(1+e^{t\omega} + e^{\frac{t(\kappa+\omega)}{2}}\right)} \\
\Phi_3(t) &= \frac{e^{\frac{t\omega}{2}} \omega}{\sqrt{(1+e^{t\omega}) \left(1+e^{t\omega} + e^{\frac{t(\kappa+\omega)}{2}}\right)}} \tag{6.51}
\end{aligned}$$

Integrating eq.s(6.27) with this new choice of the left hand side we obtain:

$$\begin{aligned}
h_1(t) &= \frac{t(-\kappa + \omega) - 4 \log(1+e^{t\omega}) + 2 \log(1+e^{t\omega} + e^{\frac{t(\kappa+\omega)}{2}})}{4\sqrt{2}} \\
h_2(t) &= \frac{t(\kappa + 3\omega) - 6 \log(1+e^{t\omega} + e^{\frac{t(\kappa+\omega)}{2}})}{4\sqrt{6}} \\
\varphi_1(t) &= \frac{1}{1+e^{t\omega}} \\
\varphi_2(t) &= -\frac{(1+e^{t\omega})}{1+e^{t\omega} + e^{\frac{t(\kappa+\omega)}{2}}} \\
W(t) &= \frac{2e^{t\omega} \omega}{(1+e^{t\omega}) \left(1+e^{t\omega} + e^{\frac{t(\kappa+\omega)}{2}}\right)} \tag{6.52}
\end{aligned}$$

In chapter 8.2.2 we will see how this σ -model solution can be oxidized, among other choices, to an interesting $S3/S1$ -brane solution of type IIB supergravity.

Chapter 7

The E_8 Lie algebra: Reduction, Oxidation and subalgebra embeddings

We come now to a close examination of the E_8 Lie algebra and we show how the hierarchical dimensional *reduction/oxidation* [104]–[105] of supergravity backgrounds is algebraically encoded in the hierarchical embedding of subalgebras into the E_8 algebra. Similarly the structure of the bosonic lagrangians of type II A/B supergravities in $D = 10$ [106, 107, 108, 109] is encoded in the decomposition of the solvable Lie algebra $Solv(E_{8(8)}/SO(16))$ according to irreducible representations of two inequivalent subgroups $GL(7, \mathbb{R})_{A/B} \subset E_{8(8)}$, respectively associated with the moduli space of flat metrics on a torus T^7 in compactified type II A or type II B theory [47, 48, 110].

In order to carry out our programme we begin by spelling out the E_8 Lie algebra in our chosen conventions.

Using the Cartan–Weyl basis the Lie algebra can be written in the standard form:

$$\begin{aligned} [\mathcal{H}_i, \mathcal{H}_j] &= 0 \\ [\mathcal{H}_i, E_\alpha] &= \alpha_i E_\alpha & \forall \alpha \in \Delta_+ \\ [\mathcal{H}_i, E_{-\alpha}] &= -\alpha_i E_{-\alpha} \\ [E_\alpha, E_\beta] &= N_{\alpha\beta} E_{\alpha+\beta} & \text{if } \alpha + \beta \in \Delta_+ \\ [E_\alpha, E_\beta] &= 0 & \text{if } \alpha + \beta \notin \Delta_+ \\ [E_\alpha, E_{-\beta}] &= \delta_{\alpha\beta} \alpha^i H_i \end{aligned} \tag{7.1}$$

where $\mathcal{H}_i$ are the 8 Cartan generators, E_α are the 120 step operators associated

with the positive roots $\alpha \in \Delta_+$.¹

Our choice of the simple roots as vectors in an Euclidean $\mathbb{R}^8$ space is the following one

$$\begin{aligned}
 \alpha_1 &= \{0, 1, -1, 0, 0, 0, 0, 0\} \\
 \alpha_2 &= \{0, 0, 1, -1, 0, 0, 0, 0\} \\
 \alpha_3 &= \{0, 0, 0, 1, -1, 0, 0, 0\} \\
 \alpha_4 &= \{0, 0, 0, 0, 1, -1, 0, 0\} \\
 \alpha_5 &= \{0, 0, 0, 0, 0, 1, -1, 0\} \\
 \alpha_6 &= \{0, 0, 0, 0, 0, 1, 1, 0\} \\
 \alpha_7 &= \{-\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\} \\
 \alpha_8 &= \{1, -1, 0, 0, 0, 0, 0, 0\}
 \end{aligned}$$

The Dynkin diagrams corresponding to $GL(7, \mathbb{R})_{A/B}$ are defined by the following simple roots:

$$\begin{aligned}
 GL(7, \mathbb{R})_A &\leftrightarrow \{\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_6, \alpha_8\} \\
 GL(7, \mathbb{R})_B &\leftrightarrow \{\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_8\}
 \end{aligned} \tag{7.2}$$

These two choices are illustrated in fig.7.1 where the roots belonging to the $SL(7, \mathbb{R}) \subset GL(7, \mathbb{R})$ subgroup of the metric group are painted white. The roots eventually corresponding to a B-field are instead painted black, while the root eventually corresponding to a RR state are painted gray. As one sees the difference between the A and B interpretation of the same Dynkin diagram, named by us a *painting* of the same, resides in the fact that in the first case the RR root is linked to a metric, while in the second it is linked to a B-field.

In order to motivate the above identifications, let us start recalling that the T^7 metric-moduli parametrize the coset $\mathcal{M}_g^{A/B} = GL(7, \mathbb{R})_{A/B}/SO(7)$ in the type IIA or B frameworks. If we describe $\mathcal{M}_g^{A/B}$ as a solvable Lie group generated by the solvable Lie algebra $Solv(\mathcal{M}_g^{A/B})$ [110, 35] then its coset representative $L^p_{\hat{q}}$ (in our notation the hatted indices are rigid, i.e. are acted on by the compact isotropy group) will be a solvable group element which, in virtue of the Iwasawa decomposition can be expressed as the product of a matrix $\mathcal{N}^{-1T}$, which is the exponent of a nilpotent matrix, times a diagonal one $\mathcal{H}^{-1}$: $L = \mathcal{N}^{-1T} \mathcal{H}^{-1}$. Indeed the matrix $\mathcal{N}^{-1T}$ is the exponential of the sub-algebra $\mathcal{A}$ of $Solv(\mathcal{M}_g^{A/B})$ spanned by the shift operators corresponding to the $GL(7, \mathbb{R})_{A/B}$ positive roots, while $\mathcal{H}^{-1}$ is the exponential of the six-dimensional

¹The values of the constants $N_{\alpha\beta}$, that enable to construct explicitly the representation of $E_{8(8)}$, used in this paper, are given in the hidden appendix. To see it, download the source file, delete the tag `end{document}` after the bibliography and *LaTeX*.




Figure 7.1: There are two different ways of embedding the $SL(7, \mathbb{R})$ Lie algebra in $E_{8(8)}$ which correspond to the type IIA and type IIB interpretation of the same sigma model. This can also be seen as a different way of painting the same Dynkin diagram with blobs that are either associated with the metric (white) or with the B -field (gray) or with the Ramond-Ramond field (black). Furthermore the T-duality transforming the A painting into the B one is just the change in sign of the ϵ_6 vector in Euclidean space. Indeed this corresponds, physically to inverting one of the torus radii $R_9 \rightarrow \alpha'/R_9$.

$GL(7, \mathbb{R})_{A/B}$ Cartan subalgebra. The vielbein $E_p^{\hat{q}}$ corresponding to the T^7 metric g_{pq} will have the following expression :

$$\begin{aligned} E &= L^{-1T} = \mathcal{N} \mathcal{H}, \\ g &= E E^T = \mathcal{N} \mathcal{H} \mathcal{H}^T \mathcal{N}^T. \end{aligned} \quad (7.3)$$

The matrix $\mathcal{N}$ is non-trivial only if T^7 has off-diagonal metric-moduli. In the case of a straight torus, namely when $g_{pq} = e^{2\sigma_p} \delta_{pq} = R_p^2 \delta_{pq}$ the diagonal entries of $\mathcal{H}$ are just the radii R_p : $\mathcal{H}_p^{\hat{q}} = R_p \delta_p^{\hat{q}}$.

The decomposition of $Solv(E_{8(8)}/SO(16)) = Solv_8$ with respect to $Solv(GL(7, \mathbb{R})/SO(7)) = Solv_7^{A/B}$ has the following form:

$$Solv_8 = Solv_7^{A/B} + o(1, 1) + \mathcal{A}^{[1]} + \mathbf{B}^{[1]} + \mathbf{B}^{[2]} + \sum_k \mathbf{C}_{A/B}^{[k]} \quad (7.4)$$

where $o(1,1)$ denotes the Cartan generator $H_{\alpha[7]}$ parametrized by the ten dimensional dilaton ϕ , $\mathbf{B}^{[2]} = B_{2+p,2+q}$ $\mathbf{B}^{p,q}$ is the subalgebra parametrized by the internal components of the Kalb–Ramond field and

$\mathbf{C}_{A/B}^{[k]} = C_{2+p_1 \dots 2+p_k}$ $\mathbf{C}^{p_1 \dots p_k}$ the subalgebra spanned by the internal components of the R–R k -form (in our conventions $C_{2+p_1, \dots, 2+p_k}$ for $k > 4$ are the *dualized* vectors $C_{2+q_1 \dots 2+q_{7-k}, \mu}$, with $\epsilon^{p_1 \dots p_k q_1 \dots q_{7-k}} \neq 0$). Finally the spaces $\mathcal{A}^{[1]}$ and $\mathbf{B}^{[1]}$ are parametrized by the dualized Kaluza–Klein and Kalb–Ramon vectors: $g_{p\mu}$, $B_{2+p, \mu}$. It can be verified that the shift generators corresponding to $E_{8(8)}$ positive roots decompose into order- k antisymmetric tensorial representations $\mathbf{T}^{[k]} = \{\mathbf{T}^{p_1 \dots p_k}\}$ with respect to the adjoint action of $GL(7, \mathbb{R})_{A/B}$:

$$\mathbf{E} \in GL(7, \mathbb{R})_{A/B} : \mathbf{E} \cdot \mathbf{T}^{p_1 \dots p_k} \cdot \mathbf{E}^{-1} = E^{p_1}_{q_1} \dots E^{p_k}_{q_k} \mathbf{T}^{q_1 \dots q_k} \quad (7.5)$$

From the definitions (7.2) we see that the shift generators corresponding to positive roots decompose with respect to $GL(7, \mathbb{R})_A$ into the subspaces $\mathbf{B}^{[2]}$ and $\mathbf{C}_A^{[k]}$, $k = 1, 3, 5, 7$ and with respect to $GL(7, \mathbb{R})_B$ into $\mathbf{B}^{[2]}$ and $\mathbf{C}_B^{[k]}$, $k = 0, 2, 4, 6$.

As far as the R–R scalars are concerned, these representations correspond indeed to the tensorial structure of the type IIA spectrum

$$C_{2+p}, C_{2+p, 2+q, 2+r}, C_{\mu}, C_{2+p, 2+q, \mu}$$

and type IIB spectrum

$$C, C_{2+p, 2+q}, C_{2+p, 2+q, 2+r, 2+s}, C_{2+p, \mu}.$$

We can now define a one-to-one correspondence between axions and $E_{8(8)}$ positive roots. The T^7 moduli space is $SO(7, 7)_T/[SO(7) \times SO(7)] = \text{Exp}(\text{Solv}_T)$ parametrized by the scalars $g_{2+p, 2+q}$ and $B_{2+p, 2+q}$, where:

$$\text{Solv}_T = \text{Solv}_7^{A/B} + \mathbf{B}^{[2]} \quad (7.6)$$

In three dimensions the scalar fields deriving from the dualization of $g_{p\mu}$ and $B_{p\mu}$ together with the dilaton ϕ enlarge the manifold $SO(7, 7)_T/[SO(7) \times SO(7)]$ to $SO(8, 8)/[SO(8) \times SO(8)] = \text{Exp}(\text{Solv}_{NS})$ where now:

$$\text{Solv}_{NS} = \text{Solv}_7^{A/B} + o(1, 1) + \mathcal{A}^{[1]} + \mathbf{B}^{[1]} + \mathbf{B}^{[2]}, \quad (7.7)$$

This manifold is parametrized by the 64 NS scalar fields. If we decompose Solv_8 with respect to Solv_{NS} we may achieve an intrinsic group-theoretical characterization of the NS and R–R scalars. From this point of view the R–R scalar fields span the 64-dimensional subalgebra $\text{Solv}_8/\text{Solv}_{NS}$ which coincides with a spinorial representation of $SO(7, 7)_T$ with a definite chirality. Therefore the corresponding positive roots have grading one with respect to the $SO(7, 7)_T$ spinorial root $\alpha[7]$. Finally the higher-dimensional origin of the three dimensional scalar fields can be determined by decomposing Solv_8 with respect to the

solvable algebra $Solv_{11-D}$ generating the scalar manifold $E_{11-D(11-D)}/H$ of the D -dimensional maximal supergravity. This decomposition is defined by the embedding of the higher-dimensional duality groups $E_{11-D(11-D)}$ inside the three dimensional one. The Dynkin diagrams of the $E_{11-D(11-D)}$ nested Lie algebras are arranged according to the the pictures displayed in Fig. 7.2 and Fig. 7.3.

Let us now comment on the geometrical relation between the Type IIA and IIB representations. The two $SL(7, \mathbb{R})_{A/B}$ Dynkin diagrams are mapped into each other by the $SO(7, 7)$ outer automorphism $\epsilon_7 \rightarrow -\epsilon_7$ which corresponds, in the light of our parametrization of the $E_{8(8)}$ Cartan generators, to a T-duality along the direction x^9 . To show that this operation is indeed a T-duality (see [111] and also [112] for a geometrical definition of T-duality in the solvable Lie algebra formalism) let us recall the parametrization of the Cartan subalgebra in our setup:

Type IIB:

$$\vec{h} \cdot \vec{H} = \sum_{p=1}^7 \sigma_p (\epsilon_p - \epsilon_8) - \frac{\phi}{2} \alpha_7 = \sum_{p=1}^7 \tilde{\sigma}_p (\epsilon_p - \epsilon_8) + 2\phi \epsilon_8 \quad (7.8a)$$

Type IIA:

$$\vec{h} \cdot \vec{H} = \sum_{p=1}^7 \sigma_p (\epsilon'_p - \epsilon'_8) - \frac{\phi}{2} (\alpha_7 + \epsilon_7) = \sum_{p=1}^7 \tilde{\sigma}_p (\epsilon'_p - \epsilon'_8) + 2\phi \epsilon'_8 \quad (7.8b)$$

$$\text{with } \epsilon'_w = \begin{cases} \epsilon_w & \text{if } w \neq 7 \\ \epsilon_w & \text{if } w = 7 \end{cases}$$

where, in the case of a compactification on a straight torus, $\sigma_p = \log(R_p)$ and $\tilde{\sigma}_p = \log(\tilde{R}_p)$, R_p and $\tilde{R}_p$ being the T^7 radii in the ten-dimensional Einstein- or string-frame respectively. Let us consider a T-duality along directions $x^{i_1}, \dots, x^{i_k}$: $\tilde{R}_{i_r} \rightarrow 1/\tilde{R}_{i_r}$ ($r = 1, \dots, k$, $\alpha' = 1$). The transformation $\epsilon_{i_r} \rightarrow -\epsilon_{i_r}$ in the expression of $\vec{h} \cdot \vec{H}$ can be absorbed by the transformation $\tilde{\sigma}_{i_r} \rightarrow -\tilde{\sigma}_{i_r}$ and $\phi \rightarrow \phi - \sum_{r=1}^k \tilde{\sigma}_{i_r}$ which is indeed the effect of the T-duality.

As a result of this analysis the precise one-to-one correspondence between axions and positive roots can now be given in the following form:

Type IIB:

$$\begin{aligned}
C_{2+p_1 \dots 2+p_k} &\leftrightarrow \alpha_7 + \epsilon_{p_1} + \dots \epsilon_{p_k}, \quad (k = 2, 4), \\
C_{2+p, \mu} &\leftrightarrow \alpha_7 + \epsilon_{q_1} + \dots \epsilon_{q_6}, \quad (\epsilon^{pq_1 \dots q_6} \neq 0), \\
B_{2+p, 2+q} &\leftrightarrow \epsilon_p + \epsilon_q, \\
B_{2+p, \mu} &\leftrightarrow -\epsilon_p - \epsilon_8, \\
\gamma_{2+p}^{2+q} &\leftrightarrow \epsilon_p - \epsilon_q \\
\gamma_\mu^{2+q} &\leftrightarrow \epsilon_q - \epsilon_8
\end{aligned}$$

Type IIA:

$$\begin{aligned}
C_{2+p_1 \dots 2+p_k} &\leftrightarrow \alpha_7 + \epsilon_7 + \epsilon'_{p_1} + \dots \epsilon'_{p_k}, \quad (k = 1, 3), \\
C_{2+p_1, 2+p_2, \mu} &\leftrightarrow \alpha_7 + \epsilon_7 + \epsilon'_{q_1} + \dots \epsilon'_{q_5}, \quad (\epsilon^{p_1 p_2 q_1 \dots q_5} \neq 0), \\
C_\mu &\leftrightarrow \alpha_7 + \epsilon_7 + \epsilon'_1 + \dots \epsilon'_7, \\
B_{2+p, 2+q} &\leftrightarrow \epsilon'_p + \epsilon'_q, \\
B_{2+p, \mu} &\leftrightarrow -\epsilon'_p - \epsilon'_8, \\
\gamma_{2+p}^{2+q} &\leftrightarrow \epsilon'_p - \epsilon'_q \\
\gamma_\mu^{2+q} &\leftrightarrow \epsilon'_q - \epsilon'_8
\end{aligned}$$

(7.9)

where γ_{2+p}^{2+q} are the parameters entering the matrix $\mathcal{N}$ and which determine the off-diagonal entries of the T^7 vielbein $E_p^{\hat{q}}$:

$$\mathcal{N} \equiv \exp(\gamma_{2+p}^{2+q} \mathcal{A}_q^p) \quad (7.10)$$

for a precise definition of the above exponential representation see [110]. The fields γ_μ^q denote the scalars dual to the Kaluza–Klein vectors.




Figure 7.2: The Dynkin diagrams of $E_{3(3)} \subset E_{4(4)} \subset E_{5(5)}$ and the labeling of simple roots




Figure 7.3: The Dynkin diagrams of $E_{6(6)} \subset E_{7(7)} \subset E_{8(8)}$ and the labeling of simple roots

Chapter 8

Oxidation of the A_2 solutions

In this chapter, as a working illustration of the oxidation process we derive two full fledged $D = 10$ supergravity backgrounds corresponding to the two A_2 sigma model solutions derived in previous chapters. As we already emphasized in our introduction the correspondence is not one-to-one, rather it is one-to-many. This has two reasons. First of all we can either oxidize to a type IIA or to a type IIB configuration. Secondly, even within the same supergravity choice (A or B), there are several different oxidations of the same abstract sigma model solution, just as many as the different ways of embedding the solvable $Solv(A_2)$ algebra into the solvable $Solv(E_8/SO(16))$ algebra. These embeddings lead to quite different physical interpretations of the same abstract sigma model solution.

Our first task is the classification of these inequivalent embeddings.

8.1 Possible embeddings of the A_2 algebra

In order to study the possible embeddings it is convenient to rely on a compact notation and on the following graded structure of the Solvable Lie algebra

$Solv(E_8/SO(16))$ characterized by the following non vanishing commutators:

$$[\mathcal{A}, \mathcal{A}] = \mathcal{A} \quad (8.1a)$$

$$[\mathcal{A}, \mathcal{A}^{[1]}] = \mathcal{A}^{[1]} \quad (8.1b)$$

$$[\mathbf{B}^{[2]}, \mathbf{B}^{[1]}] = \mathcal{A}^{[1]} \quad (8.1c)$$

$$[\mathcal{A}, \mathbf{B}^{[2]}] = \mathbf{B}^{[2]} \quad (8.1d)$$

$$[\mathcal{A}, \mathbf{C}^{[k]}] = \mathbf{C}^{[k]} \quad (8.1e)$$

$$[\mathbf{B}^{[2]}, \mathbf{C}^{[k]}] = \mathbf{C}^{[k+2]} \quad (8.1f)$$

$$[\mathbf{C}^{[k]}, \mathbf{C}^{[6-k]}] = \mathbf{B}^{[1]} \quad (8.1g)$$

$$[\mathbf{C}^{[4]}, \mathbf{C}^{[4]}] = \mathbf{A}^{[1]} \quad (8.1h)$$

In eq. (8.1) $\mathcal{A}^{[1]}$, $\mathbf{B}^{[2]}$, $\mathbf{B}^{[1]}$ and $\mathbf{C}^{[k]}$ are the spaces of nilpotent generators defined in the previous chapter. While $\mathcal{A}$ is the $Solv_7^{A/B}$ Lie algebra. In view of the above graded structure there are essentially 8 physically different ways of embedding the A_2 algebra into $E_{8(8)}$.

- 1 Every root $\beta_{1,2,3}$ is a metric generator $\mathcal{A}$. In this case the A_2 Lie algebra is embedded into the $SL(7, \mathbb{R})$ subalgebra of $E_{8(8)}$ and the corresponding oxidation leads to a purely gravitational background of supergravity which is identical in the type IIA or type IIB theory.
- 2 The two simple roots $\beta_{1,2}$ are respectively associated with a metric generator $\mathcal{A}$ and a B -field generator $\mathbf{B}$. The composite root β_3 is associated with a second B -field generator. This is so because the $\mathbf{B}$ -generators span an antisymmetric rank 2 representation of $SL(7, \mathbb{R})$. In this case oxidation leads to a purely NS configuration, shared by type II A and type II B theory, involving the metric, the dilaton and the B -field alone.
- 3 The two simple roots $\beta_{1,2}$ are respectively associated with a metric generator $\mathcal{A}$ and with a RR k -form generator $\mathbf{C}^{[k]}$. The composite root β_3 is associated with a second RR generator $\mathbf{C}^{[k]}$ pertaining to the same k . This follows again from the fact that the $\mathbf{C}^{[k]}$ generators span an $SL(7, \mathbb{R})$ representation. In this case oxidation leads to different results in type II A and type II B theories, although the metric is the same for the two cases and it has non trivial off-diagonal parts.
- 4 The simple roots $\beta_{1,2}$ are respectively associated with a RR k -form generator $\mathbf{C}^{[k]}$ and with a B -field generator $\mathbf{B}$. The composite root β_3 is associated with a $k+2$ form generator $\mathbf{C}^{k+2}$. In this case the metric is purely diagonal and we have non trivial B -fields and RR forms. Type IIA and type IIB

oxidations are just different in this sector. The NS sector is the same for both.

- 5 The two simple roots $\beta_{1,2}$ are respectively associated with a RR generator $\mathbf{C}^{[k]}$ and a RR generator $\mathbf{C}^{[6-k]}$. The composite root β_3 is associated with a $\mathbf{B}^{[1]}$ generator. The oxidation properties of this case are just similar to those of the previous case. Also here the metric is diagonal.
- 6 The root β_1 corresponds to an off diagonal element of the internal metric, namely belongs to $\mathcal{A}$, while $\beta_{2,3}$ correspond to scalars dual to the Kaluza–Klein vectors $g_{2+p,\mu}$ namely belong to $\mathcal{A}^{[1]}$.
- 7 The root $\beta_1 \in \mathbf{B}^{[2]}$, namely it describes an internal component of the B–field. The root $\beta_2 \in \mathbf{B}^{[1]}$ namely it corresponds to a B–field with mixed indices. The root $\beta_3 \in \mathcal{A}^{[1]}$ is associated with a mixed component of the metric.
- 8 In type IIB theory the roots $\beta_{1,2}$ belong to $C^{[4]}$ namely are associated with two different components of the internal 4–form, while $\beta_3 \in \mathcal{A}^{[1]}$ describes a mixed component of the metric.

8.2 Choice of one embedding example

As an illustration, out of the above list we choose one example of embedding that has an immediate and nice physical interpretation in terms of a brane system. We consider the case 4, with a RR generator $\mathbf{C}^{[2]}$ and a B–field generator respectively associated with $\beta_{1,2}$ and a $\mathbf{C}^{[4]}$ generator associated with the composite root β_3 . In particular we set:

$$\begin{aligned}\beta_1 &\rightarrow \mathbf{B}^{34} \\ \beta_2 &\rightarrow \mathbf{C}^{89} \\ \beta_3 &\rightarrow \mathbf{C}^{3489} \sim \mathbf{C}^{\mu 567}\end{aligned}\tag{8.2}$$

More precisely this corresponds to identifying $\beta_{1,2,3}$ with the following roots of $E_{8(8)}$ according to their classification given in the appendix:

$$\begin{aligned}\beta_1 &\hookrightarrow \alpha[69] = \epsilon_1 + \epsilon_2 && \leftrightarrow B_{34} \\ \beta_2 &\hookrightarrow \alpha[15] = \alpha[7] + \epsilon_6 + \epsilon_7 && \leftrightarrow C_{89} \\ \beta_3 &\hookrightarrow \alpha[80] = \alpha[7] + \epsilon_1 + \epsilon_1 + \epsilon_6 + \epsilon_7 && \leftrightarrow C_{3489} \sim C_{\mu 567}\end{aligned}\tag{8.3}$$

where $\alpha[7] = \{-\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$ is the spinorial simple root of $E_{8(8)}$.

Next given the explicit form of the two roots $\beta_1 \hookrightarrow \alpha[69]$ and $\beta_2 \hookrightarrow \alpha[15]$ we construct the 2–dimensional subspace of the Cartan subalgebra which is orthogonal to the orthogonal complement of $\alpha[69]$ and $\alpha[15]$ in $\mathbb{R}^8$. We immediately

see that this subspace is spanned by all 8 vectors of the form:

$$\vec{h} = \{x, x, y, y, y, -y, -y, y\} \quad (8.4)$$

so that we find:

$$\vec{h} \cdot \vec{\alpha}[69] = 2x \quad ; \quad \vec{h} \cdot \vec{\alpha}[15] = -(x + 3y) \quad (8.5)$$

Then we relate the fields x and y to the diagonal part of the ten dimensional metric.

To this effect we start from the general relations between the ten-dimensional metric in the *Einstein frame* and the fields in three-dimensions evaluated in the $D = 3$ Einstein frame, then we specialize such relations to our particular case.

General relations in dimensional reduction The Einstein frame metric in $D = 10$ can be written as:

$$G_{MN}^{(Einstein)} = \left(\frac{\exp[4\phi_3 - \frac{1}{2}\phi] g_{\mu\nu}^{(E,3)} + G_{ij} \mathcal{A}_\mu^i \mathcal{A}_\nu^j}{G_{jk} \mathcal{A}_\nu^k} \middle| \frac{G_{ik} \mathcal{A}_\mu^k}{G_{ij}} \right) \quad (8.6)$$

where $g_{\mu\nu}^{(E,3)}$ is the three dimensional Einstein frame metric (5.10) determined by the solution of the $D = 3$ -sigma model via equations (5.17) and (5.18,5.19). On the other hand G_{ij} is the Einstein frame metric in the internal seven directions. It parametrizes the coset:

$$\frac{\text{GL}(7, \mathbb{R})}{\text{SO}(7)} = O(1, 1) \times \frac{\text{SL}(7, \mathbb{R})}{\text{SO}(7)} \quad (8.7)$$

In full generality, recalling eq.s(7.3) we can set ([110, 35]):

$$G = E E^T \quad ; \quad E = \mathcal{N} \mathcal{H} \quad (8.8)$$

where, in this case:

$$\mathcal{N}_{ij} = \delta_{ij} \quad (8.9)$$

since there are no roots associated with metric generators, while the diagonal matrix:

$$H_{ij} = \exp[\sigma_i] \delta_{ij} \quad (8.10)$$

parametrizes the degrees of freedom associated with the Cartan subalgebra of $O(1, 1) \times \text{SL}(7, \mathbb{R})$. The relation of the fields σ_i with the dilaton field and the Cartan fields of $E_{8(8)}$ is obtained through the following general formulae:

$$\begin{aligned} \vec{h} &= \sum_{p=1}^7 (\sigma_p + \frac{1}{4}\phi) \vec{\epsilon}_p + 2\phi_3 \vec{\epsilon}_8 \\ \phi_3 &= \frac{1}{8}\phi - \frac{1}{2} \sum_{p=1}^7 \sigma_p \end{aligned} \quad (8.11)$$

ϕ being the dilaton in $D = 10$ and ϕ_3 its counterpart in $D = 3$. The above formula follows immediately from eq.(7.8)

We also stress the following general property of the parametrization (8.6) for the $D = 10$ metric:

$$\sqrt{-\det G} G^{00} = \sqrt{-\det g} g^{00} \quad (8.12)$$

having denoted G the full Einstein metric in ten dimension and g the Einstein metric in three dimension.

Specializing to our example Hence, in our example the ansatz for G_{ij} is diagonal

$$G_{ij} = \exp[2\sigma_i] \delta_{ij} \quad ; \quad i = 1, \dots, 7 \quad (8.13)$$

and we obtain the following relation between the fields x and y and the diagonal entries of the metric and the dilaton:

$$\begin{aligned} \phi = -\vec{h} \cdot \vec{\alpha}[7] &= x + y \\ \sigma_{1,2} &= \frac{3x - y}{4} \\ \sigma_{3,4,5} &= \frac{3y - x}{4} \\ \sigma_{6,7} &= -\frac{5y + x}{4} \end{aligned} \quad (8.14)$$

Calling $\tilde{h}_{1,2}$ the Cartan fields in the abstract A_2 model discussed in chapter 6, we have:

$$\tilde{h} \cdot \beta_1 = \sqrt{2} \tilde{h}_1 \quad ; \quad \tilde{h} \cdot \beta_2 = -\frac{1}{\sqrt{2}} \tilde{h}_1 + \sqrt{\frac{3}{2}} \tilde{h}_1 \quad (8.15)$$

so that we can conclude:

$$x = \frac{1}{\sqrt{2}} \tilde{h}_1 \quad ; \quad y = -\frac{1}{\sqrt{6}} \tilde{h}_1 \quad (8.16)$$

We can also immediately conclude that:

$$Q^2 = \frac{d}{dt} h \cdot \frac{d}{dt} h = \frac{d}{dt} \tilde{h} \cdot \frac{d}{dt} \tilde{h} = |\chi_1|^2 + |\chi_2|^2 \quad (8.17)$$

On the other hand the interpretation of Q^2 is the following. Consider the parameter ϖ^2 appearing in the three-dimensional metric determined from the sigma model by Einstein equations. It is defined as:

$$\varpi^2 = h_{IJ} \dot{\phi}^I \dot{\phi}^J = \sum_{i=1}^8 |\chi_i|^2 + \sum_{\alpha=1}^{120} |\Phi|^2 \quad (8.18)$$

If we calculate ϖ^2 using the generating solution or any other solution obtained from it by compensating H-transformations, its value, which is a constant, does not change. So we have:

$$\varpi^2 = \sum_{i=1}^8 |\chi_i^{(\text{gen.sol.})}|^2 \quad (8.19)$$

and in the lifting of our A_2 solutions we can conclude that $Q^2 = \varpi^2$. Let us calculate this crucial parameter for the case of the non trivial A_2 solutions discussed above. By means of straightforward algebra we get:

$$\varpi^2 = |\chi_1^{(\text{gen.sol.})}|^2 + |\chi_2^{(\text{gen.sol.})}|^2 = \frac{1}{24} (\kappa^2 + 3\omega^2) \quad (8.20)$$

Next we turn to the identification of the p -forms. As we will explicitly verify by checking type IIB supergravity field equations, the appropriately normalized identifications are the following ones:

$$\begin{aligned} B_{[2]} &= \varphi_1(t) dx_3 \wedge dx_4 \\ C_{[2]} &= \varphi_2(t) dx_8 \wedge dx_9 \\ C_{[4]} &= \varphi_3(t) dx_3 \wedge dx_4 \wedge dx_8 \wedge dx_9 + U \end{aligned} \quad (8.21)$$

where U is the appropriate 4-form needed to make the corresponding field strength self dual.

In this way recalling the normalizations of type IIB field strengths as given in appendix we get:

$$\begin{aligned} F_{[3]034}^{NS} &= \frac{1}{6} \varphi_1'(t) \\ F_{[3]089}^{RR} &= \frac{1}{6} \varphi_2'(t) \\ F_{[5]03489}^{RR} &= \frac{1}{240} W(t) \\ F_{[5]12567}^{RR} &= \frac{1}{240} W(t) \sqrt{-\det g} \frac{1}{g_{00} g_{33} g_{44} g_{88} g_{99}} \end{aligned} \quad (8.22)$$

and we recognize that the combination $W(t)$ defined in eq.(6.28) is just the self-dual 5-form field strength including Chern-Simons factors.

8.2.1 Full oxidation of the A_2 solution with only one root switched on

Let us now focus on the A_2 solution involving only the highest root (similar solutions were obtained in [11, 12, 13, 14, 15, 16, 17, 18, 113, 114, 115, 116, 117]), namely on eq.s (6.47) and (6.48). Inserting the explicit form of the Cartan fields in eq.s(8.16) and then using (8.14) we obtain the complete form of the metric

$$\begin{aligned} ds^2 &= -r_{[0]}^2(t) dt^2 + r_{[1|2]}^2(t) (dx_1^2 + dx_2^2) + r_{[3|4]}^2(t) (dx_3^2 + dx_4^2) \\ &\quad + r_{[5|6|7]}^2(t) (dx_5^2 + dx_6^2 + dx_7^2) + r_{[8|9]}^2(t) (dx_8^2 + dx_9^2) \end{aligned} \quad (8.23)$$

which is diagonal and it is parametrized by five time dependent *scale factors*

$$\begin{aligned}
r_{[0]}^2(t) &= e^{t\sqrt{\frac{\kappa^2}{3}+\omega^2}} \sqrt{\cosh \frac{t\omega}{2}} \\
r_{[1|2]}^2(t) &= e^{\frac{t\sqrt{\frac{\kappa^2}{3}+\omega^2}}{2}} \sqrt{\cosh \frac{t\omega}{2}} \\
r_{[3|4]}^2(t) &= \frac{1}{e^{\frac{t\kappa}{6}} \sqrt{\cosh \frac{t\omega}{2}}} \\
r_{[5|6|7]}^2(t) &= \sqrt{\cosh \frac{t\omega}{2}} \\
r_{[8|9]}^2(t) &= \frac{e^{\frac{t\kappa}{6}}}{\sqrt{\cosh \frac{t\omega}{2}}}
\end{aligned} \tag{8.24}$$

We also obtain the explicit form of the dilaton, which turns out to be linear in time:

$$\phi = -\frac{1}{6} \kappa t \tag{8.25}$$

Calculating the Ricci tensor of the metric (8.23) we find that it is also diagonal and it has five independent eigenvalues respectively given by:

$$\begin{aligned}
Ric_{00} &= \frac{(\kappa^2 + 9\omega^2 + \kappa^2 \cosh t\omega) \operatorname{sech}^2(\frac{t\omega}{2})}{288} \\
Ric_{11} = Ric_{22} &= \frac{\omega^2 \operatorname{sech}^2(\frac{t\omega}{2})}{32 e^{\frac{t\sqrt{\frac{\kappa^2}{3}+\omega^2}}{2}}} \\
Ric_{33} = Ric_{44} &= \frac{\omega^2 \operatorname{sech}^2(\frac{t\omega}{2})}{32 e^{\frac{t\sqrt{\frac{\kappa^2}{3}+\omega^2}}{2}}} \\
Ric_{55} = Ric_{66} = Ric_{77} &= \frac{-\omega^2 \operatorname{sech}^3(\frac{t\omega}{2})}{32 e^{\frac{t(\kappa+6\sqrt{\frac{\kappa^2}{3}+\omega^2})}{6}}} \\
Ric_{88} = Ric_{99} &= \frac{-\omega^2 \operatorname{sech}^3(\frac{t\omega}{2})}{32 e^{\frac{t(\kappa+6\sqrt{\frac{\kappa^2}{3}+\omega^2})}{6}}}
\end{aligned} \tag{8.26}$$

On the other hand inserting the explicit values of scalar fields (6.48) into equations (8.21) we obtain:

$$\begin{aligned}
F_{[3]}^{NS} &= 0 \\
F_{[3]}^{RR} &= 0 \\
F_{[5]}^{RR} &= \frac{\omega dt \wedge dx_3 \wedge dx_4 \wedge dx_8 \wedge dx_9}{1 + \cosh t\omega} + \frac{\omega dx_1 \wedge dx_2 \wedge dx_5 \wedge dx_6 \wedge dx_7}{2}
\end{aligned} \tag{8.27}$$

Considering eq.s(8.27) and (8.25) together the physical interpretation of the parameters ω and κ labeling the generating solution, becomes clear. They are respectively associated to the charges of the $D3$ and $D5$ branes which originate this classical supergravity solution. Indeed, as it is obvious from the last of

eq.s (8.27), there is a dyonic $D3$ -brane whose magnetic charge is uniformly distributed on the Euclidean hyperplane 12567 while the electric charge is attached to the Minkowskian hyperplane 03489. The magnetic charge per unit volume is $\omega/2$. With our choice of the A_2 subalgebra, there should also be a $D5$ -brane magnetically dual to an Euclidean D -string extending in the directions 89. In this particular solution, where $\varphi_{1,2} = 0$ the $F_{[3]}^{RR}$ vanishes, yet the presence of the $D5$ brane is revealed by the dilaton. Indeed in a pure $D3$ brane solution the dilaton would be constant. The linear behaviour (8.25) of ϕ , with coefficient $-\kappa/6$ is due to the $D5$ brane which couples non trivially to the dilaton field. Such an interpretation will become completely evident when we consider the oxidation of the solution obtained from this by a further $SO(3)$ rotation which switches on all the roots. This we do in the next section. Then we will discuss how both oxidations do indeed satisfy the field equations of type IIB supergravity and we will illustrate their physical properties as cosmic backgrounds.

8.2.2 Full oxidation of the A_2 solution with all three roots switched on

Let us then turn to the A_2 solution involving all the three nilpotent fields, namely to eq.s (6.51) and (6.52). Just as before, by inserting the explicit form of the Cartan fields in eq.s(8.16) and then using (8.14) we obtain the complete form of the new metric, which has the same diagonal structure as in the previous example, namely

$$ds^2 = -\bar{r}_{[0]}^2(t) dt^2 + \bar{r}_{[1|2]}^2(t) (dx_1^2 + dx_2^2) + \bar{r}_{[3|4]}^2(t) (dx_3^2 + dx_4^2) + \bar{r}_{[5|6|7]}^2(t) (dx_5^2 + dx_6^2 + dx_7^2) + \bar{r}_{[8|9]}^2(t) (dx_8^2 + dx_9^2) \quad (8.28)$$

now, however, the *scale factors* are given by:

$$\begin{aligned} \bar{r}_{[0]}^2(t) &= e^{t\left(\frac{-\omega}{4} + \sqrt{\frac{\kappa^2}{3} + \omega^2}\right)} (1 + e^{t\omega})^{\frac{1}{4}} \left(1 + e^{t\omega} + e^{\frac{t(\kappa+\omega)}{2}}\right)^{\frac{1}{4}} \\ \bar{r}_{[1|2]}^2(t) &= e^{\frac{t(-3\omega+2\sqrt{3}\sqrt{\kappa^2+3\omega^2})}{12}} (1 + e^{t\omega})^{\frac{1}{4}} \left(1 + e^{t\omega} + e^{\frac{t(\kappa+\omega)}{2}}\right)^{\frac{1}{4}} \\ \bar{r}_{[3|4]}^2(t) &= \frac{e^{\frac{-(t\kappa)}{6} + \frac{t\omega}{4}} \left(1 + e^{t\omega} + e^{\frac{t(\kappa+\omega)}{2}}\right)^{\frac{1}{4}}}{(1 + e^{t\omega})^{\frac{3}{4}}} \\ \bar{r}_{[5|6|7]}^2(t) &= \frac{(1 + e^{t\omega})^{\frac{1}{4}} \left(1 + e^{t\omega} + e^{\frac{t(\kappa+\omega)}{2}}\right)^{\frac{1}{4}}}{e^{\frac{t\omega}{4}}} \\ \bar{r}_{[8|9]}^2(t) &= \frac{e^{\frac{t(2\kappa+3\omega)}{12}} (1 + e^{t\omega})^{\frac{1}{4}}}{\left(1 + e^{t\omega} + e^{\frac{t(\kappa+\omega)}{2}}\right)^{\frac{3}{4}}} \end{aligned} \quad (8.29)$$




Figure 8.1: Plots of the scale factors $r_{[\alpha]}^2$, $\alpha = 1|2, 3|4, 5|6|7, 8|9$ as functions of the cosmic time $t = \tau(T)$ in the case of the choice of parameters $\omega = 1$, $\kappa = 0.5$ and for the A_2 solution with only the highest root switched on.

and the dilaton is no longer linear in time, rather it is given by:

$$\phi = \frac{-(t\kappa) - 3 \log(1 + e^{t\omega}) + 3 \log(1 + e^{t\omega} + e^{\frac{t(\kappa+\omega)}{2}})}{6} \quad (8.30)$$

Calculating the Ricci tensor of the metric (8.28,8.29) we find it diagonal with five different eigenvalues, just as in the previous case, but with a modified time dependence, namely:

$$\begin{aligned} Ric_{00} = & \frac{1}{576 (1 + e^{t\omega})^2 \left(1 + e^{t\omega} + e^{\frac{t(\kappa+\omega)}{2}}\right)^2} \\ & \times \left[(1 + e^{t\omega})^2 \left(4 + 8e^{t\omega} + 4e^{2t\omega} + 23e^{\frac{t(\kappa+\omega)}{2}} + e^{t(\kappa+\omega)} + 23e^{\frac{t(\kappa+3\omega)}{2}}\right) \kappa^2 \right. \\ & \quad \left. - 6e^{\frac{t(\kappa+\omega)}{2}} (-1 + e^{2t\omega}) \left(7 + 7e^{t\omega} + e^{\frac{t(\kappa+\omega)}{2}}\right) \kappa \omega \right. \\ & \quad \left. 9 \left(8e^{t\omega} + 16e^{2t\omega} + 8e^{3t\omega} + 3e^{\frac{t(\kappa+\omega)}{2}} + e^{t(\kappa+\omega)} + 10e^{t(\kappa+2\omega)} \right. \right. \\ & \quad \left. \left. + 17e^{\frac{t(\kappa+3\omega)}{2}} + e^{t(\kappa+3\omega)} + 17e^{\frac{t(\kappa+5\omega)}{2}} + 3e^{\frac{t(\kappa+7\omega)}{2}}\right) \omega^2 \right] \quad (8.31a) \end{aligned}$$




Figure 8.2: Plots of the scale factors $r_{[\alpha]}^2$, $\alpha = 1/2, 3/4, 5/6, 7/8, 8/9$ as functions of the cosmic time $t = \tau(T)$ in the case of the choice of parameters $\omega = 0$, $\kappa = 1$ and for the A_2 solution with only the highest root switched on.




Figure 8.3: Plots of the scale factors $r_{[\alpha]}^2$, $\alpha = 1/2, 3/4, 5/6, 7/8, 8/9$ as functions of the cosmic time $t = \tau(T)$ in the case of the choice of parameters $\omega = 1$, $\kappa = 0$ and for the A_2 solution with only the highest root switched on.

$$\begin{aligned}
Ric_{11} = Ric_{22} = & \frac{1}{64 (1 + e^{t\omega})^2 \left(1 + e^{t\omega} + e^{\frac{t(\kappa+\omega)}{2}}\right)^2} \\
& \times \left[e^{\frac{t\left(\omega - \sqrt{\frac{\kappa^2}{3} + \omega^2}\right)}{2}} \left(e^{\frac{t(\kappa+6\omega)}{2}} (\kappa - \omega)^2 + 8 e^{\frac{t\omega}{2}} \omega^2 + 16 e^{\frac{3t\omega}{2}} \omega^2 + 8 e^{\frac{5t\omega}{2}} \omega^2 \right. \right. \\
& \quad \left. \left. + 4 e^{t\left(\kappa + \frac{3\omega}{2}\right)} \omega^2 + e^{\frac{t\kappa}{2}} (\kappa + \omega)^2 + e^{\frac{t(\kappa+4\omega)}{2}} (3\kappa^2 - 2\kappa\omega + 11\omega^2) \right. \right. \\
& \quad \left. \left. + e^{\left(\frac{t\kappa}{2} + t\omega\right)} (3\kappa^2 + 2\kappa\omega + 11\omega^2) \right) \right] \quad (8.31b)
\end{aligned}$$

$$\begin{aligned}
Ric_{33} = Ric_{44} = & \frac{1}{64 e^{\frac{t\left(\kappa - 6\omega + 6\sqrt{\frac{\kappa^2}{3} + \omega^2}\right)}{6}} (1 + e^{t\omega})^3 \left(1 + e^{t\omega} + e^{\frac{t(\kappa+\omega)}{2}}\right)^2} \\
& \times \left[e^{\frac{t(\kappa+6\omega)}{2}} (\kappa - \omega)^2 - 8 e^{\frac{t\omega}{2}} \omega^2 - 16 e^{\frac{3t\omega}{2}} \omega^2 - 8 e^{\frac{5t\omega}{2}} \omega^2 - 12 e^{t\left(\kappa + \frac{3\omega}{2}\right)} \omega^2 \right. \\
& \quad \left. + e^{\frac{t\kappa}{2}} (\kappa + \omega)^2 + e^{\frac{t(\kappa+4\omega)}{2}} (3\kappa^2 - 2\kappa\omega - 21\omega^2) \right. \\
& \quad \left. + e^{\frac{t(\kappa+2\omega)}{2}} (3\kappa^2 + 2\kappa\omega - 21\omega^2) \right] \quad (8.31c)
\end{aligned}$$

$$\begin{aligned}
Ric_{55} = Ric_{66} = Ric_{77} = & \frac{1}{64 (1 + e^{t\omega})^2 \left(1 + e^{t\omega} + e^{\frac{t(\kappa+\omega)}{2}}\right)^2} \\
& \times \left[e^{\frac{t\left(\omega - 2\sqrt{\frac{\kappa^2}{3} + \omega^2}\right)}{2}} \left[e^{\frac{t(\kappa+6\omega)}{2}} (\kappa - \omega)^2 + 8 e^{\frac{t\omega}{2}} \omega^2 + 16 e^{\frac{3t\omega}{2}} \omega^2 + 8 e^{\frac{5t\omega}{2}} \omega^2 \right. \right. \\
& \quad \left. \left. + 4 e^{t\left(\kappa + \frac{3\omega}{2}\right)} \omega^2 + e^{\frac{t\kappa}{2}} (\kappa + \omega)^2 e^{\frac{t(\kappa+4\omega)}{2}} (3\kappa^2 - 2\kappa\omega + 11\omega^2) \right. \right. \\
& \quad \left. \left. + e^{\frac{t(\kappa+2\omega)}{2}} (3\kappa^2 + 2\kappa\omega + 11\omega^2) + \right] \right] \quad (8.31d)
\end{aligned}$$

$$\begin{aligned}
Ric_{88} = Ric_{99} = & - \frac{1}{64 (1 + e^{t\omega})^2 \left(1 + e^{t\omega} + e^{\frac{t(\kappa+\omega)}{2}}\right)^3} \\
& \times \left[e^{\frac{t\left(\kappa + 6\omega - 6\sqrt{\frac{\kappa^2}{3} + \omega^2}\right)}{6}} \left[3 e^{\frac{t(\kappa+6\omega)}{2}} (\kappa - \omega)^2 + 8 e^{\frac{t\omega}{2}} \omega^2 + 16 e^{\frac{3t\omega}{2}} \omega^2 + 8 e^{\frac{5t\omega}{2}} \omega^2 \right. \right. \\
& \quad \left. \left. - 4 e^{t\left(\kappa + \frac{3\omega}{2}\right)} \omega^2 + e^{\frac{t(\kappa+4\omega)}{2}} (-3\kappa + \omega)^2 + 3 e^{\frac{t\kappa}{2}} (\kappa + \omega)^2 + e^{\frac{t(\kappa+2\omega)}{2}} (3\kappa + \omega)^2 \right] \right] \quad (8.31e)
\end{aligned}$$

On the other hand inserting the explicit values of scalar fields (6.52) into

equations (8.21) we obtain:

$$F_{[3]}^{NS} = -\frac{1}{4}\omega \operatorname{sech}^2 \frac{t\omega}{2} dt \wedge dx_3 \wedge dx_4 \quad (8.32a)$$

$$F_{[3]}^{RR} = \frac{e^{\frac{t(\kappa+\omega)}{2}} (\kappa + e^{t\omega}(\kappa - \omega) + \omega)}{2 \left(1 + e^{t\omega} + e^{\frac{t(\kappa+\omega)}{2}}\right)^2} dt \wedge dx_8 \wedge dx_9 \quad (8.32b)$$

$$F_{[5]}^{RR} = \frac{e^{t\omega} \omega dt \wedge dx_3 \wedge dx_4 \wedge dx_8 \wedge dx_9}{(1 + e^{t\omega}) \left(1 + e^{t\omega} + e^{\frac{t(\kappa+\omega)}{2}}\right)} - \omega dx_1 \wedge dx_2 \wedge dx_5 \wedge dx_6 \wedge dx_7 \quad (8.32c)$$

This formula completes the oxidation also of the second sigma model solution to a full fledged $D = 10$ type IIB configuration. As expected in both cases the ten dimensional fields obtained by oxidation satisfy the field equations of supergravity as formulated in the appendix. We discuss this in the next chapter.

8.2.3 How the supergravity field equations are satisfied and their cosmological interpretation

Taking into account that the Ramond scalar C_0 vanishes the effective bosonic field equations of supergravity reduce to:

$$d \star d\phi = \frac{1}{2} (e^{-\phi} F_3^{NS} \wedge \star F_3^{NS} - e^{\phi} F_3^{RR} \wedge \star F_3^{RR}) \quad (8.33)$$

$$0 = F_3^{NS} \wedge \star F_3^{NS} \quad (8.34)$$

$$d(e^{-\phi} F_3^{NS}) = -F_3^{RR} \wedge \star F_5^{RR} \quad (8.35)$$

$$d(e^{\phi} F_3^{RR}) = F_3^{NS} \wedge \star F_5^{RR} \quad (8.36)$$

$$d(\star F_3^{RR}) = -F_3^{NS} \wedge F_3^{RR} \quad (8.37)$$

$$-2Ric_{MN} = \hat{T}_{MN} \quad (8.38)$$

where the reduced stress energy tensor $\hat{T}_{MN}$ is the superposition of two contributions that we respectively attribute to the $D3$ brane and to the $D5$ -brane, namely:

$$\hat{T}_{MN} = \hat{T}_{MN}^{[D3]} + \hat{T}_{MN}^{[D5]} \quad (8.39)$$

$$\hat{T}_{MN}^{[D3]} \equiv 150 F_{[5]M\dots} F_{[5]N}^{\dots} \quad (8.40)$$

$$\begin{aligned} \hat{T}_{MN}^{[D5]} &\equiv \frac{1}{2} \partial_M \varphi \partial_N \varphi + 9 \left(e^{-\varphi} F_{[3]M\dots}^{NS} F_{[3]N\dots}^{NS} + e^{\varphi} F_{[3]M\dots}^{RR} F_{[3]N\dots}^{RR} \right) \\ &\quad - \frac{3}{4} g_{MN} \left(e^{-\varphi} F_{[3]\dots}^{NS} F_{[3]\dots}^{NS} + e^{\varphi} F_{[3]\dots}^{RR} F_{[3]\dots}^{RR} \right) \end{aligned} \quad (8.41)$$

By means of laborious algebraic manipulations that can be easily performed on a computer with the help of MATHEMATICA, we have explicitly verified that in both cases, that of chapter 8.2.1 and that of chapter 8.2.2 the field eq.s(8.33)-(8.38) are indeed satisfied, so that the oxidation procedure we have described turns out to be well tuned and fully correct.




Figure 8.4: Plots of the energy densities in the case of the choice of parameters $\omega = 1$, $\kappa = 0.5$ and for the A_2 solution with only the highest root switched on. The first picture plots the total density $\rho^{tot}(\tau)$. The second picture plots the ratio $\rho^{d3}(\tau)/\rho^{tot}(\tau)$ and the third plots the ratio $\rho^{d3}(\tau)/\rho^{d5}(\tau)$




Figure 8.5: Plots of the pressure eigenvalues $P_{[\alpha]}$, $\alpha = 1|2, 3|4, 5|6|7, 8|9$ as functions of the cosmic time $t = \tau(T)$ in the case of the choice of parameters $\omega = 1$, $\kappa = 0.5$ and for the A_2 solution with only the highest root switched on.

In order to enlighten the physical meaning of the type IIB superstring backgrounds we have eventually constructed it is worth to analyze the structure of

the stress energy tensor. First, reintroducing the missing traces we define:

$$\begin{aligned} T_{MN}^{[D3]} &= \hat{T}_{MN}^{[D3]} - \frac{1}{2} g_{MN} \hat{T}_{RS}^{[D3]} g^{RS} \\ T_{MN}^{[D5]} &= \hat{T}_{MN}^{[D5]} - \frac{1}{2} g_{MN} \hat{T}_{RS}^{[D5]} g^{RS} \\ T_{MN}^{tot} &= T_{MN}^{[D3]} + T_{MN}^{[D5]} \end{aligned} \quad (8.42)$$

It turns out that the stress energy tensors are diagonal, just as the metric, and have the form of a perfect fluid, but with different pressure eigenvalues in the various subspaces. Indeed we can write:

$$\begin{aligned} T_{00}^{tot, D3, D5} &= g_{00} \rho^{tot, D3, D5} \\ T_{i_\alpha j_\alpha}^{tot, D3, D5} &= -g_{i_\alpha j_\alpha} P_\alpha^{tot, D3, D5} \end{aligned} \quad (8.43)$$

where α denotes the four different submanifolds extending in directions:

$$\alpha = 1|2, 3|4, 5|6|7, 8|9 \quad (8.44)$$

We can now analyze the specific properties of the two example of solutions.

8.2.4 Properties of the solution with just one root switched on

In the case of the time dependent background described in chapter (8.2.1) and obtained by oxidizing the solution (6.48) we obtain for the *energy densities*:

$$\begin{aligned} \rho^{tot} &= \frac{(1 + e^{t\omega})^2 \kappa^2 + 9 e^{t\omega} \omega^2}{36 e^{t\sqrt{\frac{\kappa^2}{3} + \omega^2}} (1 + e^{t\omega})^2 \sqrt{\cosh \frac{t\omega}{2}}} \\ \rho^{d3} &= \frac{\omega^2}{16 e^{t\sqrt{\frac{\kappa^2}{3} + \omega^2}} \cosh^{\frac{5}{2}} \frac{t\omega}{2}} \\ \rho^{d5} &= \frac{\kappa^2}{36 e^{t\sqrt{\frac{\kappa^2}{3} + \omega^2}} \sqrt{\cosh \frac{t\omega}{2}}} \end{aligned} \quad (8.45)$$

for the *total pressures*:

$$\begin{aligned} P_{1|2}^{tot} &= -\frac{\kappa^2 + 9\omega^2 + \kappa^2 \cosh t\omega}{144 e^{t\sqrt{\frac{\kappa^2}{3} + \omega^2}} \cosh^{\frac{5}{2}} \frac{t\omega}{2}} \\ P_{3|4}^{tot} &= -\frac{e^{t\left(\omega - \sqrt{\frac{\kappa^2}{3} + \omega^2}\right)} (\kappa^2 - 9\omega^2 + \kappa^2 \cosh t\omega)}{36(1 + e^{t\omega})^2 \sqrt{\cosh \frac{t\omega}{2}}} \\ P_{5|6|7}^{tot} &= -\frac{\kappa^2 + 9\omega^2 + \kappa^2 \cosh t\omega}{144 e^{t\sqrt{\frac{\kappa^2}{3} + \omega^2}} \cosh^{\frac{5}{2}} \frac{t\omega}{2}} \\ P_{8|9}^{tot} &= -\frac{e^{t\left(\omega - \sqrt{\frac{\kappa^2}{3} + \omega^2}\right)} (\kappa^2 - 9\omega^2 + \kappa^2 \cosh t\omega)}{36(1 + e^{t\omega})^2 \sqrt{\cosh \frac{t\omega}{2}}} \end{aligned} \quad (8.46)$$

for the pressures associated with the $D3$ brane:

$$\begin{aligned}
P_{1|2}^{D3} &= \frac{-\omega^2}{16 e^t \sqrt{\frac{\kappa^2}{3} + \omega^2} \cosh^{\frac{5}{2}} \frac{t\omega}{2}} \\
P_{3|4}^{D3} &= \frac{\omega^2}{16 e^t \sqrt{\frac{\kappa^2}{3} + \omega^2} \cosh^{\frac{5}{2}} \frac{t\omega}{2}} \\
P_{5|6|7}^{D3} &= \frac{-\omega^2}{16 e^t \sqrt{\frac{\kappa^2}{3} + \omega^2} \cosh^{\frac{5}{2}} \frac{t\omega}{2}} \\
P_{8|9}^{D3} &= \frac{\omega^2}{16 e^t \sqrt{\frac{\kappa^2}{3} + \omega^2} \cosh^{\frac{5}{2}} \frac{t\omega}{2}}
\end{aligned} \tag{8.47}$$

and for the pressures associated with the $D5$ brane:

$$\begin{aligned}
P_{1|2}^{D5} &= \frac{-\kappa^2}{72 e^t \sqrt{\frac{\kappa^2}{3} + \omega^2} \sqrt{\cosh \frac{t\omega}{2}}} \\
P_{3|4}^{D5} &= \frac{-\kappa^2}{72 e^t \sqrt{\frac{\kappa^2}{3} + \omega^2} \sqrt{\cosh \frac{t\omega}{2}}} \\
P_{5|6|7}^{D5} &= \frac{-\kappa^2}{72 e^t \sqrt{\frac{\kappa^2}{3} + \omega^2} \sqrt{\cosh \frac{t\omega}{2}}} \\
P_{8|9}^{D5} &= \frac{-\kappa^2}{72 e^t \sqrt{\frac{\kappa^2}{3} + \omega^2} \sqrt{\cosh \frac{t\omega}{2}}}
\end{aligned} \tag{8.48}$$

As we see from its analytic expression the total energy density is an exponentially decreasing function of time which tends to zero at asymptotically late times ($t \mapsto \infty$). What happens instead at asymptotically early times ($t \mapsto -\infty$) depends on the value of κ . For $\kappa = 0$ we have $\lim_{t \mapsto -\infty} \rho^{tot}(t) = 0$, while for $\kappa \neq 0$ we always have $\lim_{t \mapsto -\infty} \rho^{tot}(t) = \infty$. This is illustrated, for instance, in figs.(8) and (10). This phenomenon is related to the presence or absence of a $D5$ brane as it is evident from eq.s (8.45) which shows that the *dilaton*-($D5$) brane contribution to the energy density is proportional to κ^2 and it is always divergent at asymptotically early times, while the $D3$ brane contribution tends to zero in the same regime.

We also note, comparing eq.s(8.48) with eq.s(8.47) that the pressure contributed by the *dilaton*- $D5$ -brane system is the same in all directions 1–9, while the pressure contributed by the $D3$ -brane system is just opposite in the direction 3489 and in the transverse directions 12567. This is the origin of the *cosmic billiard phenomenon* that we observe in the behaviour of the metric scale factors. Indeed the presence of the $D3$ -brane causes, at a certain instant of time, a switch in the cosmic expansion. Dimensions that were previously shrinking begin to expand and dimensions that were expanding begin to shrink. It is like




Figure 8.6: Plot of the energy density as function of the cosmic time $t = \tau(T)$ in the case of a pure $D3$ brane system, namely for the the choice of parameters $\omega = 1$, $\kappa = 0$ and for the A_2 solution with only the highest root switched on.

a ball that hits a wall and inverts its speed. In the exact solution that we have constructed through reduction to three dimensions this occurs in a smooth way. There is a maximum and respectively a minimum in the behaviour of certain scale factors, which is in relation with a predominance of the $D3$ -brane energy density with respect to the total energy density. The cosmic $D3$ -brane behaves just as an instanton. Its contribution to the total energy is originally almost zero, then it raises and dominates for some time, then it exponentially decays again. This is the smooth exact realization of the potential walls envisaged by Damour et al.

To appreciate such a behaviour it is convenient to consider some plots of the scale factors, the energy densities and the pressures. In order to present such plots we first reduce the metric (8.23) to a standard cosmological form, by introducing a new time variable τ such that:

$$r_{[0]}(t) dt = d\tau \quad (8.49)$$

Explicitly we set:

$$\tau = \int_0^T r_{[0]}(t) dt \quad (8.50)$$

and inserting the explicit form of the scale factor $r_{[0]}(t)$ as given in eq.(8.24) we obtain:

$$\begin{aligned} \tau(T) = & \frac{4 \cdot 2^{\frac{3}{4}}}{\omega - 4 \sqrt{\frac{\kappa^2}{3} + \omega^2}} \left[{}_2F_1 \left(-\frac{1}{8} + \frac{\sqrt{\frac{\kappa^2}{3} + \omega^2}}{2\omega}, -\frac{1}{4}, \frac{7}{8} + \frac{\sqrt{\frac{\kappa^2}{3} + \omega^2}}{2\omega}, -1 \right) \right. \\ & \left. - e^{\frac{\tau(-\omega + 4 \sqrt{\frac{\kappa^2}{3} + \omega^2})}{8}} {}_2F_1 \left(-\frac{1}{8} + \frac{\sqrt{\frac{\kappa^2}{3} + \omega^2}}{2\omega}, -\frac{1}{4}, \frac{7}{8} + \frac{\sqrt{\frac{\kappa^2}{3} + \omega^2}}{2\omega}, -e^{T\omega} \right) \right] \quad (8.51) \end{aligned}$$

which expresses τ in terms of hypergeometric functions and exponentials.

In fig.(8.1) we observe the billiard phenomenon in a generic case where both parameters ω and κ are non vanishing. Since the value of ω can always be

rescaled by a rescaling of the original time coordinate t , we can just set it to 1 and what matters is to distinguish the case $\omega \neq 0$ where the $D3$ brane is present from the case $\omega = 0$ corresponding to its absence. Hence fig.(8.1) corresponds to the presence of both a $D3$ -brane and a *dilaton- $D5$* -brane system. A very different behavior occurs in fig.(8.2) where $\omega = 0$. In this case there is no billiard and the dimensions either shrink or expand uniformly. On the other hand in fig.(8.3) we observe the pure billiard phenomenon induced by the $D3$ -brane in the case where no *dilaton- $D5$* -brane is present, namely when we set $\kappa = 0$. In this case, as we see, the parallel directions to the Euclidean $D3$ -brane, namely 3489 have exactly the same behaviour: they first inflate and then they deflate, namely there is a maximum in the scale factor. The transverse directions to the $D3$ brane 567 have the opposite behaviour. They display a minimum at the same point where the parallel directions display a maximum. In all cases the directions 12 corresponding to the spatial directions of the three dimensional sigma model world suffer a uniform expansion.

Let us now consider the behavior of the energy densities. In fig.(8.4) we focus on the mixed case $\omega = 1$, $\kappa = 0.5$ characterized by the presence of both a $D3$ brane and *dilaton- $D5$* -brane system. As we see the total energy density exponentially decreases at late times and has a singularity at asymptotically early times. This is like in a standard Big Bang cosmological model with an indefinite expansion starting from an initial singularity. Yet the ratio of the $D3$ energy with respect to the total energy has a maximum at some instant of time and this is the cause of the billiard phenomenon in the behaviour of the scale factors respectively parallel and transverse to the $D3$ brane itself. The two contributions to the energy density from the $D3$ -brane and from the dilaton have the same sign and the plot of their ratio displays a maximum in correspondence with the billiard time. With the same choice of parameters $\omega = 1$, $\kappa = 0.5$ the physical behavior of the system can be appreciated by looking at the plots of the pressure eigenvalues. They are displayed in fig.(8.5). We observe that the pressure is negative in the directions transverse to the $D3$ brane 12 and 567. Slowly, but uniformly it increases to zero in these directions. In the directions parallel to the brane the pressure is instead always positive and it displays a sharp maximum at the instant of time where the billiard phenomenon occurs.

For a pure $D3$ brane system, namely for $\kappa = 0$ and $\omega = 1$ the energy density starts at zero, develops a maximum and then decays again to zero. This can be seen in fig.(8.6). The plot of the pressures is displayed, for this case in fig.(8.7). In this case the pressure in the directions transverse to the brane, i.e. 12567 is negative and it is just the opposite of the pressure in the directions parallel to the brane, namely 3489. This behavior causes the corresponding scale factors to suffer a minimum and a maximum, respectively.




Figure 8.7: Plots of the pressure eigenvalues $P_{[\alpha]}$, $\alpha = 1|2, 3|4, 5|6|7, 8|9$ as functions of the cosmic time $t = \tau(T)$ in the case of the choice of parameters $\omega = 1$, $\kappa = 0$ and for the A_2 solution with only the highest root switched on. This case corresponds to a pure $D3$ brane system.

8.2.5 Properties of the solution with all roots switched on

Let us now discuss the properties of the second solution where all the roots have been excited. In chapter 8.2.2 we considered the oxidation of such a sigma model solution and we constructed the corresponding $D = 10$ supergravity background given by the metric (8.28, 8.29) and by the field strengths (8.32). Looking at eq.s (8.32) we see that the interpretation of the parameter ω is still the same as it was before, namely it represents the magnetic charge of the dyonic $D3$ -brane. At $\omega = 0$ the $D3$ -brane disappears. Yet it appears from eq.s (8.32) that there is no obvious interpretation of the parameter κ as a pure $D5$ -brane charge. Indeed there is no choice of κ which suppresses both the NS and the RR 3-form field strengths.

Following the same procedure as in the previous case we calculate the energy density and the pressures and we separate the contributions due to the $D3$ -brane and to the *dilaton*- $D5$ -brane system. After straightforward but lengthy algebraic manipulations, implemented on a computer with MATHEMATICA we obtain:

$$\rho^{d3} = \frac{\omega^2}{4e^{t\left(\frac{-5\omega}{4} + \sqrt{\frac{\kappa^2}{3} + \omega^2}\right)} (1 + e^{t\omega})^{\frac{5}{4}} \left(1 + e^{t\omega} + e^{\frac{t(\kappa + \omega)}{2}}\right)^{\frac{5}{4}}} \quad (8.52a)$$

$$\begin{aligned}
\rho^{d5} = & \frac{e^{\frac{t}{4} \left(\frac{\omega}{4} - \sqrt{\frac{\kappa^2}{3} + \omega^2} \right)}}{192 (1 + e^{t\omega})^{\frac{9}{4}} \left(1 + e^{t\omega} + e^{\frac{t(\kappa+\omega)}{2}} \right)^{\frac{9}{4}}} \\
& \times \left[(1 + e^{t\omega})^2 \left(4 + 8e^{t\omega} + 4e^{2t\omega} + 20e^{\frac{t(\kappa+\omega)}{2}} + e^{t(\kappa+\omega)} + 20e^{\frac{t(\kappa+3\omega)}{2}} \right) \kappa^2 \right. \\
& - 6e^{\frac{t(\kappa+\omega)}{2}} (-1 + e^{2t\omega}) \left(6 + 6e^{t\omega} + e^{\frac{t(\kappa+\omega)}{2}} \right) \kappa \\
& + 3 \left(8e^{\frac{t(\kappa+\omega)}{2}} + 3e^{t(\kappa+\omega)} + 26e^{t(\kappa+2\omega)} + 24e^{\frac{t(\kappa+3\omega)}{2}} \right. \\
& \left. \left. + 3e^{t(\kappa+3\omega)} + 24e^{\frac{t(\kappa+5\omega)}{2}} + 8e^{\frac{t(\kappa+7\omega)}{2}} \right) \omega^2 \right] \quad (8.52b)
\end{aligned}$$

$$\begin{aligned}
\rho^{tot} = & \frac{e^{\frac{t}{4} \left(\frac{\omega}{4} - \sqrt{\frac{\kappa^2}{3} + \omega^2} \right)}}{192 (1 + e^{t\omega})^{\frac{9}{4}} \left(1 + e^{t\omega} + e^{\frac{t(\kappa+\omega)}{2}} \right)^{\frac{9}{4}}} \\
& \times \left[(1 + e^{t\omega})^2 \left(4 + 8e^{t\omega} + 4e^{2t\omega} + 20e^{\frac{t(\kappa+\omega)}{2}} + e^{t(\kappa+\omega)} + 20e^{\frac{t(\kappa+3\omega)}{2}} \right) \kappa^2 \right. \\
& - 6e^{\frac{t(\kappa+\omega)}{2}} (-1 + e^{2t\omega}) \left(6 + 6e^{t\omega} + e^{\frac{t(\kappa+\omega)}{2}} \right) \kappa \omega \\
& + 3 \left(16e^{t\omega} + 32e^{2t\omega} + 16e^{3t\omega} + 8e^{\frac{t(\kappa+\omega)}{2}} + 3e^{t(\kappa+\omega)} \right. \\
& \left. + 26e^{t(\kappa+2\omega)} + 40e^{\frac{t(\kappa+3\omega)}{2}} + 3e^{t(\kappa+3\omega)} + 40e^{\frac{t(\kappa+5\omega)}{2}} + 8e^{\frac{t(\kappa+7\omega)}{2}} \right) \omega^2 \left. \right] \quad (8.52c)
\end{aligned}$$

We see from the above formulae that the energy density contributed by the $D3$ -brane system is proportional to ω^2 as before and vanishes at $\omega = 0$. However there is no choice of the parameter κ which suppresses the *dilaton*– $D5$ -contribution leaving the $D3$ -contribution non-zero.

The pressure eigenvalues can also be calculated just as in the previous example but the resulting analytic formulae are quite messy and we do not feel them worthy to be displayed. It is rather convenient to consider a few more plots.

Just as in the previous case we define the cosmic time through the formula (8.50). In this case, however, the integral does not lead to a closed formula in terms of special functions and we just have an implicit definition:

$$\tau(T) \equiv \int_0^T e^{\frac{t}{2} \left(\frac{-\omega}{4} + \sqrt{\frac{\kappa^2}{3} + \omega^2} \right)} (1 + e^{t\omega})^{\frac{1}{8}} \left(1 + e^{t\omega} + e^{\frac{t(\kappa+\omega)}{2}} \right)^{\frac{1}{8}} dt \quad (8.53)$$

Let us now observe from eq.s (8.29), (8.32) that there are the following critical values of the parameters:

- 1 For $\omega = 0$ and $\kappa \neq 0$ there is no $D3$ -brane and there is just a D -string dual to a $D5$ -brane.
- 2 For $\kappa = \pm \frac{3}{2}\omega$ the scale factor in the directions 34 tends to a finite asymptotic value respectively at very early or very late times.




Figure 8.8: Plots of the scale factors $\bar{r}_{[\alpha]}^2$, $\alpha = 1|2, 3|4, 5|6|7, 8|9$ as functions of the cosmic time $t = \tau(T)$ with the parameter choice $\omega = 0$, $\kappa = 3/2$ and for the A_2 solution with all the roots switched on.




Figure 8.9: Plots of the pressure eigenvalues $\bar{P}_{[\alpha]}$, $\alpha = 1|2, 3|4, 5|6|7, 8|9$ as functions of the cosmic time $t = \tau(T)$ with the parameter choice $\omega = 0$, $\kappa = 3/2$ and for the A_2 solution with all the roots switched on.

The plot of the scale factors for the choice $\omega = 0$, $\kappa = 3/2$ is given in fig.(8.8)

As already stressed, this a pure D -string system and indeed the billiard phenomenon occurs only in the directions 89 that correspond to the euclidean D -string world-sheet. In all the other directions there is a monotonous behavior of the scale factors. The D -string nature of the solution is best appreciated by looking at the behavior of the pressure eigenvalues, displayed in fig.(8.9)

As we see the positive bump in the pressure now occurs only in the D -string




Figure 8.10: Plot of the energy density as function of the cosmic time $t = \tau(T)$ with the parameter choice $\omega = 0$, $\kappa = 3/2$ and for the A_2 solution with all the roots switched on.




Figure 8.11: Plots of the scale factors $\bar{r}_{[\alpha]}^2$, $\alpha = 1|2, 3|4, 5|6|7, 8|9$ as functions of the cosmic time $t = \tau(T)$ with the parameter choice $\omega = 1$, $\kappa = 0.8$ and for the A_2 solution with all the roots switched on.

directions 89, while in all the other directions the pressure is the same and rises monotonously to zero from large negative values. The pressure bump is in correspondence with the billiard phenomenon. The energy density is instead a monotonously decreasing function of time (see fig.(8.10)).

An intermediate case is provided by the parameter choice $\omega = 1$, $\kappa = 0.8 < 3/2$. The plots of the scale factors are given in fig.(8.11)

The mixture of $D3$ and $D5$ systems is evident from the pictures. Indeed we have now a billiard phenomenon in both the directions 34 and 89 as we expect from a $D3$ -brane, but the maximum in 34 is much sharper than in 89. The maximum in 89 is broader because it takes contribution both from the $D3$ brane and from the D -string.

The phenomenon is best appreciated by considering the plots of the pressure eigenvalues (see figs. (8.12)) and of the energy density (see figs. (8.13)). In the




Figure 8.12: Plots of the pressure eigenvalues $\overline{P}_{[\alpha]}$, $\alpha = 1|2, 3|4, 5|6|7, 8|9$ as functions of the cosmic time $t = \tau(T)$ with the parameter choice $\omega = 1$, $\kappa = 0.8$ and for the A_2 solution with all the roots switched on.

pressure plots we see that there is a positive bump both in the directions 34 and 89, yet the bump in 89 is anticipated at earlier times and it is bigger than the bump in 34, the reason being the cooperation between the $D3$ -brane and D -string contributions. Even more instructive is the plot of the energy densities.

In fig.(8.13) we see that the energy density of the $D3$ -brane has the usual positive bump, while the energy density of the *dilaton*- D -string system has a positive bump followed by a smaller negative one, so that it passes through zero.

At the critical value $\kappa = \frac{3}{2}\omega$ something very interesting occurs in the behaviour of the scale factors.

As we see from fig.(8.14), the scale factor in the direction 34, rather than starting from zero as in all other cases starts from a finite value and then always decreases without suffering a billiard bump. The bump is only in the scale factor 89. Essentially this means that the positive energy of the $D3$ brane and the negative one of the D -string exactly compensate at the origin of time for these critical value of the parameters.

8.2.6 Summarizing the above discussion and the cosmological billiard

Summarizing what we have learned from the numerical analysis of the type IIB cosmological backgrounds obtained by a specific oxidation of the A_2 sigma model solutions we can say what follows.

The expansion or contraction of the cosmological scale factors in the diagonal metric is driven by the presence of euclidean D -branes which behave like




Figure 8.13: Plots of the energy densities as functions of the cosmic time $t = \tau(T)$ with the parameter choice $\omega = 1$, $\kappa = 0.8$ and for the A_2 solution with all the roots switched on. The first picture plots the behaviour of the total energy density. The second plots the ratio of the $D3$ -brane contribution to the energy density with respect to the total density. The third plots the ratio of the $D3$ -brane contribution with respect to the contribution of the dilaton $D5$ -brane system. The fourth and the fifth picture plot the energy density of the $D3$ -brane and of the *dilaton*- D -string systems, respectively.

instantons (S-branes). Their energy-density and charge are localized functions of time. Alternatively we see that these branes contribute rather sharp bumps in the eigenvalues of the spatial part of the stress-energy tensor which we have named *pressures*. Typically there are *maxima* of these pressures in the space directions parallel to the euclidean brane world-volume and *minima* of the same in the directions transverse to the brane. These maxima and minima in the pressures correspond to maxima and minima of the scale factors in the same directions. Such inversions in the rate of expansion/contraction of the scale factors is the *cosmological billiard* phenomenon originally envisaged by Damour et al. In the toy A_2 model we have presented, we observe just one scattering, but this is due to the insufficient number of branes (roots in the Lie algebra language) that we have excited. Indeed it is like we had only one wall of a Weyl chamber. In subsequent publications we plan to study the phenomenon in more




Figure 8.14: Plots of the scale factors $\bar{r}_{[\alpha]}^2$, $\alpha = 1|2, 3|4, 5|6|7, 8|9$ as functions of the cosmic time $t = \tau(T)$ with the critical choice of parameters $\omega = 1$, $\kappa = 3/2$ and for the A_2 solution with all the roots switched on.

complex situations with more algebraic roots switched on. What is relevant in our opinion is that we were able to see the postulated *bumping phenomenon* in the context of exact smooth solutions rather than in asymptotic limiting regimes.

8.3 Conclusions and perspectives

In [49] we developed a convenient mathematical framework within three dimensional (ungauged) maximal supergravity where to study homogeneous cosmological solutions of type II A or II B theories. Our approach exploited the correspondence between homogeneous time-dependent solutions in ten and three dimensions. This mapping was realized through *toroidal dimensional reduction* from $D = 10$ to $D = 3$ or through *oxidation* from $D = 3$ to $D = 10$. The starting point of our study was the $E_{8(8)}$ orbit described by three dimensional homogeneous solutions and we defined the precise method for constructing a generic representative of the orbit from the generating solution which is defined only by the radii of the internal seven-torus and the dilaton. Exploiting the solvable Lie algebra (or Iwasawa) representation of the scalar manifold in the three dimensional theory it was possible to control the ten dimensional interpretation of the various bosonic fields. This allows for instance to construct a ten dimensional solution characterized by certain (off-diagonal) components of the metric or of the tensor fields by oxidizing a three dimensional solution in which the scalar fields associated with the corresponding $E_{8(8)}$ roots are switched on. As an example we worked out in three dimensions the general

homogeneous time-dependent solution of an A_2 model in which the scalar fields span a $SL(3, \mathbb{R})/SO(3)$ submanifold of $E_{8(8)}/SO(16)$. It was shown that, depending on the embedding of $SL(3, \mathbb{R})$ within $E_{8(8)}$ the ten dimensional solution obtained upon oxidation of the three dimensional one can have radically different physical interpretations in terms of ten dimensional fields. We then focused on one particular embedding for which the axionic fields are interpreted as the components B_{34} , C_{89} and C_{3489} of the type II B tensor fields, and accomplished the oxidation of the solution to ten dimensions. Its behavior, which has been described in detail in the previous section, is characterized by an exchange of energy between the tensor fields and the gravitational field which results in consecutive phases of expansion and contraction of the cosmological scale factors along the directions defined by the non vanishing components of the tensor fields. This background could be interpreted microscopically in terms of a system of space-like or *S-branes* (or *SD-branes*) [10] along the directions 89 and 3489, coupled to the Kalb–Ramond field. It is interesting to make contact with the *cosmological billiard* phenomenon describing the behavior of solutions to Einstein equations near space-like singularities. In this limit the evolution of the scale-factors/dilaton is described by a null trajectory in a hyperbolic space which is reflected by *walls* or hyper-surfaces where the energy density of the axionic fields diverges. Although ours is a different kind of analysis which aims at the construction of exact smooth cosmological solutions, we may retrieve a similar *qualitative* description of the evolution of the scale-factors/dilaton in relation to the evolution of the axionic fields. In our formalism the logarithm σ_i of the scale factors associated with the internal directions together with the ten dimensional dilaton ϕ are described by the vector $h(t)$ in the Euclidean eight-dimensional space of the $E_{8(8)}$ Cartan subalgebra. The kinetic term of an axion χ associated with the positive $E_{8(8)}$ root α contains the characteristic exponential factor $\exp(-2\alpha \cdot h)$. The corresponding wall in the space of h is defined by the equation $\alpha \cdot h = 0$ and the *billiard* region by $\alpha \cdot h \geq 0$. As it can be inferred from our solution, the evolution of $h(t)$ is such that, if we denote by $h^\parallel$ the component of h along α and by $h^\perp$ its projection on the hyperplane perpendicular to α (the wall), as the energy density of χ reaches its maximum (this temporal region corresponds to the *thickness* of the S-branes in the A_2 solution) h undergoes a reflection. This is most easily illustrated for example in the A_2 solution with just one root switched on (namely $\alpha[80]$). In this case the component of $\dot{h}$ parallel to $\alpha[80]$ undergoes a continuous sign inversion from negative values to positive ones while $|\dot{h}^\perp|$ is constant and proportional to the time derivative of the dilaton :

$$\dot{h}^\parallel \rightarrow -\dot{h}^\parallel ; \quad |\dot{h}^\perp| = \frac{k}{2\sqrt{6}} \propto \dot{\phi} = \text{const.} \quad (8.54)$$

The roots in the A_2 system are not enough to define a finite volume billiard which would result in an oscillatory behavior of the solution. Indeed, using for h the parametrization in terms of the variables x, y , namely $h = \{x, x, y, y, y, -y, -y, y\}$, the billiard region is defined by the dominant walls $\alpha[69] \cdot h = 0$, $\alpha[15] \cdot h = 0$:

$$x \geq 0; \quad y \leq -\frac{x}{3} \quad (8.55)$$

which is an open region. This explains the Kasner-like (non-oscillatory) behavior of our solution for $t \rightarrow +\infty$.

More general ten dimensional homogeneous solutions deriving from the coupling of gravity with purely metric 10 dimensional backgrounds were analyzed by some of the authors of [49] in [118], in which they addressed the related question to define how many independent orbits of cosmological solutions there are under the action of the U -duality group.

In particular, exploiting first the classification of all the possible regular embeddings $\mathbf{G}_r \hookrightarrow E_8, r \leq 8$ subalgebras, then the classification of the Weyl orbits of the $\mathbf{G}_r$ root system within the E_8 root system, the authors proved that all regular embeddings of each A_r subalgebras inside $E_{8(8)}$ fall into a single Weyl orbit, with the exception of A_7 , which falls into two distinct one.

They chose a purely metric canonical representative embedding of the A_2 sigma model, then oxidized the three-dimensional solutions found [49] and eventually studied their geometrical and physical properties. The simplest solution, which has only the highest of the three roots associated to nilpotents fields switched on and the fundamental parameter κ set to zero, provides an exact example of Bianchi type 2A metric in four dimensions. Moreover, the solution with $\kappa \neq 0$ leads to a nontrivial evolution of the scale factor in the fifth dimension which, after a Kaluza Klein reduction can be reinterpreted as a Bianchi type 2A metric with scalar matter content. The same holds for the solutions with all the roots switched on: they produce further examples of type 2A Bianchi cosmologies with scalar and vector matter contents.

All these solutions are homogeneous but not isotropic, and they show the peculiar cosmological billiard feature.

A concrete step towards the understanding of billiard dynamics has been done in [119]. The main investigation in this sense would involve derivation of exact solutions directly in a $D = 2$ or $D = 1$ context where Kač–Moody symmetries become manifest. Although the appearance of Kač–Moody extensions is algebraically well established, their exploitation in deriving solutions is not as clear as the exploitation of ordinary symmetries. In [119], the authors try to clarify the field theoretical realization of the Kač–Moody extensions, this being the prerequisite for their utilization in deriving billiard dynamics. In particular they have shown that there is a general mechanism underlying the affine Kač–Moody extension of the $D=3$ algebra $\mathbb{U}_{D=3}$ when stepping down to $D = 2$ and

that this mechanism follows a general algebraic pattern for all supergravity theories, independently of the number of supercharges N_Q . This mechanism relies on the existence of two different reduction schemes from $D = 4$ to $D = 2$, respectively named the Ehlers reduction and the Matzner–Missner reduction, which are non locally related to each other. Nicolai observed this phenomenon time ago in the case of pure gravity (or better of N=1 pure supergravity)[120] and showed that one obtains two identical lagrangians, each displaying an $SL(2, \mathbb{R})$ symmetry. The fields appearing in one lagrangian have a non local relation to those of the other lagrangian and one can put together both $SL(2, \mathbb{R})$ algebras. One algebra generates local transformations on one set of fields the other algebra generates non local ones. Together the six generators of the two $SL(2, \mathbb{R})$ provide a Chevalley basis for the Kač–Moody extension $SL(2, \mathbb{R})^\wedge$ namely for $A_1^\wedge$. The analysis in [119] is an extension of the argument by Nicolai. For a generic supergravity theory, the two reduction schemes Ehlers and Matzner–Missner lead to two different lagrangians with different local symmetries. The first is a normal σ -model the second is a *twisted* σ -model. The authors discuss in detail the symmetries of both theories, thus writing down a precise field theoretic realization of the affine symmetries setting the basis to exploit them in billiard dynamics.

Part III

APPENDICES

Appendix A

Projection of generalized Langland's boundary states

This appendix contains a careful computation of the projection of generalized Langlands boundary states (2.58) and (2.59) on the constraints in equation (2.62). Dealing with the compactified boson, the only chiral fields involved besides Virasoro fields are the holomorphic and antiholomorphic currents $J(\zeta)$ $\bar{J}(\bar{\zeta})$ generating the Heisenberg algebra (2.35). Being the current algebra abelian, the possible gluing maps reduces to:

$$J(\zeta) = \bar{J}(\bar{\zeta}) \quad (\text{A.1a})$$

$$J(\zeta) = -\bar{J}(\bar{\zeta}) \quad (\text{A.1b})$$

which, through radial quantization, are respectively mapped into

$$(\mathbf{a}_n + \bar{\mathbf{a}}_{-n}) \|B\rangle\rangle = 0, \quad (\text{A.2a})$$

$$(\mathbf{a}_n - \bar{\mathbf{a}}_{-n}) \|B\rangle\rangle = 0. \quad (\text{A.2b})$$

The Sugawara construction (2.6) ensures that this conditions are sufficient to enforce conformal invariance encoded in (2.14).

To work out the projection of (A.2) over Generalized Langlands boundary states, It is convenient to introduce, for each given $n \geq 1$, the auxiliary states

$$\left| \mathbf{r}_{(\mu, \nu)}^\alpha (S_{\varepsilon(k)}^{(-)}) \right\rangle_n \doteq \sum_{m_1, m_2} \mathbb{A}_{m_1, m_2}^n (a_n^\alpha, a_{-n}^\alpha) \frac{(\mathbf{a}_{-n}^\alpha)^{m_1} (\bar{\mathbf{a}}_{-n}^\alpha)^{m_2}}{\sqrt{n^{m_1+m_2} m_1! m_2!}} |(\mu^\alpha, \nu^\alpha)\rangle, \quad (\text{A.3})$$

(one defines similarly the states $\left| \mathbf{l}_{(\mu, \nu)}^\alpha (S_{\varepsilon(k)}^{(+)}) \right\rangle_n$ associated with $\left| \mathbf{r}_{(\mu, \nu)}^\alpha (S_{\varepsilon(k)}^{(+)}) \right\rangle$).

From condition (A.2b) (for $n = 0$) we get

$$\lambda_{(\mu, \nu)}^\alpha - \bar{\lambda}_{(\mu, \nu)}^\alpha = 0. \quad (\text{A.4})$$

which selects, at a generic value of the ratio $\frac{L(k)}{R^\alpha(k)}$, the highest weight state

$$|(\mu^\alpha, 0)\rangle = \left| \lambda_{(\mu,0)}^\alpha \right\rangle \otimes \left| \bar{\lambda}_{(\mu,0)}^\alpha \right\rangle \quad (\text{A.5})$$

i.e. the winding number does not contribute to the left and right momenta associated with this Verma module

$$\begin{aligned} \mathfrak{a}_0^\alpha |(\mu^\alpha, 0)\rangle &= \mu^\alpha \frac{L}{R^\alpha} |(\mu^\alpha, 0)\rangle \\ \bar{\mathfrak{a}}_0^\alpha |(\mu^\alpha, 0)\rangle &= \mu^\alpha \frac{L}{R^\alpha} |(\mu^\alpha, 0)\rangle \end{aligned} \quad \mu^\alpha \in \mathbb{Z} . \quad (\text{A.6})$$

Similarly, the constraint (A.2a) provides the condition

$$\lambda_{(\mu,\nu)}^\alpha + \bar{\lambda}_{(\mu,\nu)}^\alpha = 0. \quad (\text{A.7})$$

which selects, at generic value of the ratio $\frac{L(k)}{R^\alpha(k)}$, the highest weight state:

$$|(0, \nu^\alpha)\rangle = \left| \lambda_{(0,\nu)}^\alpha \right\rangle \otimes \left| \bar{\lambda}_{(0,\nu)}^\alpha \right\rangle \quad (\text{A.8})$$

such that

$$\begin{aligned} \mathfrak{a}_0^\alpha |(0, \nu^\alpha)\rangle &= \frac{1}{2} \nu^\alpha \frac{R^\alpha}{L} |(0, \nu^\alpha)\rangle \\ \bar{\mathfrak{a}}_0^\alpha |(0, \nu^\alpha)\rangle &= -\frac{1}{2} \nu^\alpha \frac{R^\alpha}{L} |(0, \nu^\alpha)\rangle \end{aligned} \quad \nu^\alpha \in \mathbb{Z} . \quad (\text{A.9})$$

Before considering the conditions (A.2) for $n \neq 0$, let us impose the condition projecting the $\left| \mathfrak{r}_{(\mu,\nu)}^\alpha(S_{\varepsilon(k)}^{(-)}) \right\rangle$ into an actual physical state:

$$\left(\mathbb{L}_0^\alpha - \bar{\mathbb{L}}_0^\alpha \right) \left| \mathfrak{r}_{(\mu,\nu)}^\alpha(S_{\varepsilon(k)}^{(-)}) \right\rangle = 0. \quad (\text{A.10})$$

From the commutation relation

$$[\mathfrak{a}_j^\alpha, (\mathfrak{a}_{-n}^\alpha)^{m_1}] = m_1 j \delta_{j-n,0} (\mathfrak{a}_{-n}^\alpha)^{m_1-1}, \quad (\text{A.11})$$

one computes

$$\left(\mathbb{L}_0^\alpha - \bar{\mathbb{L}}_0^\alpha \right) \left| \mathfrak{r}_{(\mu,\nu)}^\alpha(S_{\varepsilon(k)}^{(-)}) \right\rangle = \mu^\alpha \nu^\alpha \left| \mathfrak{r}_{(\mu,\nu)}^\alpha(S_{\varepsilon(k)}^{(-)}) \right\rangle + e^{\sqrt{-1}t_+^\alpha(\lambda_{(\mu,\nu)}^\alpha + \bar{\lambda}_{(\mu,\nu)}^\alpha)}. \quad (\text{A.12})$$

$$\cdot \prod_{n=1}^{\infty} \sum_{m_1, m_2} \mathbb{A}_{m_1, m_2}^n n(m_1 - m_2) \frac{(\mathfrak{a}_{-n}^\alpha)^{m_1} (\mathfrak{a}_{-n}^\alpha)^{m_2}}{\sqrt{n^{m_1+m_2} m_1! m_2!}} |(\mu^\alpha, \nu^\alpha)\rangle ,$$

which, together with the $U(1)$ constraints, (implying that $\mu^\alpha \nu^\alpha = 0$), shows that the physical states are characterized by the further requirement

$$m_1 = m_2.$$

For $j \neq 0$, the action of the operators $\mathfrak{a}_j^\alpha$ on the boundary states $\left| \mathfrak{r}_{(\mu,\nu)}^\alpha(S_{\varepsilon(k)}^{(\pm)}) \right\rangle$ is non-trivial when $j > 0$, and it can be worked out by considering the auxiliary

states $\left| l_{(\mu,\nu)}^{\alpha}(S_{\varepsilon(k)}^{(\pm)}) \right\rangle_n$ for each value of $n \geq 1$. In detail, let us consider the state $\left| l_{(\mu,\nu)}^{\alpha}(S_{\varepsilon(k)}^{(-)}) \right\rangle_n$, (with $m_1 = m_2 = m \geq 1$; the action of the annihilator $\mathfrak{a}_j^{\alpha}$ being trivial on $m = 0$),

$$\begin{aligned} \mathfrak{a}_j^{\alpha} \left| l_{(\mu,\nu)}^{\alpha}(S_{\varepsilon(k)}^{(-)}) \right\rangle_n &= \sum_{m=1}^{\infty} \mathbb{A}_m^j \frac{(mj) (\mathfrak{a}_{-j}^{\alpha})^{m-1} (\bar{\mathfrak{a}}_{-j}^{\alpha})^m}{j^m m!} |(\mu^{\alpha}, \nu^{\alpha})\rangle \\ &= \bar{\mathfrak{a}}_{-j}^{\alpha} \sum_{m=1}^{\infty} \mathbb{A}_m^j \frac{(\mathfrak{a}_{-j}^{\alpha})^{m-1} (\bar{\mathfrak{a}}_{-j}^{\alpha})^{m-1}}{j^{m-1} (m-1)!} |(\mu^{\alpha}, \nu^{\alpha})\rangle. \end{aligned} \quad (\text{A.13})$$

From the definition of $\mathbb{A}_m^j$, and the following recursion relation for Laguerre polynomials

$$L_m^{-1}(x) = L_m(x) - L_{m-1}(x), \quad (\text{A.14})$$

we get

$$\mathbb{A}_m^j = e^{2\pi s \sqrt{-1}} \mathbb{A}_{m-1}^j + e^{-2j|a_j^{\alpha}|^2} e^{2\pi s m \sqrt{-1}} L_m^{-1}(4j|a_j^{\alpha}|^2). \quad (\text{A.15})$$

Inserting this latter into (A.13) provides

$$\begin{aligned} \mathfrak{a}_j^{\alpha} \left| l_{(\mu,\nu)}^{\alpha}(S_{\varepsilon(k)}^{(-)}) \right\rangle_n &= e^{2\pi s \sqrt{-1}} \bar{\mathfrak{a}}_{-j}^{\alpha} \left| l_{(\mu,\nu)}^{\alpha}(S_{\varepsilon(k)}^{(-)}) \right\rangle_n + \\ &+ e^{-2j|a_j^{\alpha}|^2} e^{2\pi s m \sqrt{-1}} \bar{\mathfrak{a}}_{-j}^{\alpha} \sum_m L_m^{-1}(4j|a_j^{\alpha}|^2) \frac{(\mathfrak{a}_{-j}^{\alpha})^{m-1} (\bar{\mathfrak{a}}_{-j}^{\alpha})^{m-1}}{j^{m-1} (m-1)!} |(\mu^{\alpha}, \nu^{\alpha})\rangle, \end{aligned}$$

from which it immediately follows that the Neumann condition

$$(\mathfrak{a}_j^{\alpha} - \bar{\mathfrak{a}}_{-j}^{\alpha}) \left| l_{(\mu,\nu)}^{\alpha}(S_{\varepsilon(k)}^{(-)}) \right\rangle_n^{(N)} = 0 \quad (\text{A.16})$$

requires $s \in \mathbb{Z}$ and $L_m^{-1}(4j|a_j^{\alpha}|^2) = 0$, viz. $a_j^{\alpha} \equiv 0 \ \forall j$, whereas for the Dirichlet condition

$$(\mathfrak{a}_j^{\alpha} + \bar{\mathfrak{a}}_{-j}^{\alpha}) \left| l_{(\mu,\nu)}^{\alpha}(S_{\varepsilon(k)}^{(-)}) \right\rangle_n^{(D)} = 0 \quad (\text{A.17})$$

we need $s \in \frac{1}{2}\mathbb{Z}$ and $a_j^{\alpha} \equiv 0 \ \forall j$. Since $\mathbb{A}_m^n(a_n^{\alpha} = 0) = e^{2\pi m s \sqrt{-1}}$, and

$$\sum_m \mathbb{A}_m^n \frac{(\mathfrak{a}_{-n}^{\alpha})^m (\bar{\mathfrak{a}}_{-n}^{\alpha})^m}{n^m m!} = \begin{cases} \exp\left(\frac{1}{n} (\mathfrak{a}_{-n}^{\alpha}) (\bar{\mathfrak{a}}_{-n}^{\alpha})\right), & s \in \mathbb{Z} \\ \exp\left(-\frac{1}{n} (\mathfrak{a}_{-n}^{\alpha}) (\bar{\mathfrak{a}}_{-n}^{\alpha})\right), & s \in \frac{1}{2}\mathbb{Z} \end{cases}, \quad (\text{A.18})$$

the corresponding boundary states are provided by

$$\left| \mathfrak{t}_{\mu}^{\alpha}(S_{\varepsilon(k)}^{(-)}) \right\rangle^{(D)} = e^{\sqrt{-1} t_+^{\alpha} \mu^{\alpha} \frac{t_-}{R^{\alpha}}} \exp\left(\sum_{n=1}^{\infty} \frac{1}{n} (\mathfrak{a}_{-n}^{\alpha}) (\bar{\mathfrak{a}}_{-n}^{\alpha})\right) |(\mu^{\alpha}, 0)\rangle, \quad (\text{A.19})$$

and

$$\left| \mathbf{r}^\alpha(S_{\varepsilon(k)}^{(-)}) \right\rangle^{(N)} = \exp \left(\sum_{n=1}^{\infty} -\frac{1}{n} (\mathbf{a}_{-n}^\alpha) (\bar{\mathbf{a}}_{-n}^\alpha) \right) |(0, \nu^\alpha)\rangle. \quad (\text{A.20})$$

Note that (up to the weighting factor in (A.20)) these states are nothing but the Dirichlet and Neumann Ishibashi states associated with the free bosonic field $X^\alpha(k)|_-$ on the circle $S_{\varepsilon(k)}^{(-)}$.

Appendix B

Conformal properties of Boundary Insertion Operators

Boundary Insertion Operators live on the ribbon graph, thus their interactions are guided by the trivalent structure of Γ . In this appendix we report an exhaustive analysis which shows that this structure allows to define all fundamental coefficients weighting the self-interaction of boundary insertion operators. Two points functions are well defined on the edges of the graph, while OPE coefficients naturally defines the fusion of different boundary insertion operators interacting in $N_2(T)$ tri-valent vertexes of Γ . Moreover, we can define a set of sewing constraint these coefficients must satisfy. Remarkably, these constraints are perfectly analogues to the sewing constraint introduced in [73] for boundary conditions changing operator, thus enforcing the analogy between these latters and BIOS.

Due to their CVO analogue structure, we are able to specify the conformal properties of these operators. BIOS are primary operators of the boundary chiral algebra, thus they have a well-defined conformal dimension which, in this case, coincide with the highest weight of the $V_\lambda(p, q)$ module of the $U(1)$ (Virasoro) algebra:

$$H(p, q) = \frac{1}{2} \lambda(p, q)^2 \quad (\text{B.1})$$

As usual, the conformal invariance fixes to zero the one point function:

$$\langle \psi_{\lambda(p, q)}^{B(q)A(p)} \rangle = 0 \quad (\text{B.2})$$

while, for the two-points function, it is a priori possible to construct two types of correlators between BIOS which mediate the changing in boundary conditions between the two adjacent cylinder $\Delta_{\varepsilon(p)}^*$ and $\Delta_{\varepsilon(q)}^*$: a first one will correlate two operators which both mediate the changing in the “ p -to- q ” direction (or, equivalently, which mediate both the changing in the “ q -to- p ” direction). The




Figure B.1: Two-point functions

second type will correlate two BIOs which mediate one in the “ p -to- q ” direction and the other in the “ q -to- p ” one (see figure B.1).

The former situation, depicted in figure B.1a, leads to:

$$\langle \psi_{\lambda(p,q)}^{B(q)A(p)}(x_1(p,q)) \psi_{\lambda'(q,p)}^{C(p)D(q)}(x_2(q,p)) \rangle = \frac{b_{\lambda(p,q)}^{B(q)A(p)} \delta_{\lambda(p,q)\lambda'(p,q)} \delta^{A(p)C(p)} \delta^{B(q)D(q)}}{|x_1(p,q) - x_2(p,q)|^{2H(p,q)}} \quad (\text{B.3})$$

Note that the Kronecker delta function is used to let the two boundary conditions on the same oriented edge of the strip to be compatible. In the usual BCFT boundary fields correlators, the two fields are taken to perform two serial switch in boundary condition, and this leads to a different equalities between boundary conditions indexes. Here, however, the two operators act in a parallel way, and this justifies our choice for delta functions.

Obviously, the second situation, depicted in figure B.1b, leads to the same result because of transformation law (3.11):

$$\langle \psi_{\lambda(p,q)}^{B(q)A(p)}(x_1(p,q)) \psi_{\lambda'(q,p)}^{C(p)D(q)}(x_2(q,p)) \rangle = \frac{b_{\lambda(p,q)}^{B(q)A(p)} \delta_{\lambda(p,q)\lambda'(p,q)} \delta^{A(p)C(p)} \delta^{B(q)D(q)}}{|x_1(p,q) - x_2(p,q)|^{2H(p,q)}} \quad (\text{B.4})$$

To analyze the conformal properties of BIOs and define completely the equivalence between them and usual boundary operators in CFT, let us consider what happens in each vertex of the underling triangulation (the situation is the one depicted in figure B.2). Let us consider the generic vertex of the triangulation $\rho^0(p, q, r)$ and its uniformizing neighborhood unit disk $(U_{\rho^0(p,q,r)}, \omega(p, q, r))$. The BIOs are well defined on the disk because of the transition functions defined in (1.30). The vertexes of the triangulation are the natural point in which to define the fusion between BIOs coming from different edges via their OPE.

Let us take three points in an ϵ -neighborhood of the vertex $\omega = 0$, denoting




Figure B.2: Two-point functions

with

$$\omega_r = \frac{\epsilon}{3} e^{\frac{\pi}{6}} \in U_{\rho^0(p,q,r)} \cap U_{\rho^1(r,p)} \quad (\text{B.5})$$

$$\omega_p = \frac{\epsilon}{2} e^{\frac{5\pi}{6}} \in U_{\rho^0(p,q,r)} \cap U_{\rho^1(p,q)} \quad (\text{B.6})$$

$$\omega_q = \epsilon e^{\frac{3\pi}{2}} \in U_{\rho^0(p,q,r)} \cap U_{\rho^1(q,r)} \quad (\text{B.7})$$

their coordinates, and let us focus our attention the three fields $\psi_{\lambda(r,p)}^{A(p)C(r)}(\omega_r)$, $\psi_{\lambda(p,q)}^{B(q)A(p)}(\omega_p)$ and $\psi_{\lambda(q,r)}^{C(r)B(q)}(\omega_q)$ which mediate pairwise the boundary conditions between the boundaries of $\rho^2(r)$ and $\rho^2(p)$, $\rho^2(p)$ and $\rho^2(q)$ and $\rho^2(q)$ and $\rho^2(r)$ respectively (as usual, the direction of the action of BIOs is given by the notation).

In the limit $\epsilon \rightarrow 0$ the product of the two fields $\psi_{\lambda(r,p)}^{A(p)C(r)}$ and $\psi_{\lambda(q,r)}^{C(r)B(q)}$ will mediate the changing in boundary conditions from $B(q)$ to $A(p)$, thus the OPE of these two fields must be expressed as a function of a $\psi_{\lambda(q,p)}^{A(p)B(q)}$ -type field:

$$\begin{aligned} \psi_{\lambda(r,p)}^{A(p)C(r)}(\omega_r) \psi_{\lambda'(q,r)}^{C(r)B(q)}(\omega_q) \sim \\ \sum_{\lambda''(q,p) \in \mathcal{Y}} \mathcal{C}_{\lambda(r,p)\lambda'(q,r)\lambda''(q,p)}^{A(p)C(r)B(q)} |\omega_r - \omega_q|^{H(q,p) - H(r,p) - H(q,r)} \psi_{\lambda''(q,p)}^{A(p)B(q)}(\omega_q) \end{aligned} \quad (\text{B.8a})$$

This situation is described by the continuous arrows in fig. B.2

In the same way we can write the other two naturally defined OPES:

$$\begin{aligned} \psi_{\lambda(q,r)}^{C(r)B(q)}(\omega_q) \psi_{\lambda'(p,q)}^{B(q)A(p)}(\omega_p) \sim \\ \sum_{\lambda''(p,r) \in \mathcal{Y}} \mathcal{C}_{\lambda(q,r)\lambda'(p,q)\lambda''(p,r)}^{C(r)B(q)A(p)} |\omega_r - \omega_q|^{H(q,p) - H(r,p) - H(q,r)} \psi_{\lambda''(p,r)}^{C(r)A(p)}(\omega_p) \end{aligned} \quad (\text{B.8b})$$

described by the dashed arrows in fig. B.2, and

$$\begin{aligned} \psi_{\lambda(p,q)}^{B(q)A(p)}(\omega_p) \psi_{\lambda'(r,p)}^{A(p)C(r)}(\omega_r) \sim \\ \sum_{\lambda''(q,p) \in \mathcal{Y}} \mathcal{C}_{\lambda(p,q)\lambda'(r,p)\lambda''(q,p)}^{B(q)A(p)C(r)} |\omega_r - \omega_q|^{H(q,p) - H(r,p) - H(q,r)} \psi_{\lambda''(q,p)}^{B(q)C(r)}(\omega_r) \end{aligned} \quad (\text{B.8c})$$

described by the dotted arrows in fig. B.2

The complete description of the interaction of the N_0 BCFTs passes through the determination of the OPE coefficients of the BIOs $\mathcal{C}_{\lambda(\cdot, \cdot)\lambda'(\cdot, \cdot)\lambda''(\cdot, \cdot)}^{A(\cdot)B(\cdot)C(\cdot)}$, defined for $A, B, C \in \mathcal{A}$ and $\lambda, \lambda', \lambda'' \in \mathcal{Y}$.

To obtain this result, it is first of all necessary to prove that our description is totally equivalent with the usual BCFT formulation. This can be achieved via demonstrating that the OPE coefficients and the normalization factor entering in (B.3) satisfy both the cyclic symmetry and the sewing constraint which derive from the three points and four points functions in the usual formulation of a Boundary Conformal Field Theory. Let us start from considering the three points function which arise naturally in the vertex $\rho^0(p, q, r)$, which must be cyclically invariant. Thus:

$$\begin{aligned} \langle \psi_{\lambda(r,p)}^{A(p)C(r)}(\omega_r) \psi_{\lambda'(q,r)}^{C(r)B(q)}(\omega_q) \psi_{\lambda''(p,q)}^{B(q)A(p)}(\omega_p) \rangle = \\ \langle \psi_{\lambda''(p,q)}^{B(q)A(p)}(\omega_p) \psi_{\lambda(r,p)}^{A(p)C(r)}(\omega_r) \psi_{\lambda'(q,r)}^{C(r)B(q)}(\omega_q) \rangle = \\ \langle \psi_{\lambda'(q,r)}^{C(r)B(q)}(\omega_q) \psi_{\lambda''(p,q)}^{B(q)A(p)}(\omega_p) \psi_{\lambda(r,p)}^{A(p)C(r)}(\omega_r) \rangle \quad (\text{B.9}) \end{aligned}$$

If we exploit the OPEs defined in (B.8), from each identity in the previous equation we obtain the following relations between the OPEs coefficients:

$$\mathcal{C}_{\lambda(r,p)\lambda'(q,r)\lambda''(p,q)}^{A(p)C(r)B(q)} d_{\lambda''(q,p)}^{A(p)B(q)} = \mathcal{C}_{\lambda''(p,q)\lambda(r,p)\lambda'(r,q)}^{B(q)A(p)C(r)} d_{\lambda'(r,q)}^{B(q)C(r)} \quad (\text{B.10})$$

$$\mathcal{C}_{\lambda(r,p)\lambda'(q,r)\lambda''(p,q)}^{A(p)C(r)B(q)} d_{\lambda''(q,p)}^{A(p)B(q)} = \mathcal{C}_{\lambda'(q,r)\lambda''(p,q)\lambda(p,r)}^{C(r)B(q)A(p)} d_{\lambda(p,r)}^{C(r)A(p)} \quad (\text{B.11})$$

$$\mathcal{C}_{\lambda''(p,q)\lambda(r,p)\lambda'(r,q)}^{B(q)A(p)C(r)} d_{\lambda'(r,q)}^{B(q)C(r)} = \mathcal{C}_{\lambda'(q,r)\lambda''(p,q)\lambda(p,r)}^{C(r)B(q)A(p)} d_{\lambda(p,r)}^{C(r)A(p)} \quad (\text{B.12})$$

where the $d_{\lambda''(q,p)}^{A(p)B(q)}$ are the normalization coefficients of the two point functions in the $(U_{\rho^0(p,q,r)}, \omega(p, q, r))$ frame:

$$d_{\lambda(q,p)}^{A(p)B(q)} = \left| \frac{d\omega}{dz(q,p)} \right|_{\omega=\omega_q}^{-H(q,p)} \left| \frac{d\omega}{dz(q,p)} \right|_{\omega=\omega_p}^{-H(q,p)} b_{\lambda(q,p)}^{A(p)B(q)}, \quad (\text{B.13})$$

which satisfies the obvious property $d_{\lambda(q,p)}^{A(p)B(q)} = d_{\lambda(p,q)}^{B(q)A(p)}$.

Moreover, if we take the two different OPEs in the first term of (B.9), we obtain the usual sewing constraint:

$$\mathcal{C}_{\lambda(r,p)\lambda'(q,r)\lambda''(p,q)}^{A(p)C(r)B(q)} \mathcal{C}_{\lambda''(q,p)\lambda''(p,q)0}^{A(p)B(q)A(p)} = \mathcal{C}_{\lambda'(q,r)\lambda''(p,q)\lambda(p,r)}^{C(r)B(q)A(p)} \mathcal{C}_{\lambda(r,p)\lambda(p,r)0}^{A(p)C(r)A(p)} \quad (\text{B.14})$$

where 0 is the singlet label. To complete this analysis, we would need to check sewing constraints arising from the four-points function behavior. We leave this further investigation to the next chapter, in which will see that the behavior of the four points function is fundamental to define the action of BIOs on the full surface. This analysis, together with properties outlined in section proves the complete equivalence between the usual action of boundary changing conditions operators on the upper half plane (UHP) boundary, and our particular model, in which BIOs merge dynamically the N_0 copies of an UHP-defined BCFT.

Appendix C

An introduction to truly marginal deformations of boundary conformal field theories

The coupling of a String theory with a generic background field corresponds in a microscopic CFT description to a *deformation* of the model. This deformation is formally obtained modifying the action with a perturbative term:

$$S \rightarrow S' = S + g \int d^2z \mathcal{O}(z, \bar{z}) \quad (\text{C.1})$$

For the theory to remain conformally invariant, the operator must be a *marginal operator*, i.e. $\mathcal{O}(z, \bar{z})$ must at least have conformal dimension (1,1) in order to let the integrated quantity be dimensionless.

Let us assume one is given a certain Lagrangian $\mathcal{L}$ defining an exactly solvable Conformal Field Theory on a two dimensional manifold Σ (thus we know the central charge of the Virasoro algebra, the spectrum of the model and the associated Vertex algebra). A deformation of this model is obtained by adding a perturbative term to the action:

$$\mathcal{L} \rightarrow \mathcal{L}' = \mathcal{L} + \sum_i g_i f_i(z, \bar{z}) \quad (\text{C.2})$$

where $f_i(z, \bar{z})$ are some operators in the spectrum of the theory defined by L , while g_i are some coupling constants. In this connection, a Conformal Field Theory can be defined as a fixed point in the infinite dimensional space of theories parametrized by the coefficients of possible operators such as in equation (C.2).

Let us consider the simpler case in which we add to $\mathcal{L}$ a single operator $f(z, \bar{z})$ with coefficient g . In a Renormalization Group (RG) approach, once the operator is added with a certain value of the coefficient, the theory (*i.e.* the coefficient value) flows along a RG trajectory as the energy scale μ is changed. The behavior of the coupling is characterized by the β -function

$$\beta(g) \equiv \mu \frac{\partial g}{\partial \mu} \quad (\text{C.3})$$

One can imagine the existence of a fixed point of such flows when $\beta(g) = 0$ thus $g(\mu) = \bar{g}$ and the coupling tend to that specific value, as shown in fig. C.1.




Figure C.1: Flows

On the left, $\bar{g}$ is an IR fixed point, since the coupling tends to this value for decreasing μ . On the contrary, on the right, the theory is an UV fixed point, since the coupling constant tends to the fixed point value for increasing energy scale.

The spectrum of operators associated with $\mathcal{L}$ can be divided into the following categories:

- Operators whose conformal dimension is larger than $(1,1)$, which are called *irrelevant operators*. The correspondent coupling constant has a negative mass dimension. If one adds an irrelevant operator to a CFT exact Lagrangian, one finds that the coupling constant in the new Lagrangian decreases to zero in the IR limit, thus the name itself.
- Operators whose conformal dimension is smaller than $(1,1)$, which are called *relevant operators*. The corresponding coupling constant has a positive mass dimension and, adding such an operator to $\mathcal{L}$ one finds that in the new model, g flows to a larger value in the IR. Moreover, if the theory is unitary, it will flow to a smaller c value one.
- Operators whose dimension is exactly $(1,1)$. They are called *marginal operators*, and their associated coupling constants are dimensionless. They do not break the scale invariance of the model explicitly, thus they do not take away the CFT from the fixed point. However, the coupling constants

can change under renormalization. This leads the marginal operators to subdivide into three classes:

- Marginal operators whose couplings in the modified Lagrangian are IR free on the worldsheet. Thus these operators turn out to be irrelevant ones. They differ from operators irrelevant at the classical level because they decrease logarithmically and not power-like towards the infrared.
- Marginal operators whose couplings g_i under the modified Lagrangian are asymptotically free, thus increase logarithmically towards the infrared. They are then relevant operators.
- Marginal operators whose addition to the Lagrangian $\mathcal{L}$ maintains the couplings g_i dimensionless. This promotes $\mathcal{L}$ to a family of CFTs. These operators are called *truly marginal* and form a basis for a family for neighborhood of CFTs.

C.1 Deformations of a boundary conformal field theory

We want to describe the effects of perturbations of a boundary condition generated by marginal operators, in particular marginal boundary fields. We will show that a certain class of perturbations, which in [78] are called analytic deformations, are truly marginal at all orders in the perturbation expansion. These perturbations are induced by *self-local* boundary operators of dimension one.

Definition 3 *Mutually local boundary fields*

Let $\{\varphi_\lambda(z, \bar{z}) = \Phi(|\varphi_\lambda\rangle; z, \bar{z})\}$ and $\{\psi_\nu(x) = \Phi(|\psi_\nu\rangle; x)\}$ be respectively the collection of bulk and boundary fields of a given BCFT. Let us consider the correlator

$$\langle \psi_1(x_1) \cdots \psi_n(x_n) \psi_1(z_1, \bar{z}_1) \cdots \psi_N(z_N, \bar{z}_N) \rangle_\alpha, \quad x_\nu < x_{\nu+1}, \quad (\text{C.4})$$

where α denotes the given boundary condition on the real axis. These functions are analytic in the variables $z_i, i = 1, \dots, N$ throughout the whole UHP $\Im z > 0$. For the variables $x_\nu, \nu = 1, \dots, N$, the analyticity domain is restricted to the interval $x \in]x_{\nu-1}, x_{\nu+1}[$ on the boundary. In most cases, there are no unique analytic continuation of the correlation functions to exchange the position of two two neighboring boundary fields, since the result may depend on the orientation of the path we analytically continue the correlator (see fig. C.2).

In this connection, two boundary fields $\psi_{\nu_1}(x_1) = \Phi(|\psi_{\nu_1}\rangle; x_1)$ and $\psi_{\nu_2}(x_2) = \Phi(|\psi_{\nu_2}\rangle; x_2)$ are said to be **MUTUALLY LOCAL** if

$$\Phi(|\psi_{\nu_1}\rangle; x_1) \Phi(|\psi_{\nu_2}\rangle; x_2) = \Phi(|\psi_{\nu_2}\rangle; x_2) \Phi(|\psi_{\nu_1}\rangle; x_1), \quad x_1 < x_2 \quad (\text{C.5})$$




Figure C.2: Curves along which correlators of bulk and boundary operators can be analytically continued

where the last relation holds if included into an arbitrary correlator function of bulk and boundary fields. Obviously, to make sense it requires that an unique analytic continuation from $x_1 < x_2$ to $x_2 < x_1$ does exist.

Definition 4 Self-local boundary fields

A boundary field $\psi_\nu(x)$ is called SELF-LOCAL or ANALYTIC if it is mutually local with respect to itself.

Let us notice that definition 4 constraints the OPE of a self-local boundary field to contain only pole singularities. In particular, the OPE of a self-local boundary field $\psi(x)$ with $h_\psi = 1$, is:

$$\psi(x_1) \psi(x_2) \sim \frac{K}{(x_1 - x_2)^2} \quad (\text{C.6})$$

where K is a real constant.

C.1.1 The general prescription

To describe truly marginal perturbations of a CFT in a general contest, let us consider a Boundary Conformal Field Theory defined on the Upper Half plane $\Im z \geq 0$. Its state space will be labelled by a couple of parameters (Ω, α) where Ω identifies the gluing map along the real line, while α is the given boundary condition. We can use boundary operators $\psi(x) \in \Psi(\mathcal{H})$ to define a new perturbed theory. According to the general formula (C.2), the deformed model is obtained by adding to the action the operator:

$$S' = g \int_{-\infty}^{+\infty} \frac{dx}{2\pi} \psi(x) \quad (\text{C.7})$$

Thus, perturbed correlators are obtained by the unperturbed ones by the formal prescription:

$$\begin{aligned}
\langle \varphi_1(z_1, \bar{z}_1) \cdots \varphi_N(z_N, \bar{z}_N) \rangle_{\alpha; \lambda_\varphi} &= Z^{-1} \cdot \langle I_{\lambda_\varphi} \varphi_1(z_1, \bar{z}_1) \cdots \varphi_n(z_n, \bar{z}_n) \rangle_\alpha \\
&:= Z^{-1} \sum_n g^n \int \cdots \int_{x_i < x_{i+1}} \frac{dx_1}{2\pi} \cdots \frac{dx_n}{2\pi} \langle \psi(x_1) \cdots \psi(x_n) \varphi_1(z_1, \bar{z}_1) \cdots \varphi_n(z_n, \bar{z}_n) \rangle_\alpha \\
&= Z^{-1} \sum_n \frac{g^n}{n!} \sum_{\sigma \in S_n} \int \cdots \int_{x_{\sigma(i)} < x_{\sigma(i+1)}} \frac{dx_1}{2\pi} \cdots \frac{dx_n}{2\pi} \\
&\quad \times \langle \psi(x_1) \cdots \psi(x_n) \varphi_1(z_1, \bar{z}_1) \cdots \varphi_n(z_n, \bar{z}_n) \rangle_\alpha \quad (\text{C.8})
\end{aligned}$$

where g is the real coupling constant and the second sum runs over all the elements of the permutation group S_n . The operator $I_{g\varphi}$ has to be understood as the path-ordered exponential of the perturbing term (C.7), $I_{g\varphi} = \text{P exp } S'$, and the normalization is defined as the expectation value $Z = (A_0^\alpha)^{-1} \langle I_{g\varphi} \rangle$. If other boundary fields are included in the unperturbed correlator, prescription (C.8) has to be modified in order to include all boundary fields in the path ordering.

To make sense of (C.8) beyond the formal level, integrals need to be regularized, introducing suitable IR and UV cutoffs, while coupling and field have to be renormalized. In the following, we will not encounter IR-divergencies (which are usually cured by putting the system in a finite box, *i.e.* by considering finite-temperature correlators). On the other hand, we will deal with UV-divergent integrals. Thus, let us regularize them introducing a UV-cutoff ϵ , to let integrals domains be restricted to regions $|x_i - x_j| > \epsilon$ and to let integrals became UV-finite before taking the limit $\epsilon \rightarrow 0$.

In the following, we will analyze deformations generated by truly marginal operators, *i.e.* operators with $h = 1$. At RG fix-points, all CFT deformations share the fact that local properties of the bulk theory are not affected by the boundary condensate. Thus, boundary deformations actually will only involve changes in boundary conditions.

Let us start discussing the change induced by a boundary field $\psi(x)$ on the two point function of the boundary field itself $\langle \psi(x) \psi(x) \rangle_\alpha$. First order contributions will involve the following integrals:

$$\begin{aligned}
&\int_{-\infty}^{x_1 - \epsilon} dx \langle \psi(x) \psi(x_1) \psi(x_2) \rangle_\alpha + \\
&\int_{x_1 + \epsilon}^{x_2 - \epsilon} dx \langle \psi(x_1) \psi(x) \psi(x_2) \rangle_\alpha + \int_{x_2 + \epsilon}^{\infty} dx \langle \psi(x_1) \psi(x_2) \psi(x) \rangle_\alpha \quad (\text{C.9})
\end{aligned}$$

From the general form for a three points function we get

$$\langle \psi(x_1) \psi(x_2) \psi(x_3) \rangle_\alpha = \frac{C_{\psi\psi\psi}^\alpha}{(x_1 - x_2)(x_1 - x_3)(x_2 - x_3)} \quad (\text{C.10})$$

we easily notice that the first order contribution to the perturbed two points function logarithmically diverges unless the structure constants $C_{\psi\psi\psi}^\alpha$ would vanish. This divergence would let the conformal weight h_ψ flow away from $h_\psi = 1$, thus a boundary field ψ is not truly marginal unless $C_{\psi\psi\psi}^\alpha$. Moreover, if there are more boundary fields in a BCFT, then a marginal field ψ is truly marginal only if $C_{\psi\psi\psi'}^\alpha = 0$ for all marginal boundary fields $\{\psi'\}$ in the theory. Equation (C.6), together with (3.9), shows that self-local operators satisfy this (necessary but not sufficient - as a matter of fact true marginality is not guaranteed at higher orders of the perturbative expansion) first order condition.

C.1.2 Truly marginal operators: the case of chiral fields

The discussion can be pursued further: as a matter of fact, it is possible to prove that EVERY SELF-LOCAL MARGINAL BOUNDARY OPERATOR IS INDEED TRULY MARGINAL TO EACH ORDER AND IT GENERATES A DEFORMATION OF A BCFT. To this end, let us consider a perturbing self-local marginal field $\psi(x)$ and let us rewrite the regularized deformed correlators of bulk fields as:

$$\langle \varphi_1(z_1, \bar{z}_1) \cdots \varphi_N(z_N, \bar{z}_N) \rangle_{\alpha; g\varphi}^\epsilon = Z^{-1} \sum_n \frac{g^n}{n!} \int_{-\infty}^{+\infty} \cdots \int_{-\infty}^{+\infty} \frac{dx_1}{2\pi} \cdots \frac{dx_n}{2\pi} \langle \psi(x_1) \cdots \psi(x_n) \varphi_1 \cdots \varphi_N \rangle_\alpha \quad (\text{C.11})$$

where the integrations are performed on the real axes with the UV-cutoff described above. Including (C.6) in (C.11), we can show that logarithmic divergences cancel out and the $\epsilon \rightarrow 0$ limit is well defined. Due to self-locality, the final result of the limit procedure is:

$$\langle \varphi_1(z_1, \bar{z}_1) \cdots \varphi_N(z_N, \bar{z}_N) \rangle_{\alpha; g\varphi} = \lim_{\epsilon \rightarrow 0} \langle \varphi_1(z_1, \bar{z}_1) \cdots \varphi_N(z_N, \bar{z}_N) \rangle_{\alpha; g\varphi}^\epsilon = \sum_n \frac{g^n}{n!} \int_{\gamma_1} \cdots \int_{\gamma_n} \frac{dx_1}{2\pi} \cdots \frac{dx_n}{2\pi} \langle \psi(x_1) \cdots \psi(x_n) \varphi_1 \cdots \varphi_N \rangle_\alpha, \quad (\text{C.12})$$

where the various γ_i are straight lines parallel to the real axis parametrized as $\Im z = i\epsilon_i$. The rhs of equation (C.12) is finite and independent from ϵ as long as $\epsilon < \Im z_i \forall i = 1, \dots, N$, where $\{z_i\}$ are the bulk fields insertion points. Thus, formula (C.12) allows to construct the perturbed bulk fields correlators to all orders in perturbation theory.

The extension of the above formula to mixed correlator, containing both bulk and boundary fields, is not straightforward. As a matter of fact (C.12) can

be extended to:

$$\begin{aligned} & \langle \psi_1(u_1) \cdots \psi_M(u_M) \varphi_1(z_1, \bar{z}_1) \cdots \varphi_N(z_N, \bar{z}_N) \rangle_{\alpha; g\varphi} = \\ & \lim_{\epsilon \rightarrow 0} \langle \varphi_1(z_1, \bar{z}_1) \cdots \varphi_N(z_N, \bar{z}_N) \rangle_{\alpha; g\varphi}^\epsilon = \\ & Z^{-1} \sum_n \frac{g^n}{n!} \int_{-\infty}^{+\infty} \cdots \int_{-\infty}^{+\infty} \frac{dx_1}{2\pi} \cdots \frac{dx_n}{2\pi} \langle \psi(x_1) \cdots \psi(x_n) \psi_1 \cdots \psi_M \varphi_1 \cdots \varphi_N \rangle_\alpha, \end{aligned} \quad (\text{C.13})$$

if and only if the boundary fields are local with respect to the perturbing field. Even in those cases, integrals on the rhs of (C.13) diverges when $\epsilon \rightarrow 0$ whenever the iterated OPE of the perturbing field with one of the boundary fields contains an even pole. Renormalization of such divergencies is worked out introducing renormalized boundary operators:

$$\tilde{\psi}_i = \left[e^{\frac{1}{2}g\psi} \psi_i \right] (u_i) \doteq \sum_{n=0}^{\infty} \frac{g^n}{2^n n!} \oint_{C_1} \frac{dx_1}{2\pi} \oint_{C_n} \frac{dx_n}{2\pi} \psi_i(u_i) \psi(x_1) \cdots \psi(x_n) \quad (\text{C.14})$$

and contour integration (see [78] and references therein), so that the perturbed correlator becomes:

$$\begin{aligned} & \langle \psi_1(u_1) \cdots \psi_M(u_M) \varphi_1(z_1, \bar{z}_1) \cdots \varphi_N(z_N, \bar{z}_N) \rangle_{\alpha; g\varphi} = \\ & \lim_{\epsilon \rightarrow 0} \langle \varphi_1(z_1, \bar{z}_1) \cdots \varphi_N(z_N, \bar{z}_N) \rangle_{\alpha; g\varphi}^\epsilon = \\ & \sum_n \frac{g^n}{n!} \int_{\gamma_1} \cdots \int_{\gamma_n} \frac{dx_1}{2\pi} \cdots \frac{dx_n}{2\pi} \langle \psi(x_1) \cdots \psi(x_n) \tilde{\psi}_1 \cdots \tilde{\psi}_M \varphi_1 \cdots \varphi_N \rangle_\alpha, \end{aligned} \quad (\text{C.15})$$

where the C_μ are small circles surrounding the ψ_i insertion point.

Looking at (C.14) the functional form of boundary fields is left unchanged by the renormalization procedure, since integrals on the rhs of (C.14) pick up only simple poles. Thus, the fields ψ and $\tilde{\psi}$ have the same conformal dimension, and $\tilde{\psi}$ can be regarded as the ROTATION OF ψ_i GENERATED BY THE PERTURBING FIELD ψ .

Let us analyze deformations of n -point functions of the perturbing field itself. Integrals on the rhs of (C.14) vanish, since OPE (C.6) does not contain simple poles. Thus, $\tilde{\psi}$ and ψ coincide, and equation (C.6) forces all contour integrals in (C.15) to vanish if there are no bulk fields inserted on the UHP. Hence, any perturbative correction to the n -point function of ψ vanish, *i.e.* each marginal boundary operator is truly marginal to all orders of the perturbed theory.

Let us conclude this introductory section restricting our analysis to boundary condensates made up by elements of the chiral algebra, *i.e.* fields associated to elements of $\mathcal{H}^{u(1)}_1$ (see comments after formula (2.11)). These marginal fields are local with respect all bulk and boundary fields, thus formula (C.13) can be directly applied to compute deformations of arbitrary bulk and boundary fields correlators.

As remarked above, truly marginal perturbations actually generates only changes in boundary conditions. Thus, let us analyze effects of switching on a chiral boundary condensate on the gluing map Ω associate to a given BCFT. To this end, let us insert $W(z + 2i\delta) - \Omega\bar{W}(\bar{z} - 2i\delta)$, with $z = \bar{z}$, in an arbitrary unperturbed correlator of bulk fields. Then taking the $\delta \rightarrow 0^+$ limit, the chiral currents move towards the boundary and the correlator vanishes in the limiting case. To study deformations of the gluing map, let us perturb the above relation introducing operator $I_{g,J(x)} = P[e^{-S'_J}]$:

$$0 = \lim_{\delta \rightarrow 0^+} P[e^{-S'_J}] W(z + 2i\delta) - \Omega\bar{W}(\bar{z} - 2i\delta) =$$

$$\lim_{\delta \rightarrow 0^+} \sum_n \frac{g^n}{n!} \int_{\gamma_1} \cdots \int_{\gamma_n} \frac{dx_1}{2\pi} \cdots \frac{dx_n}{2\pi} J(x_1) \cdots J(x_n) W(z + 2i\delta) - \Omega\bar{W}(\bar{z} - 2i\delta)$$
(C.16)

Describing fields W and ΩW in terms of the corresponding states $|w\rangle$, $|\Omega w\rangle \in \mathcal{H}_0^{(H)}$, we get from the above integrals:

$$0 = \lim_{\delta \rightarrow 0^+} \sum_n \frac{g^n}{n!} \Phi(J_0^n |w\rangle \otimes |0\rangle; z_\delta, \bar{z}_\delta) - \Phi(|0\rangle \otimes |\Omega w\rangle; z_\delta, \bar{z}_\delta)$$

$$= e^{igJ_0} W(z) e^{-igJ_0} - \Omega\bar{W}(\bar{z})$$
(C.17)

where J_0 is the zero mode of the UHP chiral current (more details about this computation can be found in [78]). Thus, conjugation with e^{igJ_0} induces an inner automorphism on the chiral algebra $\mathcal{W}$ defined by:

$$\gamma_J(W) \doteq e^{-igJ_0} W e^{igJ_0} \quad \forall W \in \mathcal{W}$$
(C.18)

Replacing (C.18) in the second line of (C.17) we get the perturbed gluing condition under the action of a chiral marginal perturbation:

$$W(z) = \Omega \circ \gamma_J(\bar{W})(\bar{z})$$
(C.19)

Since γ_J acts trivially on the Virasoro field, the gluing condition $T = \bar{T}$ and those of all other generators which commute with J_0 remain unchanged under the chiral perturbation. Thus, these fields generate the same Ward identities as in the unperturbed theory.

Applying the construction outlined after formula (2.15), we can rephrase the result in equation (C.19) in the boundary state formalism. The chiral perturbed version of formula (2.62) turns out to be:

$$W_n = (-1)^{\bar{h}} \bar{\omega} \Omega \circ \gamma_J(\bar{W}_{-n}) \|\Omega, \alpha\|_{g,J},$$
(C.20)

where the perturbed boundary state is related to the unperturbed one via a simple rotation:

$$\|\Omega, \alpha\|_{g,J} = e^{igJ_0} \|\Omega, \alpha\|$$
(C.21)

Appendix D

Useful formulae

This section contains a collection of useful equations and formulae.

D.1 Simplicial String Duality

- **Direct products**

If a group G is the direct product of groups, $G = G_1 \times G_2$, then, given any two elements $g_1 \in G_1$ and $g_2 \in G_2$ it holds:

$$D_{m_1 n_1; m_2 n_2}^{j_1 \times j_2}(g_1 g_2) = D_{m_1 n_1}^{(1)j_1}(g_1) D_{m_2 n_2}^{(2)j_2}(g_2) \quad (D.1)$$

- **Clebsh-Gordan expansion**

The product of two Wigner functions with the same argument can be expanded in the Clebsh-Gordan series:

$$D_{m_1 n_1}^{j_1}(\Gamma) D_{m_2 n_2}^{j_2}(\Gamma) = \sum_{J=|j_1-j_2|}^{j_1+j_2} \sum_{|M|, |N| \leq J} C_{j_1 m_1 j_2 m_2}^{J M} D_M^J(\Gamma) C_{j_1 n_1 j_2 n_2}^{J N} \quad (D.2)$$

- **Unitarity relations of Clebsh-Gordan coefficients**

$$\sum_{m_1 m_2} C_{j_1 m_1 j_2 m_2}^{j m} C_{j_1 m_1 j_2 m_2}^{j' m'} = \delta_{j j'} \delta_{m m'} \quad (D.3a)$$

$$\sum_{j m} C_{j_1 m_1 j_2 m_2}^{j m} C_{j_1 m'_1 j_2 m'_2}^{j m} = \delta_{m_1 m'_1} \delta_{m_2 m'_2} \quad (D.3b)$$

D.2 Cosmological solution of supergravity

Bosonic Field Equations of type IIB supergravity

The bosonic part of the equations can be formally obtained through variation of the following action ¹:

$$\begin{aligned}
S_{IIB} = & \frac{1}{2\kappa^2} \int d^{10}x \left[-2 \sqrt{-\det g} R \right] \\
& - \frac{1}{4\kappa^2} \int d\varphi \wedge \star d\varphi + e^{-\varphi} F_{[3]}^{NS} \wedge \star F_{[3]}^{NS} + e^{2\varphi} F_{[1]}^{RR} \wedge \star F_{[1]}^{RR} \\
& + e^{\varphi} F_{[3]}^{RR} \wedge \star F_{[3]}^{RR} + \frac{1}{2} F_{[5]}^{RR} \wedge \star F_{[5]}^{RR} - C_{[4]} \wedge F_{[3]}^{NS} \wedge F_{[3]}^{RR} \quad (D.4)
\end{aligned}$$

where:

$$F_{[1]}^{RR} = dC_{[0]} \quad (D.5a)$$

$$F_{[3]}^{NS} = dB_{[2]} \quad (D.5b)$$

$$F_{[3]}^{RR} = dC_{[2]} - C_{[0]} dB_{[2]} \quad (D.5c)$$

$$F_{[5]}^{RR} = dC_{[4]} - \frac{1}{2} (B_{[2]} \wedge dC_{[2]} - C_{[2]} \wedge dB_{[2]}) \quad (D.5d)$$

It is important to stress though that the action (D.4) is to be considered only a book keeping device since the 4-form $C_{[4]}$ is not free, its field strength $F_{[5]}^{RR}$ being subject to the on-shell self-duality constraint:

$$F_{[5]}^{RR} = \star F_{[5]}^{RR} \quad (D.6)$$

From the above action the corresponding equations of motion can be obtained:

$$d \star d\varphi - e^{2\varphi} F_{[1]}^{RR} \wedge \star F_{[1]}^{RR} = -\frac{1}{2} \left(e^{-\varphi} F_{[3]}^{NS} \wedge \star F_{[3]}^{NS} - e^{\varphi} F_{[3]}^{RR} \wedge \star F_{[3]}^{RR} \right) \quad (D.7a)$$

$$d \left(e^{-\varphi} \star F_{[3]}^{NS} \right) + e^{\varphi} F_{[1]}^{RR} \wedge \star F_{[3]}^{RR} = -F_{[3]}^{RR} \wedge F_{[5]}^{RR} \quad (D.7b)$$

$$d \left(e^{2\varphi} \star F_{[1]}^{RR} \right) = -e^{\varphi} F_{[3]}^{NS} \wedge \star F_{[3]}^{RR} \quad (D.7c)$$

$$d \left(e^{\varphi} \star F_{[3]}^{RR} \right) = -F_{[5]}^{RR} \wedge F_{[3]}^{NS} \quad (D.7d)$$

$$d \star F_{[5]}^{RR} = -F_{[3]}^{NS} \wedge F_{[3]}^{RR} \quad (D.7e)$$

$$\begin{aligned}
-2 R_{MN} = & \frac{1}{2} \partial_M \varphi \partial_N \varphi + \frac{e^{2\varphi}}{2} \partial_M C_{[0]} \partial_N C_{[0]} + 150 F_{[5]M} \dots F_{[5]N}^{\dots} \\
& + 9 \left(e^{-\varphi} F_{[3]M}^{NS} F_{[3]N}^{NS} \dots + e^{\varphi} F_{[3]M}^{RR} F_{[3]N}^{RR} \dots \right) \\
& - \frac{3}{4} g_{MN} \left(e^{-\varphi} F_{[3] \dots}^{NS} F_{[3] \dots}^{NS} + e^{\varphi} F_{[3] \dots}^{RR} F_{[3] \dots}^{RR} \right) \quad (D.7f)
\end{aligned}$$

¹Note that our R is equal to $-\frac{1}{2} R^{old}$, R^{old} being the normalization of the scalar curvature usually adopted in General Relativity textbooks. The difference arises because in the traditional literature the Riemann tensor is not defined as the components of the curvature 2-form R^{ab} rather as -2 times such components.

It is not difficult to show, upon suitable identification of the massless superstring fields, that this is the correct set of equations which can be consistently obtained from the manifestly $SU(1, 1)$ covariant formulation of type IIB supergravity [107, 108, 109].

A useful integral

$$\int e^{[a x]} (\cosh[b x])^{1/4} = \frac{2^{7/4}}{4a - b} e^{[(a-b)x]} {}_2F_1\left(-\frac{1}{8} + \frac{a}{2b}, -\frac{1}{4}, \frac{7}{8} + \frac{a}{2b}; -e^{2bx}\right) \quad (\text{D.8})$$

Appendix E

Listing of the E_8 positive roots

E.1 Listing according to height

In this listing we present the roots of the E_8 Lie algebra, giving their definition both in terms of the simple roots and in the euclidian basis. The notation $a_{i|j}$ is introduced to denote the height of the root (i). The number (j) is introduced to distinguish the roots of the same height

Table E.1: Listing of positive E_8 roots according to height

label	root	Dynkin labels	Euclidian basis labels
$a_{1 1} = \alpha[1]$	$=$	$\{1,0,0,0,0,0,0,0\}$	$= \{0,1,-1,0,0,0,0,0\}$
$a_{1 2} = \alpha[2]$	$=$	$\{0,1,0,0,0,0,0,0\}$	$= \{0,0,1,-1,0,0,0,0\}$
$a_{1 3} = \alpha[3]$	$=$	$\{0,0,1,0,0,0,0,0\}$	$= \{0,0,0,1,-1,0,0,0\}$
$a_{1 4} = \alpha[4]$	$=$	$\{0,0,0,1,0,0,0,0\}$	$= \{0,0,0,0,1,-1,0,0\}$
$a_{1 5} = \alpha[5]$	$=$	$\{0,0,0,0,1,0,0,0\}$	$= \{0,0,0,0,0,1,-1,0\}$
$a_{1 6} = \alpha[6]$	$=$	$\{0,0,0,0,0,1,0,0\}$	$= \{0,0,0,0,0,1,1,0\}$
$a_{1 7} = \alpha[7]$	$=$	$\{0,0,0,0,0,0,1,0\}$	$= \{-\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$
$a_{1 8} = \alpha[8]$	$=$	$\{0,0,0,0,0,0,0,1\}$	$= \{1,-1,0,0,0,0,0,0\}$
$a_{2 1} = \alpha[9]$	$=$	$\{1,1,0,0,0,0,0,0\}$	$= \{0,1,0,-1,0,0,0,0\}$
$a_{2 2} = \alpha[10]$	$=$	$\{1,0,0,0,0,0,0,1\}$	$= \{1,0,-1,0,0,0,0,0\}$
$a_{2 3} = \alpha[11]$	$=$	$\{0,1,1,0,0,0,0,0\}$	$= \{0,0,1,0,-1,0,0,0\}$
$a_{2 4} = \alpha[12]$	$=$	$\{0,0,1,1,0,0,0,0\}$	$= \{0,0,0,1,0,-1,0,0\}$
$a_{2 5} = \alpha[13]$	$=$	$\{0,0,0,1,1,0,0,0\}$	$= \{0,0,0,0,1,0,-1,0\}$
$a_{2 6} = \alpha[14]$	$=$	$\{0,0,0,1,0,1,0,0\}$	$= \{0,0,0,0,1,0,1,0\}$
$a_{2 7} = \alpha[15]$	$=$	$\{0,0,0,0,0,1,1,0\}$	$= \{-\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\}$
Continued on next page			

Table E.1 – continued from previous page

label	root	Dynkin labels	Euclidian basis labels
$a_{3 1}$	$= \alpha[16]$	$= \{1,1,1,0,0,0,0,0\}$	$= \{0,1,0,0,-1,0,0,0\}$
$a_{3 2}$	$= \alpha[17]$	$= \{1,1,0,0,0,0,0,1\}$	$= \{1,0,0,-1,0,0,0,0\}$
$a_{3 3}$	$= \alpha[18]$	$= \{0,1,1,1,0,0,0,0\}$	$= \{0,0,1,0,0,-1,0,0\}$
$a_{3 4}$	$= \alpha[19]$	$= \{0,0,1,1,1,0,0,0\}$	$= \{0,0,0,1,0,0,-1,0\}$
$a_{3 5}$	$= \alpha[20]$	$= \{0,0,1,1,0,1,0,0\}$	$= \{0,0,0,1,0,0,1,0\}$
$a_{3 6}$	$= \alpha[21]$	$= \{0,0,0,1,0,1,1,0\}$	$= \{-\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\}$
$a_{3 7}$	$= \alpha[22]$	$= \{0,0,0,1,1,1,0,0\}$	$= \{0,0,0,0,1,1,0,0\}$
$a_{4 1}$	$= \alpha[23]$	$= \{1,1,1,1,0,0,0,0\}$	$= \{0,1,0,0,0,-1,0,0\}$
$a_{4 2}$	$= \alpha[24]$	$= \{0,1,1,1,1,0,0,0\}$	$= \{0,0,1,0,0,0,-1,0\}$
$a_{4 3}$	$= \alpha[25]$	$= \{0,1,1,1,0,1,0,0\}$	$= \{0,0,1,0,0,0,1,0\}$
$a_{4 4}$	$= \alpha[26]$	$= \{1,1,1,0,0,0,0,1\}$	$= \{1,0,0,0,-1,0,0,0\}$
$a_{4 5}$	$= \alpha[27]$	$= \{0,0,1,1,0,1,1,0\}$	$= \{-\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\}$
$a_{4 6}$	$= \alpha[28]$	$= \{0,0,1,1,1,1,0,0\}$	$= \{0,0,0,1,0,1,0,0\}$
$a_{4 7}$	$= \alpha[29]$	$= \{0,0,0,1,1,1,1,0\}$	$= \{-\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$
$a_{5 1}$	$= \alpha[30]$	$= \{1,1,1,1,1,0,0,0\}$	$= \{0,1,0,0,0,0,-1,0\}$
$a_{5 2}$	$= \alpha[31]$	$= \{1,1,1,1,0,1,0,0\}$	$= \{0,1,0,0,0,0,1,0\}$
$a_{5 3}$	$= \alpha[32]$	$= \{0,1,1,1,0,1,1,0\}$	$= \{-\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\}$
$a_{5 4}$	$= \alpha[33]$	$= \{0,1,1,1,1,1,0,0\}$	$= \{0,0,1,0,0,1,0,0\}$
$a_{5 5}$	$= \alpha[34]$	$= \{0,0,1,1,1,1,1,0\}$	$= \{-\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$
$a_{5 6}$	$= \alpha[35]$	$= \{1,1,1,1,0,0,0,1\}$	$= \{1,0,0,0,0,-1,0,0\}$
$a_{5 7}$	$= \alpha[36]$	$= \{0,0,1,2,1,1,0,0\}$	$= \{0,0,0,1,1,0,0,0\}$
$a_{6 1}$	$= \alpha[37]$	$= \{1,1,1,1,0,1,1,0\}$	$= \{-\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\}$
$a_{6 2}$	$= \alpha[38]$	$= \{1,1,1,1,1,1,0,0\}$	$= \{0,1,0,0,0,1,0,0\}$
$a_{6 3}$	$= \alpha[39]$	$= \{0,1,1,1,1,1,1,0\}$	$= \{-\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$
$a_{6 4}$	$= \alpha[40]$	$= \{0,1,1,2,1,1,0,0\}$	$= \{0,0,1,0,1,0,0,0\}$
$a_{6 5}$	$= \alpha[41]$	$= \{0,0,1,2,1,1,1,0\}$	$= \{-\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$
$a_{6 6}$	$= \alpha[42]$	$= \{1,1,1,1,1,0,0,1\}$	$= \{1,0,0,0,0,0,-1,0\}$
$a_{6 7}$	$= \alpha[43]$	$= \{1,1,1,1,0,1,0,1\}$	$= \{1,0,0,0,0,0,1,0\}$
$a_{7 1}$	$= \alpha[44]$	$= \{1,1,1,1,1,1,1,0\}$	$= \{-\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$
$a_{7 2}$	$= \alpha[45]$	$= \{1,1,1,2,1,1,0,0\}$	$= \{0,1,0,0,1,0,0,0\}$
$a_{7 3}$	$= \alpha[46]$	$= \{0,1,1,2,1,1,1,0\}$	$= \{-\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$
$a_{7 4}$	$= \alpha[47]$	$= \{0,1,2,2,1,1,0,0\}$	$= \{0,0,1,1,0,0,0,0\}$
$a_{7 5}$	$= \alpha[48]$	$= \{1,1,1,1,1,1,0,1\}$	$= \{1,0,0,0,0,1,0,0\}$
$a_{7 6}$	$= \alpha[49]$	$= \{0,0,1,2,1,2,1,0\}$	$= \{-\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$
$a_{7 7}$	$= \alpha[50]$	$= \{1,1,1,1,0,1,1,1\}$	$= \{\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\}$
$a_{8 1}$	$= \alpha[51]$	$= \{1,1,1,2,1,1,1,0\}$	$= \{-\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$
$a_{8 2}$	$= \alpha[52]$	$= \{1,1,2,2,1,1,0,0\}$	$= \{0,1,0,1,0,0,0,0\}$
$a_{8 3}$	$= \alpha[53]$	$= \{0,1,1,2,1,2,1,0\}$	$= \{-\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$

Continued on next page

Table E.1 – continued from previous page

label	root	Dynkin labels	Euclidian basis labels
$a_{8 4} = \alpha[54]$	$=$	$\{0,1,2,2,1,1,1,0\}$	$= \{-\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$
$a_{8 5} = \alpha[55]$	$=$	$\{1,1,1,2,1,1,0,1\}$	$= \{1,0,0,0,1,0,0,0\}$
$a_{8 6} = \alpha[56]$	$=$	$\{1,1,1,1,1,1,1,1\}$	$= \{\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$
$a_{9 1} = \alpha[57]$	$=$	$\{1,1,1,2,1,2,1,0\}$	$= \{-\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\}$
$a_{9 2} = \alpha[58]$	$=$	$\{1,1,2,2,1,1,1,0\}$	$= \{-\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$
$a_{9 3} = \alpha[59]$	$=$	$\{1,2,2,2,1,1,0,0\}$	$= \{0,1,1,0,0,0,0,0\}$
$a_{9 4} = \alpha[60]$	$=$	$\{0,1,2,2,1,2,1,0\}$	$= \{-\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\}$
$a_{9 5} = \alpha[61]$	$=$	$\{1,1,2,2,1,1,0,1\}$	$= \{1,0,0,1,0,0,0,0\}$
$a_{9 6} = \alpha[62]$	$=$	$\{1,1,1,2,1,1,1,1\}$	$= \{\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$
$a_{10 1} = \alpha[63]$	$=$	$\{1,1,2,2,1,2,1,0\}$	$= \{-\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\}$
$a_{10 2} = \alpha[64]$	$=$	$\{1,2,2,2,1,1,1,0\}$	$= \{-\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$
$a_{10 3} = \alpha[65]$	$=$	$\{1,2,2,2,1,1,0,1\}$	$= \{1,0,1,0,0,0,0,0\}$
$a_{10 4} = \alpha[66]$	$=$	$\{1,1,2,2,1,1,1,1\}$	$= \{\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$
$a_{10 5} = \alpha[67]$	$=$	$\{0,1,2,3,1,2,1,0\}$	$= \{-\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\}$
$a_{10 6} = \alpha[68]$	$=$	$\{1,1,1,2,1,2,1,1\}$	$= \{\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\}$
$a_{11 1} = \alpha[69]$	$=$	$\{2,2,2,2,1,1,0,1\}$	$= \{1,1,0,0,0,0,0,0\}$
$a_{11 2} = \alpha[70]$	$=$	$\{1,1,2,3,1,2,1,0\}$	$= \{-\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\}$
$a_{11 3} = \alpha[71]$	$=$	$\{1,2,2,2,1,2,1,0\}$	$= \{-\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\}$
$a_{11 4} = \alpha[72]$	$=$	$\{1,2,2,2,1,1,1,1\}$	$= \{\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$
$a_{11 5} = \alpha[73]$	$=$	$\{1,1,2,2,1,2,1,1\}$	$= \{\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\}$
$a_{11 6} = \alpha[74]$	$=$	$\{0,1,2,3,2,2,1,0\}$	$= \{-\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$
$a_{12 1} = \alpha[75]$	$=$	$\{2,2,2,2,1,1,1,1\}$	$= \{\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$
$a_{12 2} = \alpha[76]$	$=$	$\{1,1,2,3,2,2,1,0\}$	$= \{-\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$
$a_{12 3} = \alpha[77]$	$=$	$\{1,2,2,3,1,2,1,0\}$	$= \{-\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\}$
$a_{12 4} = \alpha[78]$	$=$	$\{1,2,2,2,1,2,1,1\}$	$= \{\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\}$
$a_{12 5} = \alpha[79]$	$=$	$\{1,1,2,3,1,2,1,1\}$	$= \{\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\}$
$a_{13 1} = \alpha[80]$	$=$	$\{2,2,2,2,1,2,1,1\}$	$= \{\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\}$
$a_{13 2} = \alpha[81]$	$=$	$\{1,2,2,3,2,2,1,0\}$	$= \{-\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$
$a_{13 3} = \alpha[82]$	$=$	$\{1,2,2,3,1,2,1,1\}$	$= \{\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\}$
$a_{13 4} = \alpha[83]$	$=$	$\{1,2,3,3,1,2,1,0\}$	$= \{-\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\}$
$a_{13 5} = \alpha[84]$	$=$	$\{1,1,2,3,2,2,1,1\}$	$= \{\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$
$a_{14 1} = \alpha[85]$	$=$	$\{2,2,2,3,1,2,1,1\}$	$= \{\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\}$
$a_{14 2} = \alpha[86]$	$=$	$\{1,2,2,3,2,2,1,1\}$	$= \{\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$
$a_{14 3} = \alpha[87]$	$=$	$\{1,2,3,3,2,2,1,0\}$	$= \{-\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$
$a_{14 4} = \alpha[88]$	$=$	$\{1,2,3,3,1,2,1,1\}$	$= \{\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\}$
$a_{15 1} = \alpha[89]$	$=$	$\{2,2,2,3,2,2,1,1\}$	$= \{\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$
$a_{15 2} = \alpha[90]$	$=$	$\{2,2,3,3,1,2,1,1\}$	$= \{\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\}$
$a_{15 3} = \alpha[91]$	$=$	$\{1,2,3,3,2,2,1,1\}$	$= \{\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$
Continued on next page			

Table E.1 – continued from previous page

label	root	Dynkin labels	Euclidian basis labels
$a_{15 4}$	$= \alpha[92]$	$= \{1,2,3,4,2,2,1,0\}$	$= \{-\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$
$a_{16 1}$	$= \alpha[93]$	$= \{2,2,3,3,2,2,1,1\}$	$= \{\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\}$
$a_{16 2}$	$= \alpha[94]$	$= \{2,3,3,3,1,2,1,1\}$	$= \{\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}\}$
$a_{16 3}$	$= \alpha[95]$	$= \{1,2,3,4,2,2,1,1\}$	$= \{\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$
$a_{16 4}$	$= \alpha[96]$	$= \{1,2,3,4,2,3,1,0\}$	$= \{-\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\}$
$a_{17 1}$	$= \alpha[97]$	$= \{2,2,3,4,2,2,1,1\}$	$= \{\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$
$a_{17 2}$	$= \alpha[98]$	$= \{2,3,3,3,2,2,1,1\}$	$= \{\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\}$
$a_{17 3}$	$= \alpha[99]$	$= \{1,2,3,4,2,3,1,1\}$	$= \{\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\}$
$a_{17 4}$	$= \alpha[100]$	$= \{1,2,3,4,2,3,2,0\}$	$= \{-1,0,0,0,0,0,-1\}$
$a_{18 1}$	$= \alpha[101]$	$= \{2,2,3,4,2,3,1,1\}$	$= \{\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\}$
$a_{18 2}$	$= \alpha[102]$	$= \{2,3,3,4,2,2,1,1\}$	$= \{\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$
$a_{18 3}$	$= \alpha[103]$	$= \{1,2,3,4,2,3,2,1\}$	$= \{0,-1,0,0,0,0,-1\}$
$a_{19 1}$	$= \alpha[104]$	$= \{2,2,3,4,2,3,2,1\}$	$= \{0,0,-1,0,0,0,-1\}$
$a_{19 2}$	$= \alpha[105]$	$= \{2,3,3,4,2,3,1,1\}$	$= \{\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\}$
$a_{19 3}$	$= \alpha[106]$	$= \{2,3,4,4,2,2,1,1\}$	$= \{\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$
$a_{20 1}$	$= \alpha[107]$	$= \{2,3,3,4,2,3,2,1\}$	$= \{0,0,0,-1,0,0,-1\}$
$a_{20 2}$	$= \alpha[108]$	$= \{2,3,4,4,2,3,1,1\}$	$= \{\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}\}$
$a_{21 1}$	$= \alpha[109]$	$= \{2,3,4,4,2,3,2,1\}$	$= \{0,0,0,0,-1,0,0,-1\}$
$a_{21 2}$	$= \alpha[110]$	$= \{2,3,4,5,2,3,1,1\}$	$= \{\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$
$a_{22 1}$	$= \alpha[111]$	$= \{2,3,4,5,2,3,2,1\}$	$= \{0,0,0,0,0,-1,0,-1\}$
$a_{22 2}$	$= \alpha[112]$	$= \{2,3,4,5,3,3,1,1\}$	$= \{\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$
$a_{23 1}$	$= \alpha[113]$	$= \{2,3,4,5,3,3,2,1\}$	$= \{0,0,0,0,0,0,-1,-1\}$
$a_{23 2}$	$= \alpha[114]$	$= \{2,3,4,5,2,4,2,1\}$	$= \{0,0,0,0,0,0,1,-1\}$
$a_{24 1}$	$= \alpha[115]$	$= \{2,3,4,5,3,4,2,1\}$	$= \{0,0,0,0,0,1,0,-1\}$
$a_{25 1}$	$= \alpha[116]$	$= \{2,3,4,6,3,4,2,1\}$	$= \{0,0,0,0,1,0,0,-1\}$
$a_{26 1}$	$= \alpha[117]$	$= \{2,3,5,6,3,4,2,1\}$	$= \{0,0,0,1,0,0,0,-1\}$
$a_{27 1}$	$= \alpha[118]$	$= \{2,4,5,6,3,4,2,1\}$	$= \{0,0,1,0,0,0,0,-1\}$
$a_{28 1}$	$= \alpha[119]$	$= \{3,4,5,6,3,4,2,1\}$	$= \{0,1,0,0,0,0,0,-1\}$
$a_{29 1}$	$= \alpha[120]$	$= \{3,4,5,6,3,4,2,2\}$	$= \{1,0,0,0,0,0,0,-1\}$

E.2 Listing the roots according to the dimensional filtration

Roots in D[1], D[2], D[3], D[4], D[5] and D[6]

Table E.2: Listing of positive E_8 roots according to dimensional filtration: roots in D[1], D[2], D[3], D[4], D[5] and D[6]

label	root #	Dynkin labels	Type IIB	type IIA
$d_{1 1} = \alpha[7] = \{0,0,0,0,0,0,1,0\}$			ρ	C_9
$d_{2 1} = \alpha[6] = \{0,0,0,0,0,1,0,0\}$			B_{89}	γ_8^9
$d_{2 2} = \alpha[5] = \{0,0,0,0,1,0,0,0\}$			γ_8^9	γ_8^9
$d_{2 3} = \alpha[15] = \{0,0,0,0,0,1,1,0\}$			C_{89}	C_8
$d_{3 1} = \alpha[22] = \{0,0,0,1,1,1,0,0\}$			B_{78}	γ_7^8
$d_{3 2} = \alpha[4] = \{0,0,0,1,0,0,0,0\}$			γ_7^8	γ_7^8
$d_{3 3} = \alpha[14] = \{0,0,0,1,0,1,0,0\}$			B_{79}	γ_7^9
$d_{3 4} = \alpha[13] = \{0,0,0,1,1,0,0,0\}$			γ_7^9	B_{79}
$d_{3 5} = \alpha[29] = \{0,0,0,1,1,1,1,0\}$			C_{78}	C_{789}
$d_{3 6} = \alpha[21] = \{0,0,0,1,0,1,1,0\}$			C_{79}	C_7
$d_{4 1} = \alpha[36] = \{0,0,1,2,1,1,0,0\}$			B_{67}	B_{67}
$d_{4 2} = \alpha[3] = \{0,0,1,0,0,0,0,0\}$			γ_6^7	γ_6^7
$d_{4 3} = \alpha[28] = \{0,0,1,1,1,1,0,0\}$			B_{68}	B_{68}
$d_{4 4} = \alpha[12] = \{0,0,1,1,0,0,0,0\}$			γ_6^8	γ_6^8
$d_{4 5} = \alpha[20] = \{0,0,1,1,0,1,0,0\}$			B_{69}	γ_6^9
$d_{4 6} = \alpha[19] = \{0,0,1,1,1,0,0,0\}$			γ_6^9	B_{69}
$d_{4 7} = \alpha[41] = \{0,0,1,2,1,1,1,0\}$			C_{67}	C_{679}
$d_{4 8} = \alpha[34] = \{0,0,1,1,1,1,1,0\}$			C_{68}	C_{689}
$d_{4 9} = \alpha[27] = \{0,0,1,1,0,1,1,0\}$			C_{69}	C_6
$d_{4 10} = \alpha[49] = \{0,0,1,2,1,2,1,0\}$			C_{6789}	C_{678}
$d_{5 1} = \alpha[47] = \{0,1,2,2,1,1,0,0\}$			B_{56}	B_{56}
$d_{5 2} = \alpha[2] = \{0,1,0,0,0,0,0,0\}$			γ_5^6	γ_5^6
$d_{5 3} = \alpha[40] = \{0,1,1,2,1,1,0,0\}$			B_{57}	B_{57}
$d_{5 4} = \alpha[11] = \{0,1,1,0,0,0,0,0\}$			γ_5^7	γ_5^7
$d_{5 5} = \alpha[33] = \{0,1,1,1,1,1,0,0\}$			B_{58}	B_{58}
$d_{5 6} = \alpha[18] = \{0,1,1,1,0,0,0,0\}$			γ_5^8	γ_5^8
$d_{5 7} = \alpha[25] = \{0,1,1,1,0,1,0,0\}$			B_{59}	γ_5^9
$d_{5 8} = \alpha[24] = \{0,1,1,1,1,0,0,0\}$			γ_5^9	B_{59}
$d_{5 9} = \alpha[54] = \{0,1,2,2,1,1,1,0\}$			C_{56}	C_{569}
$d_{5 10} = \alpha[46] = \{0,1,1,2,1,1,1,0\}$			C_{57}	C_{579}
$d_{5 11} = \alpha[39] = \{0,1,1,1,1,1,1,0\}$			C_{58}	C_{589}
$d_{5 12} = \alpha[32] = \{0,1,1,1,0,1,1,0\}$			C_{59}	C_5
$d_{5 13} = \alpha[53] = \{0,1,1,2,1,2,1,0\}$			C_{5789}	C_{578}
$d_{5 14} = \alpha[60] = \{0,1,2,2,1,2,1,0\}$			C_{5689}	C_{568}
$d_{5 15} = \alpha[67] = \{0,1,2,3,1,2,1,0\}$			C_{5679}	C_{567}
$d_{5 16} = \alpha[74] = \{0,1,2,3,2,2,1,0\}$			C_{5678}	$C_{34\mu}$
Continued on next page				

Table E.2 – continued from previous page

label	root #	Dynkin labels	Type IIB	type IIA
$d_{6 1}$	$= \alpha[59]$	$= \{1,2,2,2,1,1,0,0\}$	B_{45}	B_{45}
$d_{6 2}$	$= \alpha[1]$	$= \{1,0,0,0,0,0,0,0\}$	γ_4^5	γ_4^5
$d_{6 3}$	$= \alpha[52]$	$= \{1,1,2,2,1,1,0,0\}$	B_{46}	B_{46}
$d_{6 4}$	$= \alpha[9]$	$= \{1,1,0,0,0,0,0,0\}$	γ_4^6	γ_4^6
$d_{6 5}$	$= \alpha[45]$	$= \{1,1,1,2,1,1,0,0\}$	B_{47}	B_{47}
$d_{6 6}$	$= \alpha[16]$	$= \{1,1,1,0,0,0,0,0\}$	γ_4^7	γ_4^7
$d_{6 7}$	$= \alpha[38]$	$= \{1,1,1,1,1,1,0,0\}$	B_{48}	B_{48}
$d_{6 8}$	$= \alpha[23]$	$= \{1,1,1,1,0,0,0,0\}$	γ_4^8	γ_4^8
$d_{6 9}$	$= \alpha[31]$	$= \{1,1,1,1,0,1,0,0\}$	B_{49}	γ_4^9
$d_{6 10}$	$= \alpha[30]$	$= \{1,1,1,1,1,0,0,0\}$	γ_4^9	B_{49}
$d_{6 11}$	$= \alpha[100]$	$= \{1,2,3,4,2,3,2,0\}$	$B_{3\mu}$	$B_{3\mu}$
$d_{6 12}$	$= \alpha[96]$	$= \{1,2,3,4,2,3,1,0\}$	$C_{3\mu}$	$C_{39\mu}$
$d_{6 13}$	$= \alpha[64]$	$= \{1,2,2,2,1,1,1,0\}$	C_{45}	C_{459}
$d_{6 14}$	$= \alpha[58]$	$= \{1,1,2,2,1,1,1,0\}$	C_{46}	C_{469}
$d_{6 15}$	$= \alpha[51]$	$= \{1,1,1,2,1,1,1,0\}$	C_{47}	C_{479}
$d_{6 16}$	$= \alpha[44]$	$= \{1,1,1,1,1,1,1,0\}$	C_{48}	C_{489}
$d_{6 17}$	$= \alpha[37]$	$= \{1,1,1,1,0,1,1,0\}$	C_{49}	C_4
$d_{6 18}$	$= \alpha[57]$	$= \{1,1,1,2,1,2,1,0\}$	C_{4789}	C_{478}
$d_{6 19}$	$= \alpha[63]$	$= \{1,1,2,2,1,2,1,0\}$	C_{4689}	C_{468}
$d_{6 20}$	$= \alpha[70]$	$= \{1,1,2,3,1,2,1,0\}$	C_{4679}	C_{467}
$d_{6 21}$	$= \alpha[76]$	$= \{1,1,2,3,2,2,1,0\}$	C_{4678}	$C_{35\mu}$
$d_{6 22}$	$= \alpha[71]$	$= \{1,2,2,2,1,2,1,0\}$	C_{4589}	C_{458}
$d_{6 23}$	$= \alpha[77]$	$= \{1,2,2,3,1,2,1,0\}$	C_{4579}	C_{457}
$d_{6 24}$	$= \alpha[81]$	$= \{1,2,2,3,2,2,1,0\}$	C_{4578}	$C_{36\mu}$
$d_{6 25}$	$= \alpha[83]$	$= \{1,2,3,3,1,2,1,0\}$	C_{4569}	C_{456}
$d_{6 26}$	$= \alpha[87]$	$= \{1,2,3,3,2,2,1,0\}$	C_{4568}	$C_{37\mu}$
$d_{6 27}$	$= \alpha[92]$	$= \{1,2,3,4,2,2,1,0\}$	C_{4567}	$C_{38\mu}$

Roots in D[7]

- Electric with respect to the electric subgroup $SL(8) \subset E_{7(7)} \subset E_{8(8)}$

Table E.3: Roots in D[7] electric with respect to $SL(8) \subset E_{7(7)} \subset E_{8(8)}$

label	root #	Dynkin labels	q-vector	Type IIB	type IIA
$d_{7 1}$	$= \alpha[50]$	$= \{1,1,1,1,0,1,1,1\} \Rightarrow$	$\{1,1,1,1,0,1,1\}$	C_{39}	C_3
$d_{7 2}$	$= \alpha[99]$	$= \{1,2,3,4,2,3,1,1\} \Rightarrow$	$\{1,2,3,4,2,3,1\}$	$C_{4\mu}$	$C_{49\mu}$
$d_{7 3}$	$= \alpha[101]$	$= \{2,2,3,4,2,3,1,1\} \Rightarrow$	$\{2,2,3,4,2,3,1\}$	$C_{5\mu}$	$C_{59\mu}$
Continued on next page					

Table E.3 – continued from previous page

label	root #	Dynkin labels	q-vector	Type IIB	type IIA
$d_{7 4} = \alpha[105]$	$=$	$\{2,3,3,4,2,3,1,1\} \Rightarrow$	$\{2,3,3,4,2,3,1\}$	$C_{6\mu}$	$C_{69\mu}$
$d_{7 5} = \alpha[108]$	$=$	$\{2,3,4,4,2,3,1,1\} \Rightarrow$	$\{2,3,4,4,2,3,1\}$	$C_{7\mu}$	$C_{79\mu}$
$d_{7 6} = \alpha[110]$	$=$	$\{2,3,4,5,2,3,1,1\} \Rightarrow$	$\{2,3,4,5,2,3,1\}$	$C_{8\mu}$	$C_{89\mu}$
$d_{7 7} = \alpha[84]$	$=$	$\{1,1,2,3,2,2,1,1\} \Rightarrow$	$\{1,1,2,3,2,2,1\}$	C_{3678}	$C_{45\mu}$
$d_{7 8} = \alpha[86]$	$=$	$\{1,2,2,3,2,2,1,1\} \Rightarrow$	$\{1,2,2,3,2,2,1\}$	C_{3578}	$C_{46\mu}$
$d_{7 9} = \alpha[91]$	$=$	$\{1,2,3,3,2,2,1,1\} \Rightarrow$	$\{1,2,3,3,2,2,1\}$	C_{3568}	$C_{47\mu}$
$d_{7 10} = \alpha[95]$	$=$	$\{1,2,3,4,2,2,1,1\} \Rightarrow$	$\{1,2,3,4,2,2,1\}$	C_{3567}	$C_{48\mu}$
$d_{7 11} = \alpha[97]$	$=$	$\{2,2,3,4,2,2,1,1\} \Rightarrow$	$\{2,2,3,4,2,2,1\}$	C_{3467}	$C_{58\mu}$
$d_{7 12} = \alpha[102]$	$=$	$\{2,3,3,4,2,2,1,1\} \Rightarrow$	$\{2,3,3,4,2,2,1\}$	C_{3457}	$C_{68\mu}$
$d_{7 13} = \alpha[106]$	$=$	$\{2,3,4,4,2,2,1,1\} \Rightarrow$	$\{2,3,4,4,2,2,1\}$	C_{3456}	$C_{78\mu}$
$d_{7 14} = \alpha[93]$	$=$	$\{2,2,3,3,2,2,1,1\} \Rightarrow$	$\{2,2,3,3,2,2,1\}$	C_{3468}	$C_{57\mu}$
$d_{7 15} = \alpha[89]$	$=$	$\{2,2,2,3,2,2,1,1\} \Rightarrow$	$\{2,2,2,3,2,2,1\}$	C_{3478}	$C_{56\mu}$
$d_{7 16} = \alpha[98]$	$=$	$\{2,3,3,3,2,2,1,1\} \Rightarrow$	$\{2,3,3,3,2,2,1\}$	C_{3458}	$C_{67\mu}$
$d_{7 17} = \alpha[103]$	$=$	$\{1,2,3,4,2,3,2,1\} \Rightarrow$	$\{1,2,3,4,2,3,2\}$	$B_{4\mu}$	$B_{4\mu}$
$d_{7 18} = \alpha[104]$	$=$	$\{2,2,3,4,2,3,2,1\} \Rightarrow$	$\{2,2,3,4,2,3,2\}$	$B_{5\mu}$	$B_{5\mu}$
$d_{7 19} = \alpha[107]$	$=$	$\{2,3,3,4,2,3,2,1\} \Rightarrow$	$\{2,3,3,4,2,3,2\}$	$B_{6\mu}$	$B_{6\mu}$
$d_{7 20} = \alpha[109]$	$=$	$\{2,3,4,4,2,3,2,1\} \Rightarrow$	$\{2,3,4,4,2,3,2\}$	$B_{7\mu}$	$B_{7\mu}$
$d_{7 21} = \alpha[111]$	$=$	$\{2,3,4,5,2,3,2,1\} \Rightarrow$	$\{2,3,4,5,2,3,2\}$	$B_{8\mu}$	$B_{8\mu}$
$d_{7 22} = \alpha[114]$	$=$	$\{2,3,4,5,2,4,2,1\} \Rightarrow$	$\{2,3,4,5,2,4,2\}$	γ_μ^9	$B_{9\mu}$
$d_{7 23} = \alpha[8]$	$=$	$\{0,0,0,0,0,0,0,1\} \Rightarrow$	$\{0,0,0,0,0,0,0\}$	γ_3^4	γ_3^4
$d_{7 24} = \alpha[10]$	$=$	$\{1,0,0,0,0,0,0,1\} \Rightarrow$	$\{1,0,0,0,0,0,0\}$	γ_3^5	γ_3^5
$d_{7 25} = \alpha[17]$	$=$	$\{1,1,0,0,0,0,0,1\} \Rightarrow$	$\{1,1,0,0,0,0,0\}$	γ_3^6	γ_3^6
$d_{7 26} = \alpha[26]$	$=$	$\{1,1,1,0,0,0,0,1\} \Rightarrow$	$\{1,1,1,0,0,0,0\}$	γ_3^7	γ_3^7
$d_{7 27} = \alpha[35]$	$=$	$\{1,1,1,1,0,0,0,1\} \Rightarrow$	$\{1,1,1,1,0,0,0\}$	γ_3^8	γ_3^8
$d_{7 28} = \alpha[43]$	$=$	$\{1,1,1,1,0,1,0,1\} \Rightarrow$	$\{1,1,1,1,0,1,0\}$	B_{39}	γ_3^9

- Magnetic with respect to the electric subgroup $SL(8) \subset E_{7(7)} \subset E_{8(8)}$

Table E.4: Roots in $D[7]$ magnetic with respect to $SL(8) \subset E_{7(7)} \subset E_{8(8)}$

label	root #	Dynkin labels	q-vector	Type IIB	type IIA
$d_{7 29} = \alpha[112]$	$=$	$\{2,3,4,5,3,3,1,1\} \Rightarrow$	$\{2,3,4,5,3,3,1\}$	$C_{9\mu}$	C_μ
$d_{7 30} = \alpha[75]$	$=$	$\{2,2,2,2,1,1,1,1\} \Rightarrow$	$\{2,2,2,2,1,1,1\}$	C_{34}	C_{349}
$d_{7 31} = \alpha[72]$	$=$	$\{1,2,2,2,1,1,1,1\} \Rightarrow$	$\{1,2,2,2,1,1,1\}$	C_{35}	C_{359}
$d_{7 32} = \alpha[66]$	$=$	$\{1,1,2,2,1,1,1,1\} \Rightarrow$	$\{1,1,2,2,1,1,1\}$	C_{36}	C_{369}
$d_{7 33} = \alpha[62]$	$=$	$\{1,1,1,2,1,1,1,1\} \Rightarrow$	$\{1,1,1,2,1,1,1\}$	C_{37}	C_{379}
$d_{7 34} = \alpha[56]$	$=$	$\{1,1,1,1,1,1,1,1\} \Rightarrow$	$\{1,1,1,1,1,1,1\}$	C_{38}	C_{389}

Continued on next page

Table E.4 – continued from previous page

label	root #	Dynkin labels	q-vector	Type IIB	type IIA
$d_{7 35} = \alpha[94]$	$=$	$\{2,3,3,3,1,2,1,1\} \Rightarrow$	$\{2,3,3,3,1,2,1\}$	C_{3459}	C_{345}
$d_{7 36} = \alpha[90]$	$=$	$\{2,2,3,3,1,2,1,1\} \Rightarrow$	$\{2,2,3,3,1,2,1\}$	C_{3469}	C_{346}
$d_{7 37} = \alpha[85]$	$=$	$\{2,2,2,3,1,2,1,1\} \Rightarrow$	$\{2,2,2,3,1,2,1\}$	C_{3479}	C_{347}
$d_{7 38} = \alpha[80]$	$=$	$\{2,2,2,2,1,2,1,1\} \Rightarrow$	$\{2,2,2,2,1,2,1\}$	C_{3489}	C_{348}
$d_{7 39} = \alpha[78]$	$=$	$\{1,2,2,2,1,2,1,1\} \Rightarrow$	$\{1,2,2,2,1,2,1\}$	C_{3589}	C_{358}
$d_{7 40} = \alpha[73]$	$=$	$\{1,1,2,2,1,2,1,1\} \Rightarrow$	$\{1,1,2,2,1,2,1\}$	C_{3689}	C_{368}
$d_{7 41} = \alpha[68]$	$=$	$\{1,1,1,2,1,2,1,1\} \Rightarrow$	$\{1,1,1,2,1,2,1\}$	C_{3789}	C_{378}
$d_{7 42} = \alpha[82]$	$=$	$\{1,2,2,3,1,2,1,1\} \Rightarrow$	$\{1,2,2,3,1,2,1\}$	C_{3579}	C_{357}
$d_{7 43} = \alpha[88]$	$=$	$\{1,2,3,3,1,2,1,1\} \Rightarrow$	$\{1,2,3,3,1,2,1\}$	C_{3569}	C_{356}
$d_{7 44} = \alpha[79]$	$=$	$\{1,1,2,3,1,2,1,1\} \Rightarrow$	$\{1,1,2,3,1,2,1\}$	C_{3679}	C_{367}
$d_{7 45} = \alpha[69]$	$=$	$\{2,2,2,2,1,1,0,1\} \Rightarrow$	$\{2,2,2,2,1,1,0\}$	B_{34}	B_{34}
$d_{7 46} = \alpha[65]$	$=$	$\{1,2,2,2,1,1,0,1\} \Rightarrow$	$\{1,2,2,2,1,1,0\}$	B_{35}	B_{35}
$d_{7 47} = \alpha[61]$	$=$	$\{1,1,2,2,1,1,0,1\} \Rightarrow$	$\{1,1,2,2,1,1,0\}$	B_{36}	B_{36}
$d_{7 48} = \alpha[55]$	$=$	$\{1,1,1,2,1,1,0,1\} \Rightarrow$	$\{1,1,1,2,1,1,0\}$	B_{37}	B_{37}
$d_{7 49} = \alpha[48]$	$=$	$\{1,1,1,1,1,1,0,1\} \Rightarrow$	$\{1,1,1,1,1,1,0\}$	B_{38}	B_{38}
$d_{7 50} = \alpha[42]$	$=$	$\{1,1,1,1,1,0,0,1\} \Rightarrow$	$\{1,1,1,1,1,0,0\}$	γ_3^9	B_{39}
$d_{7 51} = \alpha[119]$	$=$	$\{3,4,5,6,3,4,2,1\} \Rightarrow$	$\{3,4,5,6,3,4,2\}$	γ_μ^4	γ_μ^4
$d_{7 52} = \alpha[118]$	$=$	$\{2,4,5,6,3,4,2,1\} \Rightarrow$	$\{2,4,5,6,3,4,2\}$	γ_μ^5	γ_μ^5
$d_{7 53} = \alpha[117]$	$=$	$\{2,3,5,6,3,4,2,1\} \Rightarrow$	$\{2,3,5,6,3,4,2\}$	γ_μ^6	γ_μ^6
$d_{7 54} = \alpha[116]$	$=$	$\{2,3,4,6,3,4,2,1\} \Rightarrow$	$\{2,3,4,6,3,4,2\}$	γ_μ^7	γ_μ^7
$d_{7 55} = \alpha[115]$	$=$	$\{2,3,4,5,3,4,2,1\} \Rightarrow$	$\{2,3,4,5,3,4,2\}$	γ_μ^8	γ_μ^8
$d_{7 56} = \alpha[113]$	$=$	$\{2,3,4,5,3,3,2,1\} \Rightarrow$	$\{2,3,4,5,3,3,2\}$	$B_{9\mu}$	$\gamma_{\mu 9}$

Roots in D[8]Table E.5: Listing of positive E_8 roots according to dimensional filtration: roots in D[8]

label	root #	Dynkin labels	Type IIB	type IIA
$d_{8 1} = \alpha[120]$	$=$	$\{3,4,5,6,3,4,2,2\}$	γ_μ^4	γ_μ^4

Acknowledgments

First of all I would like to thank my supervisor, Mauro Carfora, and Annalisa Marzuoli for their continuous support, help, encouragement and teachings during these three years.

I also thank my officemate, collaborator and friend, Claudio Dappiaggi, for his help, his support and for all the fruitful discussions we had these years. . . and yes, obviously also for his patience the many times I interrupted him with “Scusa, Claudio, ho una domanda da farti. . .”.

I am very grateful to Pietro Fré, Floriana Gargiulo, Ksenya Rulik, Alexander Sorin and Mario Trigiante for allowing me to work with them during my first year of Ph.D.: you taught me a lot.

I also would like to thank the CERN Theoretical Division for its kind hospitality during the past year, and all people I met there: I learnt a lot from all of you.

Especially, I thank Rodolfo Russo, for accepting to supervise my work while I was at CERN, for the fruitful collaboration started there and for the huge amount of things he taught and keep on teaching me.

I also would like to thank all people I met here in Pavia, and especially my friend Umberto, who truly shared with me this three-year experience.

Last but not least I thank Marco, obviously not only for his patience, his continuous encouragement, for the many nights he spent at my side while I was writing this thesis and for the help he gave me. Without you I would not be what I am: you are the best part of me and you will always be.

Bibliography

- [1] Adam G. Riess et al. Observational Evidence from Supernovae for an Accelerating Universe and a Cosmological Constant. *Astron. J.*, 116:1009–1038, 1998. astro-ph/9805201.
- [2] S. Perlmutter et al. Measurements of Omega and Lambda from 42 High-Redshift Supernovae. *Astrophys. J.*, 517:565–586, 1999. astro-ph/9812133.
- [3] J. L. Sievers et al. Cosmological Parameters from Cosmic Background Imager Observations and Comparisons with BOOMERANG, DASI, and MAXIMA. *Astrophys. J.*, 591:599–622, 2003. astro-ph/0205387.
- [4] Andrei D. Linde. Particle Physics and Inflationary Cosmology. 2005. hep-th/0503203.
- [5] Shamit Kachru et al. Towards inflation in string theory. *JCAP*, 0310:013, 2003. hep-th/0308055.
- [6] Shamit Kachru, Renata Kallosh, Andrei Linde, and Sandip P. Trivedi. De Sitter vacua in string theory. *Phys. Rev.*, D68:046005, 2003. hep-th/0301240.
- [7] C. P. Burgess, R. Kallosh, and F. Quevedo. de Sitter string vacua from supersymmetric D-terms. *JHEP*, 10:056, 2003. hep-th/0309187.
- [8] Pietro Fre, Mario Trigiante, and Antoine Van Proeyen. Stable de Sitter vacua from $N = 2$ supergravity. *Class. Quant. Grav.*, 19:4167–4194, 2002. hep-th/0205119.
- [9] M. de Roo, D. B. Westra, Sudhakar Panda, and M. Trigiante. Potential and mass-matrix in gauged $N = 4$ supergravity. *JHEP*, 11:022, 2003. hep-th/0310187.
- [10] Michael Gutperle and Andrew Strominger. Spacelike branes. *JHEP*, 04:018, 2002. hep-th/0202210.

- [11] V. D. Ivashchuk and V. N. Melnikov. Multidimensional classical and quantum cosmology with intersecting p-branes. *J. Math. Phys.*, 39:2866–2888, 1998. hep-th/9708157.
- [12] L. Cornalba, M. S. Costa, and C. Kounnas. A resolution of the cosmological singularity with orientifolds. *Nucl. Phys.*, B637:378–394, 2002. hep-th/0204261.
- [13] Lorenzo Cornalba and Miguel S. Costa. On the classical stability of orientifold cosmologies. *Class. Quant. Grav.*, 20:3969–3996, 2003. hep-th/0302137.
- [14] Frederic Leblond and Amanda W. Peet. A note on the singularity theorem for supergravity SD- branes. *JHEP*, 04:022, 2004. hep-th/0305059.
- [15] Martin Kruczenski, Robert C. Myers, and Amanda W. Peet. Supergravity S-branes. *JHEP*, 05:039, 2002. hep-th/0204144.
- [16] Nobuyoshi Ohta. Accelerating cosmologies from S-branes. *Phys. Rev. Lett.*, 91:061303, 2003. hep-th/0303238.
- [17] Roberto Emparan and Jaume Garriga. A note on accelerating cosmologies from compactifications and S-branes. *JHEP*, 05:028, 2003. hep-th/0304124.
- [18] Alex Buchel and Johannes Walcher. Comments on supergravity description of S-branes. *JHEP*, 05:069, 2003. hep-th/0305055.
- [19] G. Papadopoulos, J. G. Russo, and A. A. Tseytlin. Solvable model of strings in a time-dependent plane-wave background. *Class. Quant. Grav.*, 20:969–1016, 2003. hep-th/0211289.
- [20] Fernando Quevedo. Lectures on string / brane cosmology. *Class. Quant. Grav.*, 19:5721–5779, 2002. hep-th/0210292.
- [21] M. Gasperini and G. Veneziano. The pre-big bang scenario in string cosmology. *Phys. Rept.*, 373:1–212, 2003. hep-th/0207130.
- [22] Ben Craps, David Kutasov, and Govindan Rajesh. String propagation in the presence of cosmological singularities. *JHEP*, 06:053, 2002. hep-th/0205101.
- [23] Tom Banks and W. Fischler. M-theory observables for cosmological spacetimes. 2001. hep-th/0102077.
- [24] Justin Khoury, Burt A. Ovrut, Nathan Seiberg, Paul J. Steinhardt, and Neil Turok. From big crunch to big bang. *Phys. Rev.*, D65:086007, 2002. hep-th/0108187.

- [25] James E. Lidsey, David Wands, and Edmund J. Copeland. Superstring cosmology. *Phys. Rept.*, 337:343–492, 2000. hep-th/9909061.
- [26] Albion E. Lawrence and Emil J. Martinec. String field theory in curved spacetime and the resolution of spacelike singularities. *Class. Quant. Grav.*, 13:63–96, 1996. hep-th/9509149.
- [27] Ashoke Sen. Time evolution in open string theory. *JHEP*, 10:003, 2002. hep-th/0207105.
- [28] Ashoke Sen. Rolling tachyon. *JHEP*, 04:048, 2002. hep-th/0203211.
- [29] Robert Geroch. A method for generating solutions of einstein’s equations. *J. Math. Phys.*, 12:918–924, 1971.
- [30] R. Geroch. A method for generating new solutions of einstein’s equation. 2. *J. Math. Phys.*, 13:394–404, 1972.
- [31] E. Cremmer. In S. Ferrara and J.G. Taylor, editors, *Supergravity 81*, page 313, 1981.
- [32] B. L. Julia. In S. Hawking and M. Rocek, editors, *Superspace and Supergravity*. Cambridge Univ. Press, Cambridge, 1981.
- [33] B. Julia. Infinite lie algebras in physics. Invited talk given at Johns Hopkins Workshop on Current Problems in Particle Theory, Baltimore, Md., May 25-27, 1981.
- [34] V. D. Ivashchuk and V. N. Melnikov. Billiard representation for multidimensional cosmology with intersecting p-branes near the singularity. *J. Math. Phys.*, 41:6341–6363, 2000.
- [35] T. Damour, M. Henneaux, and H. Nicolai. Cosmological billiards. *Class. Quant. Grav.*, 20:R145–R200, 2003. hep-th/0212256.
- [36] Marc Henneaux and Bernard Julia. Hyperbolic billiards of pure $D = 4$ supergravities. *JHEP*, 05:047, 2003. hep-th/0304233.
- [37] Sophie de Buyl, Gaia Pinardi, and Christiane Schomblond. Einstein billiards and spatially homogeneous cosmological models. *Class. Quant. Grav.*, 20:5141–5160, 2003. hep-th/0306280.
- [38] Thibault Damour, Marc Henneaux, Alan D. Rendall, and Marsha Weaver. Kasner-like behaviour for subcritical Einstein-matter systems. *Annales Henri Poincare*, 3:1049–1111, 2002. gr-qc/0202069.
- [39] T. Damour and M. Henneaux. Chaos in superstring cosmology. *Gen. Rel. Grav.*, 32:2339–2343, 2000.

- [40] Thibault Damour, Marc Henneaux, Bernard Julia, and Hermann Nicolai. Hyperbolic Kac-Moody algebras and chaos in Kaluza-Klein models. *Phys. Lett.*, B509:323–330, 2001. hep-th/0103094.
- [41] Thibault Damour and Marc Henneaux. E(10), BE(10) and arithmetical chaos in superstring cosmology. *Phys. Rev. Lett.*, 86:4749–4752, 2001. hep-th/0012172.
- [42] Thibault Damour and Marc Henneaux. Oscillatory behaviour in homogeneous string cosmology models. *Phys. Lett.*, B488:108–116, 2000. hep-th/0006171.
- [43] Thibault Damour and Marc Henneaux. Chaos in superstring cosmology. *Phys. Rev. Lett.*, 85:920–923, 2000. hep-th/0003139.
- [44] J. Demaret, Y. De Rop, and M. Henneaux. CHAOS IN NONDIAGONAL SPATIALLY HOMOGENEOUS COSMOLOGICAL MODELS IN SPACE-TIME DIMENSIONS ≤ 10 . *Phys. Lett.*, B211:37–41, 1988.
- [45] Neil Marcus and John H. Schwarz. Three-dimensional supergravity theories. *Nucl. Phys.*, B228:145, 1983.
- [46] H. Nicolai and H. Samtleben. Compact and noncompact gauged maximal supergravities in three dimensions. *JHEP*, 04:022, 2001. hep-th/0103032.
- [47] L. Andrianopoli, R. D’Auria, S. Ferrara, P. Fre, and M. Trigiante. R-R scalars, U-duality and solvable Lie algebras. *Nucl. Phys.*, B496:617–629, 1997. hep-th/9611014.
- [48] Laura Andrianopoli et al. Solvable Lie algebras in type IIA, type IIB and M theories. *Nucl. Phys.*, B493:249–280, 1997. hep-th/9612202.
- [49] P. Fre et al. Cosmological backgrounds of superstring theory and solvable algebras: Oxidation and branes. *Nucl. Phys.*, B685:3–64, 2004.
- [50] Gerard ’t Hooft. A lanar diagram theory for trong interactions. *Nucl. Phys.*, B72:461, 1974.
- [51] Leonard Susskind. The world as a hologram. *J. Math. Phys.*, 36:6377–6396, 1995.
- [52] Gerard ’t Hooft. Dimensional reduction in quantum gravity. 1993.
- [53] Juan M. Maldacena. The large n limit of superconformal field theories and supergravity. *Adv. Theor. Math. Phys.*, 2:231–252, 1998.
- [54] Marcos Marino. Les houches lectures on matrix models and topological strings. 2004.

- [55] Antonella Grassi and Michele Rossi. Large n dualities and transitions in geometry. 2002.
- [56] Rajesh Gopakumar. From free fields to AdS. *Phys. Rev.*, D70:025009, 2004.
- [57] Rajesh Gopakumar. From free fields to AdS. II. *Phys. Rev.*, D70:025010, 2004. hep-th/0402063.
- [58] Edward Witten. Anti-de Sitter space and holography. *Adv. Theor. Math. Phys.*, 2:253–291, 1998. hep-th/9802150.
- [59] S. S. Gubser, Igor R. Klebanov, and Alexander M. Polyakov. Gauge theory correlators from non-critical string theory. *Phys. Lett.*, B428:105–114, 1998. hep-th/9802109.
- [60] M. Carfora, C. Dappiaggi, and A. Marzuoli. The modular geometry of random Regge triangulations. *Class. Quant. Grav.*, 19:5195, 2002. gr-qc/0206077.
- [61] Mauro Carfora and Annalisa Marzuoli. Conformal modes in simplicial quantum gravity and the Weil-Petersson volume of moduli space. *Adv. Theor. Math. Phys.*, 6:357–401, 2003. [math-ph/0107028].
- [62] M. Mulase and M. Penkava. Ribbon graphs, quadratic differentials on riemann surfaces and algebraic curves defined over $\overline{\mathbb{Q}}$. math-ph/9811024.
- [63] Rajesh Gopakumar. From free fields to AdS. III. 2005. hep-th/0504229.
- [64] Mauro Carfora, Claudio Dappiaggi, and Valeria Gili. Simplicial aspects of string dualities. *AIP Conf. Proc.*, 751:182–184, 2005. hep-th/0410006.
- [65] M. Carfora, C. Dappiaggi, and V. Gili. Simplicial and modular aspects of string dualities. In preparation.
- [66] Robert P. Langlands. An essay on the dynamics and statistics of critical field theories. In *Canadian Mathematical Society, Selecta 1945-1995*, volume 3, pages 173–209.
- [67] G. Arcioni, M. Carfora, C. Dappiaggi, and A. Marzuoli. The WZW model on random Regge triangulations. *Jour. Geom. Phys.*, 52:137, 2004. hep-th/0209031.
- [68] M. Nakahara. *Geometry, Topology and Physics*. Graduate Student Series in Physics. Institute of Physics Publishing, 1990.
- [69] Jan Ambjorn, Bergfinnur Durhuus, and Thordur Jonsson. *Quantum Geometry: A Statistical Field Theory Approach*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, 1997.

- [70] M Troyanov. Prescribing curvature on compact surfaces with conical singularities. *Trans. Amer. Math. Soc.*, (324):793, 1991.
- [71] W.P. Thurston. Shapes of polyhedra and tringulations of the sphere. In *Geometry end Topology Monographs. Volume 1: The Epstein Birthday Schrift.*, pages 511–549. 1988.
- [72] E. Looijenga. Intersection theory on deligne-mumford compactifications, 1992-93. Séminaire Bourbaki.
- [73] David C. Lewellen. Sewing constraints for conformal field theories on surfaces with boundaries. *Nucl. Phys.*, B372:654–682, 1992.
- [74] John L. Cardy and David C. Lewellen. Bulk and boundary operators in conformal field theory. *Phys. Lett.*, B259:274–278, 1991.
- [75] M. R. Gaberdiel. D-branes from conformal field theory. *Fortsch. Phys.*, 50:783–801, 2002.
- [76] E. Frenkel and D. Ben-Zvi. Vertex algebras and algebraic curves. Providence, USA: Am. Math. Soc. (2004) 400 p.
- [77] Mitsuhiro Kato and Tomoharu Okada. D-branes on group manifolds. *Nucl. Phys.*, B499:583–595, 1997.
- [78] Andreas Recknagel and Volker Schomerus. Boundary deformation theory and moduli spaces of d-branes. *Nucl. Phys.*, B545:233–282, 1999.
- [79] John L. Cardy. Boundary conditions, fusion rules and the verlinde formula. *Nucl. Phys.*, B324:581, 1989.
- [80] P. Di Francesco, P. Matheu, and D. Sénéchal. *Conformal Field Theory*. Springer-Verlag, New York Inc., 1996.
- [81] H. Saleur. Lectures on non perturbative field theory and quantum impurity problems. 1998.
- [82] Eric Charpentier and Krzysztof Gawedzki. Wess-Zumino-Witten conformal field theory for simply laced groups at level one. *Ann. Phys.*, 213:233–294, 1992.
- [83] Roger E. Behrend, Paul A. Pearce, Valentina B. Petkova, and Jean-Bernard Zuber. Boundary conditions in rational conformal field theories. *Nucl. Phys.*, B570:525–589, 2000.
- [84] Nobuyuki Ishibashi. The boundary and crosscap states in conformal field theories. *Mod. Phys. Lett.*, A4:251, 1989.

- [85] V. G. Kac. Contravariant form for infinite dimensional Lie algebras and superalgebras. (Talk). In *Austin 1978, Proceedings, Group Theoretical Methods In Physics*, Berlin 1979, 441-445.
- [86] Igor R. Klebanov and Alexander M. Polyakov. Interaction of discrete states in two-dimensional string theory. *Mod. Phys. Lett.*, A6:3273–3281, 1991.
- [87] Jr. Callan, Curtis G., Igor R. Klebanov, Andreas W. W. Ludwig, and Juan M. Maldacena. Exact solution of a boundary conformal field theory. *Nucl. Phys.*, B422:417–448, 1994.
- [88] Kristjan R. Kristjansson and Larus Thorlacius. Correlation functions in a $c = 1$ boundary conformal field theory. *JHEP*, 01:047, 2005.
- [89] Jr. Callan, Curtis G. and Igor R. Klebanov. Exact $c = 1$ boundary conformal field theories. *Phys. Rev. Lett.*, 72:1968–1971, 1994.
- [90] Michael B. Green and Michael Gutperle. Symmetry breaking at enhanced symmetry points. *Nucl. Phys.*, B460:77–108, 1996.
- [91] M. R. Gaberdiel, A. Recknagel, and G. M. T. Watts. The conformal boundary states for $su(2)$ at level 1. *Nucl. Phys.*, B626:344–362, 2002.
- [92] Ingo Runkel. Boundary structure constants for the a -series virasoro minimal models. *Nucl. Phys.*, B549:563–578, 1999.
- [93] Giovanni Felder, Jurg Frohlich, Jurgen Fuchs, and Christoph Schweigert. The geometry of WZW branes. *J. Geom. Phys.*, 34:162–190, 2000.
- [94] Luis Alvarez-Gaume, C. Gomez, and G. Sierra. Quantum group interpretation of some conformal field theories. *Phys. Lett.*, B220:142, 1989.
- [95] Clifford v. Johnson. *D-branes*. Cambridge Monographs on Mathematical Physics, 2003.
- [96] Amit Giveon, Massimo Porrati, and Eliezer Rabinovici. Target space duality in string theory. *Phys. Rept.*, 244:77–202, 1994.
- [97] J. E. Paton and Hong-Mo Chan. Generalized veneziano model with isospin. *Nucl. Phys.*, B10:516–520, 1969.
- [98] Jr. Callan, Curtis G., C. Lovelace, C. R. Nappi, and S. A. Yost. Loop corrections to Superstring Equations of Motion. *Nucl. Phys.*, B308:221, 1988.
- [99] Jr. Callan, Curtis G. and Igor R. Klebanov. D-brane boundary state dynamics. *Nucl. Phys.*, B465:473–486, 1996.

- [100] Ali Yegulalp. New boundary conformal field theories indexed by the simply laced Lie algebras. *Nucl. Phys.*, B450:641–662, 1995.
- [101] Paul H. Ginsparg and Gregory W. Moore. Lectures on 2-d gravity and 2-d string theory. 1993.
- [102] P. Di Francesco, Paul H. Ginsparg, and Jean Zinn-Justin. 2-d gravity and random matrices. *Phys. Rept.*, 254:1–133, 1995.
- [103] D. V. Alekseevskii. Classification of quaternionic spaces with a transitive solvable group of motion. *Math. USSR Izvestija*, 9:297–339, 1975.
- [104] Arjan Keurentjes. The group theory of oxidation. *Nucl. Phys.*, B658:303–347, 2003. hep-th/0210178.
- [105] B. Julia. Group disintegrations. Invited paper presented at Nuffield Gravity Workshop, Cambridge, Eng., Jun 22 - Jul 12, 1980.
- [106] I. C. G. Campbell and Peter C. West. N=2 D = 10 Nonchiral Supergravity and its spontaneous compactification. *Nucl. Phys.*, B243:112, 1984.
- [107] John H. Schwarz. Covariant field equations of chiral N=2 D = 10 supergravity. *Nucl. Phys.*, B226:269, 1983.
- [108] Paul S. Howe and Peter C. West. The complete n=2, d = 10 supergravity. *Nucl. Phys.*, B238:181, 1984.
- [109] L. Castellani and I. Pesando. The Complete superspace action of chiral D = 10, N=2 supergravity. *Int. J. Mod. Phys.*, A8:1125–1138, 1993.
- [110] E. Cremmer, B. Julia, Hong Lu, and C. N. Pope. Dualisation of dualities. I. *Nucl. Phys.*, B523:73–144, 1998. hep-th/9710119.
- [111] Hong Lu, C. N. Pope, and K. S. Stelle. Weyl Group Invariance and p-brane Multiplets. *Nucl. Phys.*, B476:89–117, 1996. hep-th/9602140.
- [112] Matteo Bertolini and Mario Trigiante. Regular R-R and NS-NS BPS black holes. *Int. J. Mod. Phys.*, A15:5017, 2000.
- [113] Hong Lu, S. Mukherji, C. N. Pope, and K. W. Xu. Cosmological solutions in string theories. *Phys. Rev.*, D55:7926–7935, 1997. hep-th/9610107.
- [114] Hong Lu, S. Mukherji, and C. N. Pope. From p-branes to cosmology. *Int. J. Mod. Phys.*, A14:4121–4142, 1999. hep-th/9612224.
- [115] Andre Lukas, Burt A. Ovrut, and Daniel Waldram. Cosmological solutions of type II string theory. *Phys. Lett.*, B393:65–71, 1997. hep-th/9608195.

- [116] Andre Lukas, Burt A. Ovrut, and Daniel Waldram. String and M-theory cosmological solutions with Ramond forms. *Nucl. Phys.*, B495:365–399, 1997. hep-th/9610238.
- [117] C. Grojean, F. Quevedo, G. Tasinato, and I. Zavala. Branes on charged dilatonic backgrounds: Self-tuning, Lorentz violations and cosmology. *JHEP*, 08:005, 2001. hep-th/0106120.
- [118] P. Fre, K. Rulik, and M. Trigiante. Exact solutions for bianchi type cosmological metrics, weyl orbits of $e(8(8))$ subalgebras and p-branes. *Nucl. Phys.*, B694:239–274, 2004. hep-th/0312189.
- [119] Pietro Fre', Floriana Gargiulo, Ksenya Rulik, and Mario Trigiante. The general pattern of kac moody extensions in supergravity and the issue of cosmic billiards. 2005. hep-th/0507249.
- [120] H. Nicolai. A hyperbolic lie algebra from supergravity. *Phys. Lett.*, B276:333–340, 1992.